

September 1, 2009

UNCONSTRAINED HIGHER SPINS OF MIXED SYMMETRY

II. FERMION FIELDS

A. CAMPOLEONI ^{a,1}, D. FRANCIA ^b, J. MOURAD ^b AND A. SAGNOTTI ^a*^aScuola Normale Superiore and INFN
Piazza dei Cavalieri, 7
I-56126 Pisa ITALY**e-mail: a.campoleoni@sns.it, sagnotti@sns.it**^bAstroParticule et Cosmologie (APC) ²
Université Paris VII - Campus Paris Rive Gauche
10, rue Alice Domon et Leonie Duquet
F-75205 Paris Cedex 13 FRANCE**e-mail: francia@apc.univ-paris7.fr, mourad@apc.univ-paris7.fr*

ABSTRACT

This paper is a sequel of arXiv:0810.4350 [hep-th], and is also devoted to the local “metric-like” unconstrained Lagrangians and field equations for higher-spin fields of mixed symmetry in flat space. Here we complete the previous constrained on-shell formulation of Labastida for Fermi fields, deriving the corresponding constrained Lagrangians both via the Bianchi identities and via the requirement of self-adjointness. We also describe two types of unconstrained Lagrangian formulations: a “minimal” one, containing higher derivatives of the compensator fields, and another non-minimal one, containing only one-derivative terms. We identify classes of these systems that are invariant under Weyl-like symmetry transformations.

¹Address after October 1, 2009: Max-Planck-Institut für Gravitationsphysik, Albert-Einstein-Institut, Am Mühlenberg 1, DE-14476 Golm, Germany.

²Unité mixte de Recherche du CNRS (UMR 7164.)

Contents

1	INTRODUCTION	3
2	TWO-FAMILY FERMIONIC FIELDS	6
2.1	THE LAGRANGIANS	6
2.2	THE FIELD EQUATIONS	20
2.3	ON-SHELL REDUCTION TO $\mathcal{S} = 0$	23
2.3.1	WEYL-LIKE SYMMETRIES	30
2.3.2	LAGRANGIANS THAT ARE TOTAL DERIVATIVES	37
2.4	AN EXAMPLE: REDUCIBLE SPINOR-TENSOR FIELDS OF RANK $(3, 2)$	38
2.4.1	LAGRANGIAN AND FIELD EQUATIONS	38
2.4.2	ON-SHELL REDUCTION	41
2.4.3	CURRENT EXCHANGES	44
3	GENERAL FERMIONIC FIELDS	47
3.1	THE CONSTRAINED LAGRANGIANS	47
3.2	THE UNCONSTRAINED LAGRANGIANS	53
3.3	THE FIELD EQUATIONS	58
3.4	COMMENTS ON THE ON-SHELL REDUCTION TO $\mathcal{S} = 0$	59
3.4.1	THE DEGREES OF FREEDOM DESCRIBED BY $\mathcal{S} = 0$	60
3.4.2	WEYL-LIKE SYMMETRIES	62
4	IRREDUCIBLE FIELDS OF MIXED SYMMETRY	66
4.1	AN APPLICATION: SOME LAGRANGIANS WITH WEYL-LIKE SYMMETRIES	70
5	MULTI-FORM SPINOR-TENSORS	72
6	ELIMINATION OF HIGHER DERIVATIVES	76
6.1	SYMMETRIC FERMIONS	77
6.2	MIXED-SYMMETRY FERMIONS	80
7	CONCLUSIONS	83
A	NOTATION AND CONVENTIONS	86
B	USEFUL IDENTITIES FOR FERMIONS	90
C	PROOF OF SOME RESULTS USED IN SECTION 3	93
D	SOME PROPERTIES OF THE REDUCTION FOR N -FAMILY FIELDS	101

1 INTRODUCTION

This paper is devoted to fermionic higher-spin fields [1] of mixed symmetry in flat space, and thus complements the corresponding treatment for bosonic fields presented in [2]. The general framework and the motivations for the present analysis were already illustrated in detail in [2] and will not be repeated here, where we shall confine ourselves to some additional considerations.

Actions for generic massless symmetric spinor-tensors were first constructed in the late seventies by Fang and Fronsdal [3] starting from the massive Singh-Hagen [4] construction. They were thus able to display the gauge symmetry underlying these massless Lagrangians, while also inheriting from the massive theory some surprising algebraic constraints on gauge parameters and gauge fields: in their setup the former are in fact to be γ -traceless, while the latter are to be triply γ -traceless. Other independent justifications for the constraints were provided in [5, 6] shortly thereafter.

In the eighties, crucial inputs originating from the development of String Field Theory [7] shifted the attention to tensors of mixed symmetry. Important progress was then made by a number of authors [8, 9] (with some relevant work actually dating back to previous years [10]), and most notably Labastida [11, 12] constructed bosonic Lagrangians extending what Fronsdal had attained for symmetric tensors in [13]. Those constrained Lagrangians were recently taken as a starting point for a theory of unconstrained mixed-symmetry bosons presented in [2]. Indeed, as shown in a number of previous works [14, 15, 16, 17, 18, 19, 20, 21], the need for algebraic constraints on gauge fields and parameters can be naturally foregone, either via the addition of non-local terms or via the introduction of auxiliary fields. While this could well help to simplify the structure of the interactions, one clear motivation to delve more deeply into these matters is provided by String Field Theory [7], whose equations of motion, in their simplest form and in the tensionless limit, describe reducible chains of massless higher-spin modes, without any algebraic constraints on gauge fields and gauge parameters [22]. In addition, in the free limit a geometrical formulation of interacting higher-spin gauge fields along the lines of Einstein's gravity³ would be expected to rely somehow on the linearized higher-spin curvatures introduced by de Wit and Freedman for the symmetric case [6], or on their generalizations for mixed-symmetry fields discussed in [18]. Curvature tensors are defined in terms of unconstrained gauge fields, while their properties rely on gauge transformations involving unconstrained gauge parameters, which calls for a formulation of the free dynamics that is capable of avoiding from the outset the algebraic restrictions of the Fang-Fronsdal-Labastida construction.

As we anticipated, both the Fronsdal constraints and their generalizations can be bypassed in two quite distinct, albeit related, ways. The first rests directly on the higher-spin curvatures, and was indeed shown to be possible in [15, 19] for symmetric tensors, at the price of allowing *non-local* contributions to the Lagrangians. Investigations in the same spirit were then performed for bosons of mixed symmetry in [18], and more recently for massive symmetric Bose and Fermi fields in [21], leading in particular to a formulation of massive higher spins in which, in sharp contrast with the original Singh-Hagen setting [4], *no* auxiliary fields are needed. The second way rests on the introduction of additional fields in more conventional local Lagrangian formulations [14, 17, 19]. In its “minimal form” [16, 17, 19], Fronsdal's constraints can then be foregone at the price of introducing only two additional fields and some higher-derivative terms that are anyway

³Both the present work and its companion paper [2] proceed along the lines of the metric formalism for gravity. An important alternative is the higher-spin counterpart of the frame-like formalism [23, 24], that lies at the heart of the Vasiliev setting, where the interactions are driven by an infinite-dimensional extension of the tangent-space Lorentz algebra [25, 26].

pure gauge. In particular, unconstrained Lagrangians for symmetric Fermi fields of spin $s + \frac{1}{2}$ were obtained ⁴ in [17]. Together with the gauge potential $\psi_{\mu_1 \dots \mu_s}$, they involve an additional spin- $(s-2)$ compensator $\xi_{\mu_1 \dots \mu_{s-2}}$, that first emerges for spin $s = \frac{5}{2}$, and a spin- $(s-3)$ Lagrange multiplier, that first emerges for spin $s = \frac{7}{2}$.

Even at the constrained level, the theory of mixed-symmetry fermionic fields $\psi_{\mu_1 \dots \mu_{s_1}, \nu_1 \dots \nu_{s_2}, \dots}$, bearing several “families” of fully symmetric indices, was left somewhat incomplete in the metric-like approach, since Labastida [30] did not arrive at Lagrangians generalizing the results of Fang and Fronsdal ⁵. He nonetheless proposed free fermionic equations of motion and identified the proper generalizations of the constraints on parameters and gauge fields, that in our notation read

$$\gamma_{(i} \epsilon_{j)} = 0, \quad T_{(ij} \gamma_{k)} \psi = 0. \quad (1.1)$$

Here γ_i denotes a γ -trace in the i -th index family, T_{ij} denotes a trace involving a pair of space-time indices belonging, respectively, to the i -th and j -th families, and the parentheses indicate, here and elsewhere in this paper, complete symmetrization of the enclosed indices, with the minimal number of terms needed to this end and with *unit* overall normalization. Hence, quite differently from what the symmetric case of [3] could naively suggest, but in strict analogy with the bosonic construction of [11, 12], *not all* γ -traces of the gauge parameters and *not all* triple γ -traces of the gauge fields are forced to vanish in Labastida’s formulation.

In this paper we present three main groups of results :

- we extend to the Lagrangian level the field equations proposed by Labastida for Fermi fields of mixed symmetry in D dimensions (see eqs. (3.7) and (3.28)). In Section 3.4.1 we also show that their propagating degrees of freedom fill representations of the little group $O(D-2)$;
- we propose a number of alternative formulations in which all constraints are removed at the price of introducing “minimal” sets of additional auxiliary fields, in the spirit of what was already done for bosons in [2] (see for instance eqs. (3.42) and (6.31));
- we show that the resulting Lagrangians can display Weyl-like symmetries for particular models in special space-time dimensions. In particular, in Section 2.3 we provide a full classification of Weyl-invariant Lagrangians for two-family Fermi fields, while in Sections 3.4 and 4.1 we set up the corresponding analysis for general N -family fields and display some interesting classes of examples.

For pedagogical reasons, in Section 2 we open our discussion constructing Lagrangians and field equations for the simplest generalization of the fully symmetric case, two-family fermionic gauge fields of the type $\psi_{\mu_1 \dots \mu_{s_1}, \nu_1 \dots \nu_{s_2}}$, and study their on-shell reduction to the constrained Labastida form of [30]. As in [2], we discuss in detail the emergence of Weyl-like symmetries and

⁴Previous investigations in this direction [14], based on BRST techniques [27], led to constructions involving $\mathcal{O}(s)$ different fields to describe unconstrained spin- s modes. The reduction of the corresponding equations to the minimal ones involving the compensators was first presented in [16]. See also [28] for an early discussion of the spin 3 compensator. More recently, an alternative formulation without higher-derivative terms was obtained in [20] for symmetric bosons and fermions by suitably restricting the “triplets” [22] of String Field Theory. See also [29] for a discussion of the triplets in the frame-like formalism.

⁵Actually, fermionic Lagrangians were first built in [9], within a BRST-like formalism along the lines of the first attempts to construct free covariant string equations, in an approach that is rather remote from this line of developments.

their role in the reduction procedure of the field equations to the non-Lagrangian Labastida form. We conclude Section 2 with the explicit example of reducible rank-(3, 2) spinor-tensor fields, that are relatively simple but suffice to illustrate a number of key points of our construction ⁶. In Section 3 we extend the discussion to multi-family fermionic fields, identifying via the Bianchi identities their unconstrained Lagrangians. When restricted to the case of constrained fields, our results determine the off-shell Lagrangian extension of the field equations of [30]. However, we also describe how, in the constrained case, requiring that the kinetic operator be self-adjoint, along the lines of what was previously done for bosons in [12], one can recover the same result. Moreover, we show that the non-Lagrangian Labastida equation of [30] propagates in general the correct degrees of freedom and comment on the degenerate low-dimensional cases in which the Lagrangian equations do not reduce directly to it and Weyl-like symmetries emerge.

The key ingredients of our construction are *reducible* $gl(D)$ spinor-tensors $\psi_{\mu_1 \dots \mu_{s_1}, \nu_1 \dots \nu_{s_2}, \dots}$, possessing several “families” of *symmetric* index sets, in a D -dimensional space time. Spinor-tensor fields of this kind describe a portion of the massive fermionic excitations of the five ten-dimensional superstrings, of their type-0 counterparts [31] and of various brane systems [32]. However, in Section 4 we also describe how to adapt the results of this paper to spinor-tensor fields transforming in *irreducible* representations of the Lorentz group, so as to clarify how the theory develops along lines that are more closely related to what is usually done for low spins. Both the Fang-Fronsdal-Labastida kinetic operators and the corresponding Lagrangians take the same form for reducible and for Young-projected irreducible fields, but when considering the latter the theory involves fewer independent gauge parameters and fewer independent auxiliary fields, in a way that we spell out in detail. We also display an instructive off-shell counting argument that illustrates for irreducible two-family fields how the Labastida constraints of [30] result in the propagation of irreducible representations of the Poincaré group. In Section 4.1 we show that combining the conditions for irreducibility with the constraints selecting cases with Weyl-like symmetries allows one to find explicit solutions of the latter for an interesting class of mixed-symmetry fields.

In Section 5 we adapt our formalism, following largely previous results of [18], to another relevant class of fields, multi-form spinor-tensors possessing several “families” of *antisymmetric* index sets in a D -dimensional space time, while in Section 6 we discuss how the minimal higher-derivative unconstrained Lagrangians of Sections 2 and 3 can be reduced systematically to others with a single derivative, generalizing the results of [21, 2]. Section 7 contains our conclusions, and the paper closes with a number of appendices where our notation is carefully spelled out and tedious intermediate derivations are elaborated upon.

Both here and in [2] our notation is a natural extension of the index-free description used in previous work on symmetric (spinor-) tensors by some of us [15, 16, 17, 19, 21]. It involves explicit “family” indices, that are needed to identify particular subsets of the actual tensor indices to be treated differently by the various operators entering field equations and Lagrangians. For instance, while the basic fermionic field can still be simply denoted by ψ , as in the fully symmetric case, its gauge transformation,

$$\delta \psi = \partial^i \epsilon_i, \quad (1.2)$$

now involves explicit family indices. These identify the various gauge parameters and the various sets of tensor indices taking part in the gradient, which are to be fully symmetrized (or antisymmetrized) *only* with others of the same family. At the same time, the gauge parameters

⁶In contrast with the notation used for instance in [19], in this paper the rank refers only to the vector indices carried by spinor-tensor fields.

ϵ_i also carry one lower family index, and bear correspondingly one less space-time index of the i -th family than the gauge field ψ .

From the methodological viewpoint, our approach differs somewhat from the original paper [12], since we are guided throughout by the Bianchi identities and their γ -traces. This choice was instrumental for deriving, in a relatively handy fashion, complete local Lagrangians for Fermi fields of mixed symmetry and provides a neat rationale for these rather complicated constructions.

This paper is admittedly rather long. However, Section 2 can already provide a fair view of our methods and of the subtleties to be met in the general case, both in its early general portion and in the explicit examples of Section 2.4. On the other hand, Section 3 contains our main results (eqs. (3.7) and (3.28)), but the derivation of the Lagrangians is somewhat involved and may be skipped in a first reading. Sections 4 and 5, devoted to irreducible fields and to multi-forms, should be also accessible to a large extent after reading Section 2. Finally, a glance at the first part of Section 6 can provide a fair view of the key ideas behind the formalism that we are proposing to reduce the unconstrained models to more conventional one-derivative Lagrangians.

2 TWO-FAMILY FERMIONIC FIELDS

The “metric-like” theory of fermionic fields of mixed symmetry is less developed than its bosonic counterpart, since Labastida [30] did identify both their field equations and the algebraic constraints on gauge fields and parameters but did not arrive at the corresponding Lagrangians. One of the aims of the present paper is therefore to build constrained and unconstrained metric-like Lagrangian formulations for Fermi fields of mixed symmetry in flat space. We begin, as we did for Bose fields in [2], with a discussion of the relatively handy two-family fields, in this case spinor-tensors of the type $\psi_{\mu_1 \dots \mu_{s_1}, \nu_1 \dots \nu_{s_2}}$, that can already exhibit the key features of our construction.

2.1 THE LAGRANGIANS

In the following all vector and spinor indices will be left implicit, as in [15, 16, 17, 19, 21, 2], so that a generic spinor-tensor of the type $\psi_{\mu_1 \dots \mu_{s_1}, \nu_1 \dots \nu_{s_2}}$ will be simply denoted by ψ . However, operators like gradients, divergences and (γ -)traces will bear “family” indices specifying the sets of space-time indices they are acting upon. In particular, divergences and gradients will be denoted exactly as in [2], so that, for instance,

$$\begin{aligned} \partial_1 \psi &\equiv \partial_\lambda \psi^\lambda_{\mu_1 \dots \mu_{s_1-1}, \nu_1 \dots \nu_{s_2}} , \\ \partial^1 \psi &\equiv \partial_{(\mu_1} \psi_{\mu_2 \dots \mu_{s_1+1}), \nu_1 \dots \nu_{s_2}} . \end{aligned} \tag{2.1}$$

As in [2] and as we shall explain in detail in Appendix A, an upper family index, as the one borne by the gradient above, will signal the addition of a space-time index to the corresponding group or family, while a lower family index, as the one borne by the divergence above, will signal its removal. In a similar spirit, even when dealing with Fermi fields it is convenient to introduce traces and metric tensors, carrying respectively a pair of lower or upper family indices, so that

for instance ⁷

$$\begin{aligned} T_{12} \psi &\equiv \psi^{\lambda}_{\mu_1 \dots \mu_{s_1-1}, \lambda \nu_1 \dots \nu_{s_2-1}} , \\ \eta^{12} \psi &\equiv \frac{1}{2} \sum_{n=1}^{s_2+1} \eta_{\mu_n}(\nu_1 | \psi_{\dots \mu_r \neq n \dots, | \nu_2 \dots \nu_{s_2+1}}) . \end{aligned} \quad (2.2)$$

We should stress again that in this paper groups of (space-time or family) indices surrounded by parentheses are meant to be symmetrized via the minimal number of terms needed to this end, and to be normalized with unit overall coefficient, rather than with unit strength. Aside from these building blocks, fermionic Lagrangians involve two new types of operators, both related to the space-time γ -matrix and bearing, respectively, one lower and one upper family index. The former, γ_i , removes one Lorentz index belonging to the i -th family via a γ -trace, while the latter, γ^i , adds an ordinary γ -matrix bearing a Lorentz index of the i -th family, to be properly symmetrized with all its peers. Thus, for instance, in the two-family case one is also led to consider expressions of the type

$$\begin{aligned} \gamma_1 \psi &\equiv \gamma_{\lambda} \psi^{\lambda}_{\mu_1 \dots \mu_{s_1-1}, \nu_1 \dots \nu_{s_2}} , \\ \gamma^1 \psi &\equiv \gamma_{(\mu_1} \psi_{\mu_2 \dots \mu_{s_1+1}), \nu_1 \dots \nu_{s_2}} . \end{aligned} \quad (2.3)$$

Together with these family γ -matrices, we shall also need to consider their antisymmetric combinations, that in the two-family case are only γ^{ij} and γ_{ij} . The former adds two Lorentz indices borne by a conventional antisymmetric combination of two γ 's, say $\gamma_{\mu\nu}$, while the latter signals an antisymmetric combination of two γ -traces. In general, it is also convenient to introduce the compact notations

$$\not{\partial} = \gamma_{\mu} \partial^{\mu} , \quad \not{\psi}_i = \gamma_i \psi , \quad (2.4)$$

where γ_{μ} clearly denotes a conventional space-time γ -matrix.

In complete analogy with the bosonic case, the starting point to build the Lagrangians is the Fang-Fronsdal-Labastida kinetic tensor [30]

$$\mathcal{S} = i (\not{\partial} \psi - \partial^i \not{\psi}_i) , \quad (2.5)$$

supplemented with the *unconstrained* gauge transformation of the two-family spinor-tensor ψ ,

$$\delta \psi = \partial^i \epsilon_i . \quad (2.6)$$

As in the bosonic case, this gauge symmetry is reducible, and the corresponding gauge-for-gauge transformations read

$$\delta \epsilon_i = \partial^j \epsilon_{[ij]} , \quad (2.7)$$

where the parameter $\epsilon_{[ij]}$ is antisymmetric in its family indices. At two families the chain stops here, while in the general multi-family setting one must clearly face longer chains of gauge-for-gauge transformations.

The gauge variation of (2.5),

$$\delta \mathcal{S} = -\frac{i}{2} \partial^i \partial^j \gamma_{(i} \epsilon_{j)} , \quad (2.8)$$

⁷The reader should notice that a somewhat unconventional but convenient factor, $\frac{1}{2}$, enters the definition of η^{12} , as in [2].

is proportional to the Labastida constraints on the gauge parameters,

$$\gamma_{(i} \epsilon_{j)} = 0, \quad (2.9)$$

and can be disposed of introducing compensator fields ξ_{ij} such that

$$\delta \xi_{ij} = \frac{1}{2} \gamma_{(i} \epsilon_{j)}. \quad (2.10)$$

As in the bosonic case, however, these compensators are nicely covariant in family notation but can not be independent in general, simply because the Labastida constraints are not independent. Indeed, gauge invariant combinations of the γ -traces of the ξ_{ij} exist⁸, so that they are to be regarded as γ -traces of independent compensators Ψ_i such that

$$\xi_{ij}(\Psi) = \frac{1}{2} \gamma_{(i} \Psi_{j)}, \quad (2.11)$$

with

$$\delta \Psi_i = \epsilon_i. \quad (2.12)$$

Now, in complete analogy with the bosonic construction of [2], or with the one-family case of [17, 19], one can move directly to the unconstrained theory, defining the gauge invariant tensor

$$\mathcal{W} = \mathcal{S} + i \partial^i \partial^j \xi_{ij}(\Psi), \quad (2.13)$$

that in the following will be the basic kinetic tensor for the theory. All our results can be easily adapted to the constrained setting via a partial gauge fixing, and for brevity in the following we shall often write ξ_{ij} to indicate these combinations, without displaying explicitly their dependence on the Ψ_i . Notice that eq. (2.13) involves *two* derivatives of the compensator field. In Section 6 we shall see how more conventional forms of the kinetic operator involving a single derivative obtain via the introduction of a few additional types of fields.

Furthermore, as was the case for bosons, the composite compensator $\xi_{ij}(\Psi)$ of eq. (2.11) would emerge after performing in the Fang-Fronsdal-Labastida tensor $\mathcal{S}$ the Stueckelberg-like shift

$$\psi \rightarrow \psi - \partial^i \Psi_i \quad (2.14)$$

to generate $\mathcal{W}$, an approach we shall elaborate upon at the end of this section⁹. Actually, this would only build the combination (2.11), as in eq. (2.13), a circumstance that reflects the presence of the constrained Labastida gauge symmetry (2.8). Notice that under the gauge-for-gauge transformations (2.7) the Ψ_i fields shift like gauge parameters would, and thus like gauge fields. This is as it should be: if they were inert like the higher-spin field ψ , part of the gauge transformations would be ineffective on them, in contradiction with the manifest possibility of removing the Ψ_i by simply undoing the Stueckelberg shift.

⁸The example presented to this effect in [2] is actually incorrect, since the combination in eq. (2.11) of that paper *is not* gauge invariant. A poignant example, however, is the more complicated expression in eq. (2.49) of [2], that on the contrary *is* gauge invariant and is available in general from two families onward. A corresponding example for Fermi fields will be presented in eq. (2.20). In the bosonic case there is another example that can be analyzed by inspection, a combination of triple traces of the α_{ijk} compensators carrying a $\{5, 3, 1\}$ Young projection in the family indices. Such an expression is gauge invariant simply because one cannot recover its structure starting from four traces of the gauge parameters Λ_i , since multiple T_{ij} lead to Young diagrams with even numbers of boxes in their rows.

⁹M.A. Vasiliev stressed to us the role of this shift in the symmetric case of [17]. In the following we shall see how to formulate the procedure in various ways for mixed-symmetry spinor-tensors.

The Bianchi identities for $\mathcal{W}$ are

$$\partial_i \mathcal{W} - \frac{1}{2} \not{\partial} \mathcal{W}_i - \frac{1}{2} \partial^j T_{ij} \mathcal{W} - \frac{1}{6} \partial^j \gamma_{ij} \mathcal{W} = \frac{1}{2} \partial^j \partial^k \mathcal{Z}_{ijk}, \quad (2.15)$$

where the gauge-invariant spinor-tensor constraints

$$\mathcal{Z}_{ijk} = \frac{i}{3} \left\{ T_{(ij} \not{\psi}_{k)} - 2 \partial_{(i} \xi_{jk)} - \not{\partial} \gamma_{(i} \xi_{jk)} - \partial^l (2 T_{(ij} \xi_{k)l} - T_{l(i} \xi_{jk)} - \gamma_{l(i} \xi_{jk)}) \right\} \quad (2.16)$$

are totally symmetric in their family indices, and when expressed in terms of the independent compensators of eq. (2.11) take the far simpler form

$$\mathcal{Z}_{ijk} = \frac{i}{3} T_{(ij} \gamma_{k)} (\psi - \partial^l \Psi_l). \quad (2.17)$$

After removing the compensators, eq. (2.17) identifies precisely the Labastida constraints on the gauge field,

$$T_{(ij} \not{\psi}_{k)} = 0, \quad (2.18)$$

that can thus be read directly from the Bianchi identities for the Fang-Fronsdal-Labastida tensor $\mathcal{S}$ of [30].

The constraints (2.18) on the gauge fields are *not* independent in general, just like the constraints (2.9) on the gauge parameters, and indeed, once the ξ_{ij} are expressed in terms of the independent compensators Ψ_i , the γ -traces of the $\mathcal{Z}_{ijk}$ spinor-tensors satisfy algebraic relations, so that for instance

$$(\gamma_{m(i} \mathcal{Z}_{j)kl} + \gamma_{m(k} \mathcal{Z}_{l)ij}) - 15 Y_{\{4,1\}} (T_{ij} \mathcal{Z}_{klm} + T_{kl} \mathcal{Z}_{ijm}) = 0, \quad (2.19)$$

where $Y_{\{4,1\}}$ denotes the Young projector onto the irreducible $\{4,1\}$ representation of the permutation group acting on the family indices¹⁰. The lack of independence of the Labastida constraints for the gauge parameters (2.9) and of their counterparts (2.18) for the gauge fields can be traced to a common origin. Indeed, starting from eq. (2.16) the part of the combination (2.19) involving the ξ_{ij} is not forced to vanish a priori. Rather, in complete analogy with the bosonic case of [2] it reduces to

$$\begin{aligned} & (\gamma_{m(i} \mathcal{Z}_{j)kl} + \gamma_{m(k} \mathcal{Z}_{l)ij}) - 15 Y_{\{4,1\}} (T_{ij} \mathcal{Z}_{klm} + T_{kl} \mathcal{Z}_{ijm}) = \\ & = \frac{i}{3} \partial^n \left\{ 8 T_{(ij} T_{kl)} \xi_{mn} - 4 T_{mn} T_{(ij} \xi_{kl)} + 4 T_{m(i} T_{n|j} \xi_{kl)} - 2 T_{(m|(i} T_{jk} \xi_{l)|n)} \right\} \\ & + \frac{i}{3} \partial^n \left\{ 2 T_{(i|(m} \gamma_{n)|j} \xi_{kl)} - 2 \gamma_{(m|(i} T_{jk} \xi_{l)|n)} \right\}, \end{aligned} \quad (2.20)$$

where the left over combination of (γ) -traces of the ξ_{ij} is gauge invariant and is available in general from two families onward. The terms entering eq. (2.20) involve four γ -traces acting on the ξ_{ij} , and in the two-family case it is not possible to obtain gauge invariant combinations containing smaller numbers of γ -traces¹¹. Therefore, the linear dependence of the Labastida

¹⁰The Young projector acts only on the last two terms in eq. (2.19) since, when they are expressed in terms of the fields via eq. (2.17), the first two terms automatically result in a $\{4,1\}$ -projected combination. Our conventions for the permutation group are spelled out at the end of Appendix A.

¹¹In order for the most general linear combination of three γ -traces of the compensators to be gauge invariant several cancelations must occur simultaneously. The resulting system apparently admits a number of solutions, but a closer analysis shows that they all involve combinations of γ -traces of the compensators that are antisymmetric in three indices, and thus vanish identically where only two index families are present.

constraints does not manifest itself in simple mixed-symmetry models with small numbers of space-time indices.

The form of the Bianchi identities (2.15) now suggests to start from the trial Lagrangian

$$\mathcal{L}_0 = \frac{1}{2} \langle \bar{\psi}, \mathcal{W} - \frac{1}{2} \gamma^i \mathcal{W}_i - \frac{1}{2} \eta^{ij} T_{ij} \mathcal{W} - \frac{1}{12} \gamma^{ij} \gamma_{ij} \mathcal{W} \rangle + \text{h.c.}, \quad (2.21)$$

where, as in [2], we have introduced a convenient scalar product that is defined in Appendix A. The gauge variation of $\mathcal{L}_0$ is then

$$\begin{aligned} \delta \mathcal{L}_0 = & -\frac{1}{2} \langle \bar{\epsilon}_i, \partial_i \mathcal{W} - \frac{1}{2} \not{\partial} \mathcal{W}_i - \frac{1}{2} \partial^j T_{ij} \mathcal{W} - \frac{1}{6} \partial^j \gamma_{ij} \mathcal{W} \rangle \\ & + \frac{1}{4} \langle \bar{\epsilon}_i, \gamma^j \partial_i \mathcal{W}_j - \frac{1}{3} \gamma^j \not{\partial} \gamma_{ij} \mathcal{W} \rangle + \frac{1}{4} \langle \bar{\epsilon}_i, \eta^{jk} \partial_i T_{jk} \mathcal{W} \rangle \\ & + \frac{1}{24} \langle \bar{\epsilon}_i, \gamma^{jk} \partial_i \gamma_{jk} \mathcal{W} \rangle + \text{h.c.} \end{aligned} \quad (2.22)$$

and, as in the bosonic case, the key ingredients to build the complete unconstrained Lagrangian are Young projections for the family indices to distinguish in $\delta \mathcal{L}_0$ contributions related to the gauge transformation (2.10) of the composite compensators ξ_{ij} from others to be dealt with via suitable γ -traces of the Bianchi identities. In the fermionic case, however, this identification is not at all straightforward, and is actually fraught with new subtleties, that we shall try to exhibit following a sequence of steps. The Bianchi identities (2.15) can relate directly the first term of eq. (2.22) to the $\mathcal{Z}_{ijk}$ constraint tensors, while the other terms must be split beforehand into different components that transform irreducibly under permutations of the family indices. One is thus led to

$$\begin{aligned} \delta \mathcal{L}_0 = & -\frac{1}{12} \langle \partial_{(i} \partial_j \bar{\epsilon}_{k)}, \mathcal{Z}_{ijk} \rangle \\ & + \frac{1}{16} \langle \bar{\epsilon}_{(i} \gamma_{j)}, \partial_{(i} \mathcal{W}_{j)} \rangle + \frac{1}{16} \langle \bar{\epsilon}_{[i} \gamma_{j]}, \partial_{[i} \mathcal{W}_{j]} - \frac{2}{3} \not{\partial} \gamma_{ij} \mathcal{W} \rangle \\ & + \frac{1}{72} \langle T_{(ij} \bar{\epsilon}_{k)}, \partial_{(i} T_{jk)} \mathcal{W} \rangle + \frac{1}{24} \langle Y_{\{2,1\}} T_{jk} \bar{\epsilon}_i, (2 \partial_i T_{jk} - \partial_{(j} T_{k)i}) \mathcal{W} \rangle \\ & + \frac{1}{72} \langle Y_{\{2,1\}} \bar{\epsilon}_i \gamma_{jk}, (2 \partial_i \gamma_{jk} - \partial_{[j} \gamma_{k]i}) \mathcal{W} \rangle + \text{h.c.}, \end{aligned} \quad (2.23)$$

after discarding the $\{1,1,1\}$ projection, that a priori $\bar{\epsilon}_i \gamma_{jk}$ would admit but is clearly not available with only two families. Aside from ordinary parentheses, that as we stated above denote symmetrizations, here we are also introducing square brackets. As is commonly the case, these denote antisymmetrizations, here built via the minimum possible number of terms and defined with unit overall coefficient.

The second and fourth terms of eq. (2.23) are both symmetrically projected and involve combinations of the type $\bar{\epsilon}_{(i} \gamma_{j)}$. Taking into account that, with the “mostly plus” convention for the metric tensor adopted in this paper,

$$\delta \bar{\xi}_{ij} = -\frac{1}{2} \bar{\epsilon}_{(i} \gamma_{j)}, \quad (2.24)$$

they can be removed introducing

$$\mathcal{L}_1 = \frac{1}{4} \langle \bar{\xi}_{ij}, \frac{1}{2} \partial_{(i} \mathcal{W}_{j)} + \frac{1}{6} \gamma^k \partial_{(i} T_{jk)} \mathcal{W} \rangle + \text{h.c.}, \quad (2.25)$$

that collects all compensator terms that are already visible in the symmetric case of [17, 19]. The total variation then becomes

$$\begin{aligned} \delta(\mathcal{L}_0 + \mathcal{L}_1) = & -\frac{1}{12} \langle \partial_{(i} \partial_j \bar{\epsilon}_k), \mathcal{Z}_{ijk} \rangle + \frac{1}{8} \langle \bar{\epsilon}_i \gamma_j, \partial_{[i} \mathcal{W}_{j]} - \frac{2}{3} \not\partial \gamma_{ij} \mathcal{W} \rangle \\ & + \frac{1}{24} \langle Y_{\{2,1\}} T_{jk} \bar{\epsilon}_i, (2 \partial_i T_{jk} - \partial_{(j} T_{k)i}) \mathcal{W} \rangle \\ & + \frac{1}{72} \langle Y_{\{2,1\}} \bar{\epsilon}_i \gamma_{jk}, (2 \partial_i \gamma_{jk} - \partial_{[j} \gamma_{k]i}) \mathcal{W} \rangle + \text{h.c.} . \end{aligned} \quad (2.26)$$

Aside from the first contribution, that contains the $\mathcal{Z}_{ijk}$ and, as we shall see, can be compensated by Lagrange multipliers λ_{ijk} , this remainder can be dealt with resorting to some consequences of the Bianchi identities. For the second term, the relevant one is the γ -trace of (2.15),

$$\begin{aligned} \partial_{[i} \mathcal{W}_{j]} - \frac{2}{3} \not\partial \gamma_{ij} \mathcal{W} + \frac{1}{6} \partial^k \gamma_{ijk} \mathcal{W} - \frac{1}{6} \partial^k (3 T_{ik} \gamma_j + T_{j[i} \gamma_{k]}) \mathcal{W} \\ = \partial^k \not\partial \mathcal{Z}_{ijk} + \frac{1}{2} \partial^k \partial^l \gamma_j \mathcal{Z}_{ikl} , \end{aligned} \quad (2.27)$$

that actually admits two independent Young projections. The first projection is a symmetric $\{2\}$, that reads

$$\partial^k T_{(ij} \mathcal{W}_{k)} = -6 \partial^k \not\partial \mathcal{Z}_{ijk} - \frac{3}{2} \partial^k \partial^l \gamma_{(i} \mathcal{Z}_{j)kl} , \quad (2.28)$$

and in complete analogy with the bosonic construction of [2] reflects an algebraic identity satisfied by $\mathcal{W}$,

$$T_{(ij} \mathcal{W}_{k)} = 3 \left\{ -2 \not\partial \mathcal{Z}_{ijk} + \partial^l \gamma_l \mathcal{Z}_{ijk} - \frac{1}{2} \partial^l \gamma_{(i} \mathcal{Z}_{jkl)} \right\} . \quad (2.29)$$

Eq. (2.28) contains indeed on the right-hand side the gradient of this expression, as can be seen taking into account that the (k, l) symmetry induced by the additional gradient makes the last two terms directly comparable. On the other hand, the second projection of eq. (2.27) is an antisymmetric $\{1, 1\}$, that reads

$$\partial_{[i} \mathcal{W}_{j]} - \frac{2}{3} \not\partial \gamma_{ij} \mathcal{W} = -\frac{1}{6} \partial^k \gamma_{ijk} \mathcal{W} + \frac{1}{3} \partial^k T_{k[i} \gamma_{j]} \mathcal{W} - \frac{1}{4} \partial^k \partial^l \gamma_{[i} \mathcal{Z}_{j]kl} , \quad (2.30)$$

where the left-hand side is exactly as needed for eq. (2.26). It should be appreciated that for fully symmetric ψ fields only eq. (2.28) would survive, so that the γ -trace of the Bianchi identity would not contribute any useful information.

The remaining two terms in eq. (2.26) present some novel features with respect to the bosonic case that are worthy of some discussion. First of all, taking a further γ -trace of the Bianchi identities gives rise to two distinct types of expressions, a trace T_{jk} and an antisymmetric γ -trace γ_{jk} . The first reads

$$\begin{aligned} \frac{1}{2} (2 \partial_i T_{jk} - \partial_{(j} T_{k)i}) \mathcal{W} + \frac{1}{6} \partial_{(j} \gamma_{k)i} \mathcal{W} - \frac{1}{2} \not\partial T_{jk} \mathcal{W}_i - \frac{1}{2} \partial^l T_{jk} T_{il} \mathcal{W} \\ - \frac{1}{6} \partial^l T_{jk} \gamma_{il} \mathcal{W} = \partial^l \partial_{(j} \mathcal{Z}_{k)i} + \frac{1}{2} \partial^l \partial^m T_{jk} \mathcal{Z}_{ilm} , \end{aligned} \quad (2.31)$$

and admits a $\{3\}$ and a $\{2, 1\}$ Young projection in family indices, while the second reads

$$\begin{aligned} \partial_{[i} \gamma_{jk]} \mathcal{W} + \frac{1}{6} \partial_{[j} \gamma_{k]i} \mathcal{W} + \frac{1}{2} \partial_{[j} T_{k]i} \mathcal{W} - \frac{5}{6} \not\partial \gamma_{ijk} \mathcal{W} + \frac{1}{6} \not\partial T_{i[j} \gamma_{k]} \mathcal{W} \\ - \frac{1}{6} \partial^l \gamma_{ijkl} \mathcal{W} - \frac{1}{2} \partial^l T_{il} \gamma_{jk} \mathcal{W} + \frac{1}{6} \partial^l (T_{i[j} \gamma_{k]l} - T_{l[j} \gamma_{k]i}) \mathcal{W} \\ + \frac{1}{6} \partial^l T_{i[j} T_{k]l} \mathcal{W} = -\partial^l \not\partial \gamma_{[j} \mathcal{Z}_{k]il} + \partial^l \partial_{[j} \mathcal{Z}_{k]il} + \frac{1}{2} \partial^l \partial^m \gamma_{jk} \mathcal{Z}_{ilm} , \end{aligned} \quad (2.32)$$

and would admit in principle a $\{2,1\}$ and a $\{1,1,1\}$ Young projection in the family indices. In analogy with the $\{2\}$ projection of eq. (2.27), the $\{3\}$ projection of eq. (2.31) simply relates structures that are proportional to the constraints $\mathcal{Z}_{ijk}$, and furthermore the $\{1,1,1\}$ projection of eq. (2.32) is clearly not available in the two-family case. The remaining $\{2,1\}$ component is common to both expressions, and will be the key ingredient of the present construction. Indeed, letting $\mathcal{B}_i$ denote for brevity the Bianchi identities (2.15), the two resulting expressions read

$$\begin{aligned}
Y_{\{2,1\}} T_{jk} \mathcal{B}_i : & \quad \frac{1}{2} (2 \partial_i T_{jk} - \partial_{(j} T_{k)i}) \mathcal{W} + \frac{1}{6} \partial_{(j} \gamma_{k)i} \mathcal{W} \\
& - \frac{1}{6} \not\partial (2 T_{jk} \gamma_i - T_{i(j} \gamma_{k)}) \mathcal{W} - \frac{1}{6} \partial^l (2 T_{jk} T_{il} - T_{i(j} T_{k)l}) \mathcal{W} \\
& - \frac{1}{18} \partial^l (2 T_{jk} \gamma_{il} - T_{i(j} \gamma_{k)l}) \mathcal{W} = -\frac{1}{3} \partial^l (2 \partial_i \mathcal{Z}_{jkl} - \partial_{(j} \mathcal{Z}_{k)il}) \\
& + \frac{1}{6} \partial^l \partial^m (2 T_{jk} \mathcal{Z}_{ilm} - T_{i(j} \mathcal{Z}_{k)lm})
\end{aligned} \tag{2.33}$$

and

$$\begin{aligned}
Y_{\{2,1\}} \gamma_{jk} \mathcal{B}_i : & \quad -\frac{1}{18} (2 \partial_i \gamma_{jk} - \partial_{[j} \gamma_{k]i}) \mathcal{W} + \frac{1}{2} \partial_{[j} T_{k]i} \mathcal{W} + \frac{1}{6} \not\partial T_{i[j} \gamma_{k]} \mathcal{W} \\
& - \frac{1}{9} \partial^l (2 T_{il} \gamma_{jk} - T_{l[j} \gamma_{k]i}) \mathcal{W} + \frac{1}{6} \partial^l T_{i[j} \gamma_{k]l} \mathcal{W} + \frac{1}{6} \partial^l T_{i[j} T_{k]l} \mathcal{W} \\
& = -\partial^l \not\partial \gamma_{[j} \mathcal{Z}_{k]il} + \partial^l \partial_{[j} \mathcal{Z}_{k]il} + \frac{1}{6} \partial^l \partial^m (2 \gamma_{jk} \mathcal{Z}_{ilm} - \gamma_{i[j} \mathcal{Z}_{k]lm}) .
\end{aligned} \tag{2.34}$$

Clearly, *neither* of these equations suffices to move past eq. (2.26), as was the case for (2.30) in the previous step, or altogether for bosonic fields in [2], since both the third and fourth contributions in (2.26) lack some of the divergences present on the left-hand sides of eqs. (2.33) and (2.34). In order to proceed, one might therefore try to combine these two consequences of the Bianchi identities, after making their manifest symmetries compatible, since at this stage their differences reflect the nature of the left entries of the corresponding scalar products in (2.26). Let us therefore construct a consequence of eq. (2.34) that is manifestly symmetric in (j, k) like eq. (2.33), and similarly a consequence of eq. (2.33) that is manifestly antisymmetric in (j, k) like eq. (2.34). The issue at stake is whether suitable combinations of these two expressions could reproduce the two sets of divergences in the last two terms of eq. (2.26). Unfortunately, however, the results of these manipulations are

$$\begin{aligned}
\gamma_{i(j} \mathcal{B}_{k)} : & \quad \frac{1}{2} (2 \partial_i T_{jk} - \partial_{(j} T_{k)i}) \mathcal{W} + \frac{1}{6} \partial_{(j} \gamma_{k)i} \mathcal{W} \\
& - \frac{1}{6} \not\partial (2 T_{jk} \gamma_i - T_{i(j} \gamma_{k)}) \mathcal{W} - \frac{1}{6} \partial^l (2 T_{jk} T_{il} - T_{i(j} T_{k)l}) \mathcal{W} \\
& + \frac{1}{2} \partial^l T_{l(j} \gamma_{k)i} \mathcal{W} - \frac{1}{6} \partial^l (2 T_{jk} \gamma_{il} - T_{i(j} \gamma_{k)l} + T_{l(j} \gamma_{k)i}) \mathcal{W} \\
& = -\partial^l \not\partial (2 \gamma_i \mathcal{Z}_{jkl} - \gamma_{(j} \mathcal{Z}_{k)lm}) + \partial^l (2 \partial_i \mathcal{Z}_{jkl} - \partial_{(j} \mathcal{Z}_{k)il}) + \frac{1}{2} \partial^l \partial^m \gamma_{i(j} \mathcal{Z}_{k)lm}
\end{aligned} \tag{2.35}$$

and

$$\begin{aligned}
T_{i[j} \mathcal{B}_{k]} : & \quad -\frac{3}{2} \partial_{[j} T_{k]i} \mathcal{W} + \frac{1}{6} (2 \partial_i \gamma_{jk} + \partial_{[j} \gamma_{k]i}) \mathcal{W} - \frac{1}{2} \not\partial T_{i[j} \mathcal{W}_{k]} \\
& - \frac{1}{2} \partial^l T_{i[j} T_{k]l} \mathcal{W} - \frac{1}{6} \partial^l T_{i[j} \gamma_{k]l} \mathcal{W} = \partial^l \partial_{[j} \mathcal{Z}_{k]il} + \frac{1}{2} \partial^l \partial^m T_{i[j} \mathcal{Z}_{k]lm} ,
\end{aligned} \tag{2.36}$$

and contain divergence terms that are identical, respectively, to those present in (2.33) and (2.34), that have the same manifest symmetries. As a result, no progress can be attained in this fashion.

The conclusion is that one can not forego the need to combine the last two terms in eq. (2.26). This is actually possible, if perhaps surprising, since $T_{jk} \bar{\epsilon}_i$ and $\bar{\epsilon}_i \gamma_{jk}$ are *not* independent, and can be related to one another up to some well-defined compensator terms. Thus, for instance,

$$Y_{\{2,1\}} \bar{\epsilon}_i \gamma_{jk} = \frac{1}{3} (2 \bar{\epsilon}_i \gamma_{jk} - \bar{\epsilon}_{[j} \gamma_{k]i}) = \frac{1}{3} (\bar{\epsilon}_{(i} \gamma_{j)k} - \bar{\epsilon}_{(i} \gamma_{k)j}) , \quad (2.37)$$

and each term in the last sum can be related to traces of the gauge parameters and to the compensators letting

$$\bar{\epsilon}_{(i} \gamma_{j)k} = \bar{\epsilon}_{(i} \gamma_{j)} \gamma_k - T_{k(i} \bar{\epsilon}_{j)} . \quad (2.38)$$

Similar considerations apply to the other combination present in eq. (2.26), so that one finally obtains

$$Y_{\{2,1\}} T_{jk} \bar{\epsilon}_i = \frac{2}{3} (2 \delta \bar{\xi}_{jk} \gamma_i - \delta \bar{\xi}_{i(j} \gamma_{k)}) + \bar{\epsilon}_{(j} \gamma_{k)i} , \quad (2.39)$$

$$Y_{\{2,1\}} \bar{\epsilon}_i \gamma_{jk} = -\frac{2}{3} \delta \bar{\xi}_{i[j} \gamma_{k]} + \frac{1}{3} T_{i[j} \bar{\epsilon}_{k]} . \quad (2.40)$$

This circumstance clearly brings about some ambiguities, since one is somehow working with a set of structures that are *not* linearly independent. The simplest way to proceed would be to choose a basis, and in the next section we shall eventually do it. However, it is quite instructive to first dwell upon these ambiguities, introducing a new parameter, κ , to put some emphasis on their role. Let us therefore cast a portion κ of the last two terms in (2.26) in the T_{jk} format, making use of eq. (2.40) in the fourth term and the remaining portion $(1 - \kappa)$ in the γ_{jk} format, making use of eq. (2.39) in the third term. Some of the resulting contributions are then manifestly proportional to the variation of the compensators, and can be canceled adding

$$\mathcal{L}_2 = \frac{1}{12} \langle \bar{\xi}_{ij} , (\kappa - 1) \gamma^k (2 \partial_k T_{ij} - \partial_{(i} T_{j)k}) \mathcal{W} + \frac{\kappa}{3} \gamma^k \partial_{(i} \gamma_{j)k} \mathcal{W} \rangle + \text{h.c.} , \quad (2.41)$$

so that

$$\begin{aligned} \delta(\mathcal{L}_0 + \mathcal{L}_1 + \mathcal{L}_2) = & -\frac{1}{12} \langle \partial_{(i} \partial_{j)} \bar{\epsilon}_k , \mathcal{Z}_{ijk} \rangle + \frac{1}{8} \langle \bar{\epsilon}_i \gamma_j , \partial_{[i} \mathcal{W}_{j]} - \frac{2}{3} \not{\partial} \gamma_{ij} \mathcal{W} \rangle \\ & + \frac{\kappa}{24} \langle T_{jk} \bar{\epsilon}_i , (2 \partial_i T_{jk} - \partial_{(j} T_{k)i}) \mathcal{W} + \frac{1}{3} \partial_{(j} \gamma_{k)i} \mathcal{W} \rangle \\ & - \frac{1 - \kappa}{8} \langle \bar{\epsilon}_i \gamma_{jk} , \partial_{[j} T_{k]i} \mathcal{W} - \frac{1}{9} (2 \partial_i \gamma_{jk} - \partial_{[j} \gamma_{k]i}) \mathcal{W} \rangle + \text{h.c.} . \end{aligned} \quad (2.42)$$

We have thus obtained two types of terms that can be taken care of via the Bianchi identities, but before proceeding a further comment is in order. The issue is that, for the special case of two-family gauge fields, some of the γ -traces of $\mathcal{W}$ that appear in eqs. (2.33), (2.34), (2.35) and (2.36) are actually proportional to the constraints, although this is not evident at first sight. Indeed, in these expressions the terms carrying four lower family indices are a priori of the form

$$\begin{array}{|c|c|} \hline & \\ \hline & \\ \hline \end{array} \otimes \begin{array}{|c|} \hline \\ \hline \end{array} \equiv \begin{array}{|c|c|c|} \hline & & \\ \hline & & \\ \hline \end{array} \oplus \begin{array}{|c|c|} \hline & \\ \hline & \\ \hline \end{array} \oplus \begin{array}{|c|} \hline \\ \hline \\ \hline \\ \hline \end{array}$$

where the first Young diagram factor is dictated by the projection while the additional box originates from the gradient. However, the subset of terms containing both a trace and a two-index γ actually admit only the $\{3, 1\}$ and $\{2, 1, 1\}$ projections, as one can see from the composition rule

$$\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} \otimes \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} \equiv \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & & \end{array} \oplus \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \\ \hline \end{array}$$

Here the symmetric boxes originate from the trace T while the antisymmetric ones originate from the two-index γ , and as we stressed repeatedly with only two families three-row projections like the $\{2, 1, 1\}$ are clearly impossible. As a result, one is finally left only with the $\{3, 1\}$ projection for these terms, that as such is related to the $\mathcal{Z}_{ijk}$. For example, a contribution of this type is present in eq. (2.33), and as we just stressed coincides with its $\{3, 1\}$ portion

$$(2T_{jk}\gamma_{il} - T_{i(j}\gamma_{k)l})\mathcal{W} \xrightarrow{\{3,1\}} \frac{1}{2}T_{(ij}\gamma_{k)l}\mathcal{W} + \frac{3}{4}(T_{il}\gamma_{j)k} + T_{(il}\gamma_{k)j})\mathcal{W}, \quad (2.43)$$

that is proportional to the $\mathcal{Z}_{ijk}$ on account of eq. (2.29), since

$$T_{(ij}\gamma_{k)l}\mathcal{W} = -\gamma_l T_{(ij}\mathcal{W}_{k)} + \frac{1}{4}\gamma_{(i}T_{jk}\mathcal{W}_{l)}. \quad (2.44)$$

Actually, an alternative possibility exists to select the components not related to the constraints, that is more suitable for an extension to generic fields with an arbitrary number of index families. It amounts to working with linear combinations of eqs. (2.33) and (2.35), or of eqs. (2.34) and (2.36), that do not contain the $\{3, 1\}$ projection of terms of this type. The relevant relations in this two-family setting are

$$\begin{aligned} \frac{9}{8}Y_{\{2,1\}}\left(T_{jk}\mathcal{B}_i - \frac{1}{9}\gamma_{i(j}\mathcal{B}_{k)}\right) &: \frac{1}{2}(2\partial_i T_{jk} - \partial_{(j}T_{k)i})\mathcal{W} + \frac{1}{6}\partial_{(j}\gamma_{k)i}\mathcal{W} \\ &- \frac{1}{6}\partial^l(2T_{jk}\gamma_i - T_{i(j}\gamma_{k)})\mathcal{W} - \frac{1}{6}\partial^l(2T_{jk}T_{il} - T_{i(j}T_{k)l})\mathcal{W} \\ &= \frac{1}{8}\partial^l\partial(\gamma_i\mathcal{Z}_{jkl} - \gamma_{(j}\mathcal{Z}_{k)il}) - \frac{1}{2}\partial^l(2\partial_i\mathcal{Z}_{jkl} - \partial_{(j}\mathcal{Z}_{k)il}) \\ &+ \frac{3}{16}\partial^l\partial^m(2T_{jk}\mathcal{Z}_{ilm} - T_{i(j}\mathcal{Z}_{k)lm}) - \frac{1}{16}\partial^l\partial^m\gamma_{i(j}\mathcal{Z}_{k)lm}, \end{aligned} \quad (2.45)$$

together with their counterparts obtained by an antisymmetrization in (i, j) and a cyclic relabeling,

$$\begin{aligned} \frac{9}{8}Y_{\{2,1\}}\left(\frac{1}{3}\gamma_{jk}\mathcal{B}_i + T_{i[j}\mathcal{B}_{k]}\right) &: -\frac{3}{2}\partial_{[j}T_{k]i}\mathcal{W} + \frac{1}{6}(2\partial_i\gamma_{jk} - \partial_{[j}\gamma_{k]i})\mathcal{W} \\ &- \frac{1}{2}\partial T_{i[j}\mathcal{W}_{k]} - \frac{1}{2}\partial^l T_{i[j}T_{k]l}\mathcal{W} = -\frac{3}{8}\partial^l\partial\gamma_{[j}\mathcal{Z}_{k]il} + \frac{3}{2}\partial^l\partial_{[j}\mathcal{Z}_{k]il} \\ &+ \frac{9}{16}\partial^l\partial^m T_{i[j}\mathcal{Z}_{k]lm} + \frac{1}{16}\partial^l\partial^m(2\gamma_{jk}\mathcal{Z}_{ilm} - \gamma_{i[j}\mathcal{Z}_{k]lm}). \end{aligned} \quad (2.46)$$

These two consequences of the Bianchi identities now reduce the total variation (2.42) to

$$\begin{aligned} \delta(\mathcal{L}_0 + \mathcal{L}_1 + \mathcal{L}_2) &= -\frac{1}{4}\langle\delta\bar{\lambda}_{ijk}, \mathcal{Z}_{ijk}\rangle + \frac{1}{24}\langle\bar{\epsilon}_i\gamma_j, \partial^k T_{k[i}\gamma_{j]}\mathcal{W}\rangle \\ &+ \frac{\kappa}{72}\langle T_{jk}\bar{\epsilon}_i, \partial(2T_{jk}\gamma_i - T_{i(j}\gamma_{k)})\mathcal{W} + \partial^l(2T_{jk}T_{il} - T_{i(j}T_{k)l})\mathcal{W}\rangle \\ &+ \frac{1-\kappa}{24}\langle\bar{\epsilon}_i\gamma_{jk}, \partial T_{i[j}\gamma_{k]}\mathcal{W} + \partial^l T_{i[j}T_{k]l}\mathcal{W}\rangle + \text{h.c.}, \end{aligned} \quad (2.47)$$

where

$$\begin{aligned}
\delta\lambda_{ijk} = & \frac{1}{3} \partial_{(i} \partial_j \epsilon_k) + \frac{1}{24} \gamma^l (\gamma_{(i} \partial_j \partial_k) \epsilon_l - \gamma_l \partial_{(i} \partial_j \epsilon_k)) \\
& + \frac{\kappa}{18} \partial^l (2T_{ij} \partial_k \epsilon_l - T_{l(i} \partial_j \epsilon_k)) + \frac{1-\kappa}{6} \partial^l \gamma_{l(i} \partial_j \epsilon_k) \\
& - \frac{\kappa}{72} \gamma^l \not\partial (2T_{ij} \partial_k \epsilon_l - T_{l(i} \partial_j \epsilon_k)) - \frac{1-\kappa}{24} \gamma^l \not\partial \gamma_{l(i} \partial_j \epsilon_k) \\
& - \frac{\kappa}{12} \eta^{lm} (T_{lm} \partial_{(i} \partial_j \epsilon_k) - T_{l(i} \partial_j \partial_k) \epsilon_m) + \frac{1-\kappa}{4} \eta^{lm} \gamma_{l(i} \partial_j \partial_k) \epsilon_m \\
& + \frac{\kappa}{72} \gamma^{lm} T_{l(i} \partial_j \partial_k) \epsilon_m - \frac{1-\kappa}{72} \gamma^{lm} (\gamma_{lm} \partial_{(i} \partial_j \epsilon_k) + \gamma_{l(i} \partial_j \partial_k) \epsilon_m) .
\end{aligned} \tag{2.48}$$

can be regarded as the gauge variation of a set of Lagrange multipliers λ_{ijk} . The rest of the variation (2.47) can be partly canceled adding

$$\begin{aligned}
\mathcal{L}_3 = & \frac{1}{2} \langle \bar{\psi}, \frac{1}{18} \eta^{ij} \gamma^k (2T_{ij} \gamma_k - T_{k(i} \gamma_{j)}) \mathcal{W} + \frac{1}{36} \eta^{ij} \eta^{kl} (2T_{ij} T_{kl} - T_{i(k} T_{l)j}) \mathcal{W} \rangle \\
& + \text{h.c.} ,
\end{aligned} \tag{2.49}$$

but one is readily to face a novel feature of the fermionic construction: an additional compensator term,

$$\mathcal{L}_4 = \frac{1}{72} \langle \bar{\xi}_{ij}, \partial^k (2T_{ij} \gamma_k - T_{k(i} \gamma_{j)}) \mathcal{W} \rangle + \text{h.c.} , \tag{2.50}$$

is needed to eliminate the symmetric portion of the gradient terms produced when varying the first contribution in eq. (2.49). At this point

$$\begin{aligned}
\delta(\mathcal{L}_0 + \dots + \mathcal{L}_4) = & -\frac{1}{4} \langle \delta \bar{\lambda}_{ijk}, \mathcal{Z}_{ijk} \rangle \\
& - \frac{1-\kappa}{72} \langle T_{jk} \bar{\epsilon}_i, \not\partial (2T_{jk} \gamma_i - T_{i(j} \gamma_{k)}) \mathcal{W} + \partial^l (2T_{jk} T_{il} - T_{i(j} T_{k)l}) \mathcal{W} \rangle \\
& + \frac{1-\kappa}{24} \langle \bar{\epsilon}_i \gamma_{jk}, \not\partial T_{i[j} \gamma_{k]} \mathcal{W} + \partial^l T_{i[j} T_{k]l} \mathcal{W} \rangle \\
& - \frac{1}{72} \langle T_{jk} \bar{\epsilon}_i \gamma_l, \partial_i (2T_{jk} \gamma_l - T_{l(j} \gamma_{k)}) \mathcal{W} \rangle \\
& - \frac{1}{288} \langle T_{jk} T_{lm} \bar{\epsilon}_i, \partial_i (2T_{jk} T_{lm} - T_{l(j} T_{k)m}) \mathcal{W} \rangle + \text{h.c.} ,
\end{aligned} \tag{2.51}$$

and one is thus confronted, again, with two apparently distinct types of structures, those present in the two terms above and proportional to $(1-\kappa)$. They can be combined making use again of eqs. (2.39) and (2.40), only to discover that they fully cancel, up to some terms proportional to gauge variation $\delta \xi_{ij}$ of the compensators, that can be taken care of adding

$$\begin{aligned}
\mathcal{L}_5 = & \frac{1-\kappa}{36} \langle \bar{\xi}_{ij}, \gamma^k \not\partial (2T_{ij} \gamma_k - T_{k(i} \gamma_{j)}) \mathcal{W} + \gamma^k \partial^l (2T_{ij} T_{kl} - T_{k(i} T_{j)l}) \mathcal{W} \rangle \\
& + \text{h.c.} .
\end{aligned} \tag{2.52}$$

Finally, *at two families*, the last two terms in eq. (2.51) can be turned into a form that can be manifestly canceled by the additional compensator terms in

$$\begin{aligned}
\mathcal{L}_6 = & -\frac{1}{288} \langle \bar{\xi}_{ij}, \gamma^{kl} (2\gamma_l T_{ij} \partial_k - T_{l(i} \gamma_j \partial_k) \mathcal{W} \\
& + 2\eta^{kl} (2T_{ij} \gamma_{(k} \partial_l) + 2T_{kl} \gamma_{(i} \partial_j) - \gamma_{(i} T_{j)(k} \partial_l - \gamma_{(k} T_{l)(i} \partial_j) \mathcal{W} \rangle \\
& - \frac{1}{288} \langle \bar{\xi}_{ij}, \eta^{kl} \gamma^m (2T_{kl} T_{ij} \partial_m - T_{k(i} T_{l|j} \partial_m) \mathcal{W} \rangle + \text{h.c.} .
\end{aligned} \tag{2.53}$$

This is clearly a further manifestation of the absence of Young tableaux with more than two rows. Its origin can be understood observing that at two families the fourth term in eq. (2.51) contains only the $\{3, 1\}$ projection

$$\begin{aligned} & \langle T_{jk} \bar{\epsilon}_i \gamma_l, \partial_i (2T_{jk} \gamma_l - T_{l(j} \gamma_{k)}) \mathcal{W} \rangle \xrightarrow{\{3,1\}} \\ & \frac{1}{24} \langle 2T_{(ij} \bar{\epsilon}_{k)} \gamma_l - T_{l(i} \bar{\epsilon}_{j} \gamma_{k)}, (2\gamma_l T_{(ij} \partial_{k)} - T_{l(i} \gamma_j \partial_{k)}) \mathcal{W} \rangle, \end{aligned} \quad (2.54)$$

and the $\{2, 2\}$ projection

$$\begin{aligned} & \langle T_{jk} \bar{\epsilon}_i \gamma_l, \partial_i (2T_{jk} \gamma_l - T_{l(j} \gamma_{k)}) \mathcal{W} \rangle \xrightarrow{\{2,2\}} \\ & \frac{1}{8} \langle T_{jk} \bar{\epsilon}_{(i} \gamma_{l)}, (2T_{jk} \gamma_{(i} \partial_{l)} + 2T_{il} \gamma_{(j} \partial_{k)} - \gamma_{(j} T_{k)(i} \partial_{l)} - \gamma_{(i} T_{l)(j} \partial_{k)}) \mathcal{W} \rangle, \end{aligned} \quad (2.55)$$

both clearly related to the gauge variation of the compensators. In a similar fashion, at two families the last term in eq. (2.51) only admits a $\{3, 2\}$ projection, and can be directly related to the variation of the compensators on account of

$$\begin{aligned} & (2T_{ij} T_{kl} - T_{i(k} T_{l)j}) \bar{\epsilon}_m \xrightarrow{\{3,2\}} \\ & \frac{1}{4} \{ 3T_{ij} T_{(kl} \bar{\epsilon}_{m)} + 3T_{kl} T_{(ij} \bar{\epsilon}_{m)} - T_{(ij|} (2T_{|kl)} \bar{\epsilon}_m + T_{m|k} \bar{\epsilon}_l) \}, \end{aligned} \quad (2.56)$$

that played a similar role also in the bosonic case of [2].

As a result, for two-family fermionic fields we have obtained a whole set of complete unconstrained Lagrangians labeled by the parameter κ ,

$$\mathcal{L}_{fin} = \frac{1}{2} \langle \bar{\psi}, \mathcal{E} \rangle + \frac{1}{4} \langle \bar{\xi}_{ij}, \Xi_{ij}(\kappa) \rangle + \frac{1}{4} \langle \bar{\lambda}_{ijk}, \mathcal{Z}_{ijk} \rangle + \text{h.c.}, \quad (2.57)$$

with the κ -independent tensor

$$\begin{aligned} \mathcal{E} \equiv & \mathcal{W} - \frac{1}{2} \gamma^i \mathcal{W}_i - \frac{1}{2} \eta^{ij} T_{ij} \mathcal{W} - \frac{1}{12} \gamma^{ij} \gamma_{ij} \mathcal{W} \\ & + \frac{1}{18} \eta^{ij} \gamma^k (2T_{ij} \gamma_k - T_{k(i} \gamma_{j)}) \mathcal{W} + \frac{1}{36} \eta^{ij} \eta^{kl} (2T_{ij} T_{kl} - T_{i(k} T_{l)j}) \mathcal{W}, \end{aligned} \quad (2.58)$$

that generalizes the Rarita-Schwinger kinetic tensor, and

$$\begin{aligned} \Xi_{ij}(\kappa) \equiv & \frac{1}{2} \partial_{(i} \mathcal{W}_{j)} + \frac{1}{6} \gamma^k \partial_{(i} T_{jk)} \mathcal{W} + \frac{1}{18} \partial^k (2T_{ij} \gamma_k - T_{k(i} \gamma_{j)}) \mathcal{W} \\ & - \frac{1-\kappa}{3} \gamma^k (2\partial_k T_{ij} - \partial_{(i} T_{j)k}) \mathcal{W} + \frac{\kappa}{9} \gamma^k \partial_{(i} \gamma_{j)k} \mathcal{W} \\ & + \frac{1-\kappa}{9} \gamma^k \left\{ \not{\partial} (2T_{ij} \gamma_k - T_{k(i} \gamma_{j)}) \mathcal{W} + \partial^l (2T_{ij} T_{kl} - T_{k(i} T_{j)l}) \mathcal{W} \right\} \\ & - \frac{1}{36} \eta^{kl} (2T_{ij} \gamma_{(k} \partial_{l)} + 2T_{kl} \gamma_{(i} \partial_{j)} - \gamma_{(i} T_{j)(k} \partial_{l)} - \gamma_{(k} T_{l)(i} \partial_{j)}) \mathcal{W} \\ & - \frac{1}{72} \gamma^{kl} (2\gamma_l T_{(ij} \partial_{k)} - T_{l(i} \gamma_j \partial_{k)}) \mathcal{W} \\ & - \frac{1}{72} \eta^{kl} \gamma^m (2T_{kl} T_{(ij} \partial_{m)} - T_{k(i} T_{l|j} \partial_{m)}) \mathcal{W}. \end{aligned} \quad (2.59)$$

At the same time, we have determined corresponding κ -dependent gauge variations of the λ_{ijk} Lagrange multipliers, that are given in eq. (2.48).

One can also explain the origin of these ambiguities noticing that, in sharp contrast with the bosonic constructions of [2], the non-trivial fermionic Bianchi identities that drive our construction carry Young projections that are *not* forbidden for the γ -traces of the compensators. Via their traces or γ -traces, in fact, the ξ_{ij} give rise in general to Young diagrams with at least two columns, that can also characterize the relevant consequences of the Bianchi identities, like the $\{2,1\}$ projections that we used repeatedly. On the other hand, for bosonic fields the consequences of the Bianchi identities that drive the construction are two-column projected, while compensator terms contain at least three columns, so that the relevant consequences of the Bianchi identities and the constraints on the gauge parameters (2.9) correspond to orthogonal Young projections. In the present example we can see explicitly this phenomenon after separating out in the Lagrangian all terms proportional to $(1 - \kappa)$, so that

$$\mathcal{L}_{fin} = \mathcal{L} + (1 - \kappa) \tilde{\mathcal{L}}, \quad (2.60)$$

with

$$\begin{aligned} \mathcal{L} = & \frac{1}{2} \langle \bar{\psi}, \mathcal{E} \rangle + \frac{1}{4} \langle \bar{\lambda}_{ijk}, \mathcal{Z}_{ijk} \rangle \\ & + \frac{1}{4} \langle \bar{\xi}_{ij}, \frac{1}{2} \partial_{(i} \mathcal{W}_{j)} + \frac{1}{6} \gamma^k \partial_{(i} T_{jk)} \mathcal{W} + \frac{1}{9} \gamma^k \partial_{(i} \gamma_{j)k} \mathcal{W} \\ & + \frac{1}{18} \partial^k (2 T_{ij} \gamma_k - T_{k(i} \gamma_{j)}) \mathcal{W} - \frac{1}{72} \gamma^{kl} (2 \gamma_l T_{(ij} \partial_{k)} - T_{l(i} \gamma_{j)} \partial_{k)}) \mathcal{W} \\ & - \frac{1}{36} \eta^{kl} (2 T_{ij} \gamma_{(k} \partial_{l)} + 2 T_{kl} \gamma_{(i} \partial_{j)} - \gamma_{(i} T_{j)(k} \partial_{l)} - \gamma_{(k} T_{l)(i} \partial_{j)}) \mathcal{W} \\ & - \frac{1}{72} \eta^{kl} \gamma^m (2 T_{kl} T_{(ij} \partial_{m)} - T_{k(i} T_{l|j} \partial_{m)}) \mathcal{W} \rangle + \text{h.c.} \end{aligned} \quad (2.61)$$

and

$$\begin{aligned} \tilde{\mathcal{L}} = & - \frac{1}{12} \langle \bar{\xi}_{ij} \gamma_k, (2 \partial_k T_{ij} - \partial_{(i} T_{j)k}) \mathcal{W} + \frac{1}{3} \partial_{(i} \gamma_{j)k} \mathcal{W} \\ & - \frac{1}{3} \not{\partial} (2 T_{ij} \gamma_k - T_{k(i} \gamma_{j)}) \mathcal{W} - \frac{1}{3} \partial^l (2 T_{ij} T_{kl} - T_{k(i} T_{j)l}) \mathcal{W} \rangle + \text{h.c.} . \end{aligned} \quad (2.62)$$

The key feature of this last set of terms is that they build exactly the left-hand side of the Bianchi identities (2.45), so that they can be turned into

$$\begin{aligned} \tilde{\mathcal{L}} = & \frac{1}{36} \langle \bar{\mathcal{Z}}_{ijk}, \partial^l (2 \gamma_l \partial_{(i} \xi_{jk)} - \gamma_{(i} \partial_j \xi_{k)l}) - \frac{1}{4} \gamma^l \not{\partial} (2 \gamma_l \partial_{(i} \xi_{jk)} - \gamma_{(i} \partial_j \xi_{k)l}) \\ & + \frac{3}{2} \eta^{lm} (\gamma_{(i} \partial_j \partial_{k)} \xi_{lm} - \gamma_l \partial_{(i} \partial_j \xi_{k)m}) + \frac{1}{4} \gamma^{lm} \gamma_l \partial_{(i} \partial_j \xi_{k)m} \rangle + \text{h.c.} , \end{aligned} \quad (2.63)$$

which is a mere redefinition of the Lagrange multipliers λ_{ijk} . A similar result, although involving a different redefinition, would obtain collecting the terms proportional to κ . In conclusion, in the fermionic case there exist field redefinitions that modify at the same time compensator terms and gauge transformations of the Lagrange multipliers, without affecting the types of projections that enter those terms or the Rarita-Schwinger-like tensor of eq. (2.58). In Section 3.2 we shall present other classes of field redefinitions affecting the Rarita-Schwinger-like tensor, that lead to convenient alternative presentations of the unconstrained Lagrangians, while in eq. (2.57) the special choice $\kappa = 1$ will lead to somewhat simpler expressions, and therefore from now on we shall often stick to it.

In conclusion, our two-family Lagrangians are finally given in eq. (2.61), a result that can be also presented in the more compact form

$$\begin{aligned}
\mathcal{L} = & \frac{1}{2} \langle \bar{\psi}, \mathcal{W} - \frac{1}{2} \gamma^i \mathcal{W}_i - \frac{1}{2} \eta^{ij} T_{ij} \mathcal{W} - \frac{1}{12} \gamma^{ij} \gamma_{ij} \mathcal{W} \\
& + \frac{1}{6} \eta^{ij} \gamma^k Y_{\{2,1\}} T_{ij} \gamma_k \mathcal{W} + \frac{1}{12} \eta^{ij} \eta^{kl} Y_{\{2,2\}} T_{ij} T_{kl} \mathcal{W} \rangle \\
& + \frac{1}{4} \langle \bar{\xi}_{ij}, \frac{1}{2} \partial_{(i} \mathcal{W}_{j)} + \frac{1}{6} \gamma^k \partial_{(i} T_{jk)} \mathcal{W} + \frac{1}{9} \gamma^k \partial_{(i} \gamma_{j)k} \mathcal{W} \\
& + \frac{1}{6} \partial^k Y_{\{2,1\}} T_{ij} \gamma_k \mathcal{W} - \frac{1}{24} \gamma^{kl} \partial_{(i} Y_{\{2,1\}} T_{|jk)} \gamma_l \mathcal{W} - \frac{1}{6} \eta^{kl} Y_{\{2,2\}} T_{ij} \gamma_{(k} \partial_{l)} \mathcal{W} \\
& - \frac{1}{24} \eta^{kl} \gamma^m \partial_{(i} Y_{\{2,2\}} T_{|jm)} T_{kl} \mathcal{W} \rangle + \frac{1}{4} \langle \bar{\lambda}_{ijk}, \mathcal{Z}_{ijk} \rangle + \text{h.c.} ,
\end{aligned} \tag{2.64}$$

where we have indicated concisely the Young projections that give rise to the various structures in eq. (2.61). The corresponding gauge transformations for the Lagrange multipliers are then

$$\begin{aligned}
\delta \lambda_{ijk} = & \frac{1}{3} \partial_{(i} \partial_j \epsilon_{k)} + \frac{1}{24} \gamma^l (\gamma_{(i} \partial_j \partial_k) \epsilon_l - \gamma_l \partial_{(i} \partial_j \epsilon_{k)}) \\
& + \frac{1}{18} \partial^l (2 T_{(ij} \partial_k) \epsilon_l - T_{l(i} \partial_j \epsilon_{k)}) - \frac{1}{72} \gamma^l \partial (2 T_{(ij} \partial_k) \epsilon_l - T_{l(i} \partial_j \epsilon_{k)}) \\
& - \frac{1}{12} \eta^{lm} (T_{lm} \partial_{(i} \partial_j \epsilon_{k)} - T_{l(i} \partial_j \partial_k) \epsilon_m) + \frac{1}{72} \gamma^{lm} T_{l(i} \partial_j \partial_k) \epsilon_m .
\end{aligned} \tag{2.65}$$

In practice this type of choice favors, at every stage, structures with higher numbers of T_{ij} tensors, that enter the various expressions via their most antisymmetric projections. These have a relatively simple structure, and as we shall see shortly this choice will prove particularly effective in leading to Lagrangians for fields with an arbitrary number of index families. In the one-family case, the κ -dependent Lagrangians of eq. (2.57) clearly reduce to the result of [17, 19]. In a notation that is essentially as in those papers but with the scalar product of Appendix A, they would read

$$\mathcal{L} = \frac{1}{2} \langle \bar{\psi}, \mathcal{W} - \frac{1}{2} \gamma \mathcal{W} - \frac{1}{2} \eta \mathcal{W}' \rangle + \frac{1}{4} \langle \bar{\xi}, \partial \cdot \mathcal{W} \rangle - \frac{1}{8} \langle \bar{\xi}, \partial \cdot \mathcal{W}' \rangle + \frac{1}{4} \langle \bar{\lambda}, \mathcal{Z} \rangle + \text{h.c.} . \tag{2.66}$$

In addition, the terms not involving the compensators or the Lagrange multipliers, that are independent of κ , are relatively simple and yet quite interesting, since they determine completely the *constrained* two-family fermionic Lagrangian leading to the (non-Lagrangian) Labastida equation of [30]:

$$\begin{aligned}
\mathcal{L} = & \frac{1}{2} \langle \bar{\psi}, \mathcal{S} - \frac{1}{2} \gamma^i \mathcal{S}_i - \frac{1}{2} \eta^{ij} T_{ij} \mathcal{S} - \frac{1}{12} \gamma^{ij} \gamma_{ij} \mathcal{S} \\
& + \frac{1}{18} \eta^{ij} \gamma^k (2 T_{ij} \gamma_k - T_{k(i} \gamma_{j)}) \mathcal{S} + \frac{1}{36} \eta^{ij} \eta^{kl} (2 T_{ij} T_{kl} - T_{i(k} T_{l)j}) \mathcal{S} \rangle + \text{h.c.} .
\end{aligned} \tag{2.67}$$

Let us conclude by noticing that, in analogy with the bosonic case of [2], once the ξ_{ij} are expressed in terms of the independent Ψ_i multipliers, eq. (2.57) acquires a further gauge symmetry, related to shifts of the Lagrange multipliers of the type

$$\delta \lambda_{ijk} = \eta^{lm} M_{ijk,lm} + \gamma^{lm} N_{ijk;lm} , \tag{2.68}$$

where $M_{ijk,lm}$ and $N_{ijk;lm}$ are $\{4,1\}$ projected in their family indices and are related according to ¹²

$$N_{ij(k;l)m} + N_{kl(i;j)m} = -\frac{1}{2} (M_{ijm,kl} + M_{klm,ij}) . \quad (2.69)$$

In fact, making use of eq. (2.17) one can see that the shift (2.68) gives rise to

$$\begin{aligned} \delta \mathcal{L} = & \frac{i}{24} \langle \bar{M}_{ijk,mn}, T_{mn} T_{(ij} \gamma_k) (\psi - \partial^l \Psi_l) \rangle \\ & - \frac{i}{12} \langle \bar{N}_{ijk,mn}, (\gamma_{mn(i} T_{jk)} - T_{(ij} T_{k)[m} \gamma_{n]}) (\psi - \partial^l \Psi_l) \rangle + \text{h.c.} . \end{aligned} \quad (2.70)$$

Indeed, the first scalar product can give contributions in the $\{5\}$, $\{4,1\}$ and $\{3,2\}$ representations. On the other hand, the first term in the second scalar product (that actually vanishes when only two families are present) can give contributions in the $\{3,1,1\}$ and $\{2,1,1,1\}$, and finally the last can give contributions in the $\{4,1\}$ and $\{3,1,1\}$. Therefore, if the M and N parameters are $\{4,1\}$ projected and are related to one another as in eq. (2.69), an interesting symmetry of the Lagrangian (2.57) emerges, that actually continues to hold in the presence of an *arbitrary* number of index families. This symmetry, as we have already stressed in [2] when discussing its bosonic counterpart, reflects the lack of independence of the Labastida constraints on the gauge fields, that we displayed in eq. (2.19). As we shall see in Section 2.3, it manifests itself rather clearly when one tries to reduce the Lagrangian field equations to the Labastida form, since all contributions that can be shifted away by (2.68) are left undetermined. Furthermore, as in the bosonic case, an important lesson to be drawn from this symmetry is that only part of the gauge transformation (2.48) of the Lagrange multipliers λ_{ijk} is effective when one works with the independent Ψ_i compensators.

One can also present the Lagrangians (2.57) in a different form, that proves very convenient when deriving their field equations. A similar rewriting was already described in [2] for Bose fields. As in that case, it can be built starting from the Lagrangians for fields ψ *not* subject to any constraints but realizing nonetheless only the constrained gauge transformations of eqs. (2.6) and (2.9),

$$\begin{aligned} \mathcal{L}_C = & \frac{1}{2} \langle \bar{\psi}, \mathcal{S} - \frac{1}{2} \gamma^i \mathcal{S}_i - \frac{1}{2} \eta^{ij} T_{ij} \mathcal{S} - \frac{1}{12} \gamma^{ij} \gamma_{ij} \mathcal{S} + \frac{1}{6} \eta^{ij} \gamma^k Y_{\{2,1\}} T_{ij} \gamma_k \mathcal{S} \\ & + \frac{1}{12} \eta^{ij} \eta^{kl} Y_{\{2,2\}} T_{ij} T_{kl} \mathcal{S} \rangle + \frac{i}{12} \langle \bar{\zeta}_{ijk}, T_{(ij} \psi_{k)} \rangle + \text{h.c.} , \end{aligned} \quad (2.71)$$

obtained introducing the ζ_{ijk} Lagrange multipliers. These enforce on-shell the Labastida constraints (2.18), and as a result are to transform as the λ_{ijk} did in eq. (2.48). Any of the κ -dependent transformations (2.48) grants the gauge invariance of (2.71), on account of the previous discussion, while the unconstrained Lagrangians of eq. (2.57) can be recovered performing the κ -dependent Stueckelberg-like shifts

$$\begin{aligned} \psi & \rightarrow \psi - \partial^i \Psi_i , \\ \zeta_{ijk} & \rightarrow \lambda_{ijk} - \Delta_{ijk}(\Psi) , \end{aligned} \quad (2.72)$$

where $\Delta_{ijk}(\Psi)$ is the combination that maps the ζ_{ijk} into gauge invariant expressions. From the technical viewpoint, the explicit rewriting of these unconstrained Lagrangians (2.57) involves

¹²Here and in the following a semi-colon separates symmetric and antisymmetric subsets of *family* indices, so that, for instance, $N_{ijk;lm}$ is symmetric under interchanges of (i,j,k) and is antisymmetric under interchanges of l and m .

precisely the same steps of the previous direct derivation, but this presentation makes it possible to relate the equations of motion for $\bar{\psi}$, the $\bar{\Psi}_i$ and the $\bar{\lambda}_{ijk}$ to the simpler ones of eq. (2.71).

Finally, we would like to call to the reader's attention that, as in the bosonic case, one could also work with independent ξ_{ij} compensators, at the cost of adding to the Lagrangian (2.57) new *gauge invariant* multipliers ρ_{ij} according to

$$\mathcal{L}_7 = \langle \bar{\rho}_{ij}, \frac{1}{2} \gamma_{(i} \Psi_{j)} - \xi_{ij} \rangle + \text{h.c.} . \quad (2.73)$$

2.2 THE FIELD EQUATIONS

In the previous section we described three different classes of Lagrangians for “two-family” Fermi fields or, equivalently, for “two-row” reducible spinorial representations of the Poincaré group. We first presented, in eq. (2.57), the Lagrangians for unconstrained spinor-tensors ψ . We then recovered from them, in eq. (2.67), their counterparts for fields that are subject to the triple γ -trace constraints (2.18). In this second presentation, the Lagrangians are only invariant under gauge transformations whose parameters are subject to the γ -trace constraints (2.9). Finally, in eq. (2.71) we exhibited Lagrangians that involve unconstrained fields but are still only invariant under constrained gauge transformations. Now we would like to derive the field equations that follow from all these Lagrangians. To this end, it is convenient to start from eq. (2.71).

The field equations for the Lagrange multipliers ζ_{ijk} read

$$E_{\bar{\zeta}} : \frac{i}{12} T_{(ij} \psi_{k)} = 0 . \quad (2.74)$$

On the other hand, the field equation for the gauge field $\bar{\psi}$ requires more work, since one must combine the terms displayed in eq. (2.71) with their hermitian conjugates, and this step requires an extensive use of some of the identities collected in Appendix B. The end result is

$$\begin{aligned} E_{\bar{\psi}} : \mathcal{E} - \frac{1}{2} \eta^{ij} \gamma^k \mathcal{Y}_{ijk} - \frac{i}{12} \left\{ \eta^{ij} \partial^k T_{(ij} \psi_{k)} - \frac{1}{6} \eta^{ij} \gamma^k \partial^l \left(\gamma_j T_{(ik} \psi_{l)} - \gamma_k T_{(ij} \psi_{l)} \right) \right. \\ \left. + \frac{1}{8} \eta^{ij} \eta^{kl} Y_{\{2,2\}} \partial^m \left(T_{ij} T_{(kl} \psi_{m)} + T_{kl} T_{(ij} \psi_{m)} \right) \right\} = 0 , \end{aligned} \quad (2.75)$$

where

$$\begin{aligned} \mathcal{E} \equiv \mathcal{S} - \frac{1}{2} \gamma^i \mathcal{S}_i - \frac{1}{2} \eta^{ij} T_{ij} \mathcal{S} - \frac{1}{12} \gamma^{ij} \gamma_{ij} \mathcal{S} \\ + \frac{1}{18} \eta^{ij} \gamma^k (2 T_{ij} \gamma_k - T_{k(i} \gamma_{j)}) \mathcal{S} + \frac{1}{36} \eta^{ij} \eta^{kl} (2 T_{ij} T_{kl} - T_{i(k} T_{l)j}) \mathcal{S} \end{aligned} \quad (2.76)$$

can be regarded as the fermionic analogue of the Einstein tensor of [12, 2], and the

$$\begin{aligned} \mathcal{Y}_{ijk} \equiv i \left\{ \zeta_{ijk} - \frac{1}{6} T_{(ij} \partial_{k)} \psi + \frac{1}{72} \gamma^l (2 \gamma_l T_{(ij} \partial_{k)} - T_{l(i} \gamma_{j)} \partial_{k)}) \psi \right. \\ \left. + \frac{1}{72} \eta^{lm} (2 T_{lm} T_{(ij} \partial_{k)} - T_{l(i} T_{m|j} \partial_{k)}) \psi \right\} \end{aligned} \quad (2.77)$$

are totally symmetric under interchanges of their family labels. Eq. (2.76) generalizes the Rarita-Schwinger spinor-tensor to the case of mixed-symmetry higher-spin Fermi fields, and can be

also recovered gauging away the compensators in eq. (2.58). For the constrained theory with Lagrange multipliers of eq. (2.71), combining eqs. (2.74) and (2.75) leads to

$$E_{\bar{\psi}} : \mathcal{E} - \frac{1}{2} \eta^{ij} \gamma^k \mathcal{Y}_{ijk} = 0. \quad (2.78)$$

Let us add a few comments on the role of eq. (2.78) in the fully constrained framework, where naively the spinor-tensors $\mathcal{Y}_{ijk}$ should not be present. In fact, when dealing with constrained fields one cannot conclude directly that

$$\mathcal{E} = 0, \quad (2.79)$$

since in general

$$T_{(ij} \gamma_k) \mathcal{E} \neq 0, \quad (2.80)$$

in apparent contrast with the properties of the gauge field ψ . However, a field equation satisfying the same constraints as the gauge field obtains after eliminating the $\mathcal{Y}_{ijk}$ tensors from eq. (2.78). The proper projection is thus encoded in the tensorial structure of the $\mathcal{Y}_{ijk}$, that hence can play a role even in the constrained theory. We shall provide an explicit example of this fact in eqs. (2.180) and (2.195), where we shall display various presentations of the equations of motion for a rank-(3, 2) field. Notice that the projected equations couple to external currents satisfying the Labastida constraints, thus extending the Fang-Fronsdal symmetric setting. However, field equations and currents that are jointly redefined to satisfy different constraints lead to the same physical current exchanges.

Finally, the equations of the unconstrained Lagrangians (2.57) follow from eqs. (2.74) and (2.75), taking into account that

$$\begin{aligned} \delta \mathcal{L}_C &= \langle \delta \bar{\psi}, E_{\bar{\psi}} \rangle + \langle \delta \bar{\zeta}_{ijk}, (E_{\bar{\zeta}})_{ijk} \rangle + \text{h.c.} \\ &= \langle \delta \bar{\psi}, E_{\bar{\psi}} \rangle + \langle \delta \bar{\Psi}_i, \partial_i E_{\bar{\psi}} - \tilde{\Delta}_{ijk} (E_{\bar{\zeta}}) \rangle + \langle \delta \bar{\lambda}_{ijk}, (E_{\bar{\zeta}})_{ijk} \rangle + \text{h.c.}, \end{aligned} \quad (2.81)$$

where the $\tilde{\Delta}_{ijk}$ are obtained integrating by parts the terms in the Δ_{ijk} of eq. (2.72). In conclusion, one thus obtains

$$\begin{aligned} E_{\bar{\psi}} : \mathcal{E} - \frac{1}{2} \eta^{ij} \gamma^k \mathcal{Y}_{ijk} - \frac{1}{4} \left\{ \eta^{ij} \partial^k \mathcal{Z}_{ijk} - \frac{1}{6} \eta^{ij} \gamma^k \partial^l \gamma_{[j} \mathcal{Z}_{k]il} \right. \\ \left. + \frac{1}{12} \eta^{ij} \eta^{kl} \partial^m (T_{ij} \mathcal{Z}_{klm} + T_{kl} \mathcal{Z}_{ijm} - T_{k(i} \mathcal{Z}_{j)lm}) \right\} = 0, \end{aligned} \quad (2.82)$$

$$\begin{aligned} E_{\bar{\Psi}} : \partial_i E_{\bar{\psi}} - \partial^j \partial^k (E_{\bar{\lambda}})_{ijk} - \frac{1}{8} \gamma^j \gamma_{[i} \partial^k \partial^l (E_{\bar{\lambda}})_{j]kl} \\ - \frac{\kappa}{4} \eta^{jk} T_{k[i} \partial^l \partial^m (E_{\bar{\lambda}})_{j]lm} - \frac{\kappa}{12} \eta^{jk} \gamma_{ij} \partial^l \partial^m (E_{\bar{\lambda}})_{klm} \\ + \frac{1-\kappa}{8} \gamma^{jk} T_{ij} \partial^l \partial^m (E_{\bar{\lambda}})_{klm} - \frac{1-\kappa}{24} \gamma^{jk} \gamma_{k(i} \partial^l \partial^m (E_{\bar{\lambda}})_{j)lm} = 0, \end{aligned} \quad (2.83)$$

$$E_{\bar{\lambda}} : \frac{1}{4} \mathcal{Z}_{ijk} = 0, \quad (2.84)$$

where now $\mathcal{E}$ denotes the Rarita-Schwinger-like tensor of eq. (2.58), while the $\mathcal{Y}_{ijk}$ tensors are built performing the Stueckelberg shifts (2.72) in eq. (2.77), barring some subtleties that we shall discuss shortly. Notice that the equations of motion for the compensators $\bar{\Psi}_i$, here written for brevity in a form that is not “normal ordered”, simply provide the conservation conditions

for an external current coupled to ψ . The latter is in fact always conserved as a result of the unconstrained gauge symmetry, as in the symmetric construction of [17, 19].

We would like to conclude this section with some comments on the λ_{ijk} Lagrange multipliers and on the Stueckelberg shifts (2.72) since, as we anticipated at the end of Section 2.1, some parts of the naive gauge transformation (2.48) of the λ_{ijk} have no effect on the Lagrangian when the compensators are expressed in terms of the Ψ_i . These parts could well be discarded, and different choices give rise to different definitions of the Stueckelberg shifts for the ζ_{ijk} in eq. (2.72). The reader familiar with the construction of [2] will probably notice a close analogy with the behavior of bosonic Lagrange multipliers. In fact, in [2] we identified redundant portions of their gauge transformations that made the $\mathcal{B}_{ijkl}$ tensors, bosonic analogs of the $\mathcal{Y}_{ijk}$ tensors, not gauge invariant. This, however, did not affect the gauge invariance of $\eta^{ij} \eta^{kl} \mathcal{B}_{ijkl}$, the combination that enters the field equations of the bosonic gauge fields. In a similar fashion, the $\mathcal{Y}_{ijk}$ tensors that enter eq. (2.82) are not invariant under the gauge transformations of eqs. (2.6) and (2.10) when these are combined with the *naive* gauge transformations of the λ_{ijk} multipliers given in eq. (2.48), since

$$\begin{aligned} \delta \mathcal{Y}_{ijk} = & -\frac{i}{12} \eta^{lm} Y_{\{4,1\}} \left\{ \kappa (T_{lm} \partial_{(i} \partial_j \epsilon_k) - T_{l(i} \partial_j \partial_k) \epsilon_m) - 3(1 - \kappa) \gamma_{l(i} \partial_j \partial_k) \epsilon_m \right\} \\ & + \frac{i}{72} \gamma^{lm} Y_{\{4,1\}} \left\{ \kappa T_{l(i} \partial_j \partial_k) \epsilon_m - (1 - \kappa) (\gamma_{lm} \partial_{(i} \partial_j \epsilon_k) + \gamma_{l(i} \partial_j \partial_k) \epsilon_m) \right\}, \end{aligned} \quad (2.85)$$

but proceeding as in (2.70) one can see that $\eta^{ij} \gamma^k \mathcal{Y}_{ijk}$ is properly gauge invariant. For both Bose and Fermi fields, dropping the redundant terms one can define gauge transformations for the multipliers that make the $\mathcal{B}_{ijkl}$ or the $\mathcal{Y}_{ijk}$ directly gauge invariant. Alternatively, if one were to work with the naive transformations of the multipliers, the resulting gauge variation (2.85) of the $\mathcal{Y}_{ijk}$ tensors could be compensated by a particular shift under the additional gauge symmetry of eq. (2.68). This, as we have seen, gives rise to the transformations

$$\delta \mathcal{Y}_{ijk} = \eta^{lm} M_{ijk,lm} + \gamma^{lm} N_{ijk;lm}, \quad (2.86)$$

with two $\{4,1\}$ projected parameters, and the two contributions present in eq. (2.85) also satisfy eq. (2.69), that relates to one another the two parameters M and N of the additional shift symmetry. At any rate, the additional shift symmetry (2.86) is present independently of the choice of gauge transformation for the multipliers, provided one works in terms of the Ψ_i .

These observations are particularly important when the unconstrained theory is built making use of Stueckelberg shifts starting from the Lagrangian (2.71): in this case the $\mathcal{Y}_{ijk}$ are gauge invariant by construction, but if one starts from the naive gauge transformations of the multipliers the result is *not* expressible in terms of the composite compensators $\xi_{ij}(\Psi)$. In order to illustrate this fact, it is convenient to split the expression obtained via the *naive* Stueckelberg shift (2.72) in eq. (2.77) as

$$\mathcal{Y}_{ijk} = \mathcal{Y}^{(1)}_{ijk} + \gamma^l \mathcal{Y}^{(2)}_{ijk,l} + \eta^{lm} \mathcal{Y}^{(3)}_{ijk,lm} + \gamma^{lm} \mathcal{Y}^{(4)}_{ijk;lm}. \quad (2.87)$$

The first term contains the contributions that are already visible in the symmetric case ¹³,

$$\begin{aligned} \mathcal{Y}^{(1)}_{ijk} = & i \left\{ \lambda_{ijk} - \frac{1}{6} T_{(ij} \partial_k) \psi + \frac{1}{6} \square \gamma_{(i} \xi_{jk)} + \frac{4\kappa - 3}{18} \partial^l \gamma_{l(i} \partial_{(i} \xi_{jk)} \right. \\ & \left. - \frac{2\kappa - 3}{18} \partial^l \partial_{(i} \gamma_j \xi_{k)l} \right\}. \end{aligned} \quad (2.88)$$

¹³Notice, however, that in the symmetric, or one-family, case the κ dependence disappears.

The second term adds some contributions that first emerge in the two-family case but are still insensitive to the redundant portion of the gauge transformations of the Lagrange multipliers, and at two families reads

$$\begin{aligned} \mathcal{Y}^{(2)}_{ijk,l} = \frac{i}{72} \Big\{ & (2\gamma_l T_{(ij} \partial_k) - T_{l(i} \gamma_j \partial_k) \psi - 2(\partial_{(i} \partial_j \xi_k)_{l)} - \partial_l \partial_{(i} \xi_{jk)}) \\ & - 2\Box \gamma_{l(i} \xi_{jk)} + 2(1-\kappa) \not{\partial} (2\gamma_l \partial_{(i} \xi_{jk)} - \partial_{(i} \gamma_j \xi_k)_{l)}) \\ & - \frac{1}{2} \partial^m (2T_{lm} \partial_{(i} \xi_{jk)} + 2T_{(ij} \partial_k) \xi_{lm} - T_{l(i} \partial_j \xi_k)_{m)} - T_{m(i} \partial_j \xi_k)_{l)}) \\ & - \frac{1}{4} \partial^m (2\gamma_{lm} \partial_{(i} \xi_{jk)} + 3\gamma_{l(i} \partial_j \xi_k)_{m)} + \gamma_{m(i} \partial_j \xi_k)_{l)}) \Big\}. \end{aligned} \quad (2.89)$$

On the other hand, the naive form of the third and the fourth terms in eq. (2.87) contains some contributions with bare Ψ_i compensators, and reads

$$\begin{aligned} \mathcal{Y}^{(3)}_{ijk,lm} = \frac{i}{24} Y_{\{3,2\}} \Big\{ & T_{lm} T_{(ij} \partial_k) \psi + 2(1-\kappa) (2\gamma_{(i} \partial_j \partial_k) \xi_{lm} + \gamma_{(l} \partial_m) \partial_{(i} \xi_{jk)}) \\ & - \frac{1}{2} \Box (T_{lm} \gamma_{(i} \xi_{jk)} + T_{(ij} \gamma_k) \xi_{lm} + \gamma_{(l} T_{(ij} \xi_k)_{|m)}) \\ & - \frac{1}{4} \partial^n (\gamma_n T_{(ij} \partial_k) \xi_{lm} + \gamma_n T_{lm} \partial_{(i} \xi_{jk)} + T_{lm} \partial_{(i} \gamma_j \xi_k)_{n)} + T_{(ij} \partial_k) \gamma_{(l} \xi_{m)n)}) \Big\} \\ & + \frac{i}{24} Y_{\{4,1\}} \Big\{ \kappa (2T_{lm} \partial_{(i} \partial_j \Psi_k) - T_{(l|i} \partial_j \partial_k) \Psi_{|m)}) - 3(1-\kappa) \gamma_{(l|i} \partial_j \partial_k) \Psi_{|m)} \Big\} \end{aligned} \quad (2.90)$$

for the third term and

$$\begin{aligned} \mathcal{Y}^{(4)}_{ijk;lm} = -\frac{i}{144} Y_{\{4,1\}} \Big\{ & \kappa T_{[l|i} \partial_j \partial_k) \Psi_{|m]} \\ & - (1-\kappa) (2\gamma_{lm} \partial_{(i} \partial_j \Psi_k) + \gamma_{[l|i} \partial_j \partial_k) \Psi_{|m]}) \Big\} \end{aligned} \quad (2.91)$$

for the last one. The terms containing bare Ψ_i compensate precisely the gauge variation (2.85) of the previous, ξ_{ij} dependent, $\mathcal{Y}_{ijk}$ tensors.

2.3 ON-SHELL REDUCTION TO $\mathcal{S} = 0$

We can now conclude our discussion of two-family fields by showing how the three sets of Lagrangian field equations described in the preceding section are equivalent to the Labastida equation of motion¹⁴

$$\mathcal{S} = 0. \quad (2.92)$$

In analogy with the case of Bose fields [2], in the following we shall discuss how this result can be attained either directly or after fixing extra Weyl-like gauge symmetries that emerge for the Lagrangians in sporadic low-dimensional cases.

To begin with, we would like to recall that the equations of motion for $\bar{\psi}$ that follow from the Lagrangians (2.67) and (2.71) can both be cast in the form (2.78). In addition, combining

¹⁴In Section 3.4.1 we shall see that eq. (2.92) and the (constrained) gauge symmetry (2.6) suffice in general to turn the spinorial $gl(D)$ representations carried by ψ fields into their $o(D-2)$ counterparts, as expected in view of the uniqueness of $\mathcal{S}$.

eqs. (2.82) and (2.84) and gauging away the compensators, the field equation for $\bar{\psi}$ following from the Lagrangian (2.57) can be also reduced to eq. (2.78). Thus, for all three Lagrangian theories that we presented one is led to discuss the reduction of

$$E_{\bar{\psi}} : \mathcal{S} - \frac{1}{2} \gamma^i \mathcal{S}_i - \frac{1}{2} \eta^{ij} T_{ij} \mathcal{S} - \frac{1}{12} \gamma^{ij} \gamma_{ij} \mathcal{S} + \frac{1}{18} \eta^{ij} \gamma^k (2 T_{ij} \gamma_k - T_{k(i} \gamma_{j)}) \mathcal{S} \\ + \frac{1}{36} \eta^{ij} \eta^{kl} (2 T_{ij} T_{kl} - T_{i(k} T_{l)j}) \mathcal{S} - \frac{1}{2} \eta^{ij} \gamma^k \mathcal{Y}_{ijk} = 0, \quad (2.93)$$

$$E_{\bar{\zeta}} : \frac{i}{12} T_{(ij} \psi_{k)} = 0 \implies T_{(ij} \mathcal{S}_{k)} = 0, \quad (2.94)$$

to the Labastida form (2.92), where of course in the constrained case the second condition holds even off shell.

To this end, one can conveniently adapt the procedure developed in [2] for N -family bosonic fields. The picture one has in mind is that whenever eq. (2.93) *does not* reduce directly to the conditions

$$\mathcal{S} = 0, \\ \mathcal{Y}_{ijk} = 0, \quad (2.95)$$

additional symmetries are present. In fact, this is equivalent to stating that the homogeneous equation (2.93) admits non-trivial solutions aside from eq. (2.95), and now we would like to classify and describe these “pathological” cases. Actually, we already came across a phenomenon of this kind in Section 2.2, where we saw that eq. (2.93) is left invariant by the shifts of the $\mathcal{Y}_{ijk}$ defined in eq. (2.86). In this case the undetermined quantities do not affect the portion of the field equation involving $\mathcal{S}$, and are thus immaterial insofar as its reduction to the Labastida form is concerned. At the same time, they can be gauged away directly, making use of the shift symmetry of the Lagrange multipliers identified in eq. (2.68).

We can now show that, in sporadic examples with low space-time dimensions, some γ -traces of the spinor-tensor $\mathcal{S}$ are left undetermined, a phenomenon that is however accompanied by the emergence of new Weyl-like gauge symmetries for the field ψ . These are precisely as needed to gauge away the undetermined quantities, thus making it possible to complete the reduction to the Labastida form. Let us stress that this peculiar behavior is already visible for symmetric spinor-tensors, since the Rarita-Schwinger equation (2.93) for a gravitino can be cast in the form

$$\mathcal{S}_\mu - \frac{1}{2} \gamma_\mu \gamma \cdot \mathcal{S} = 0. \quad (2.96)$$

In two dimensions this formal expression only involves the γ -traceless part of $\mathcal{S}$, and thus apparently one cannot use it to set to zero its γ -trace. However, precisely in two dimensions Weyl-like shifts of the form

$$\delta \psi_\mu = \gamma_\mu \Omega \quad (2.97)$$

become a symmetry and make it possible to gauge away the undetermined quantity. Again in partial analogy with the case of two-dimensional gravity, this behavior is accompanied by the vanishing of the Rarita-Schwinger Lagrangian, and thus of eq. (2.96) when it is expressed in terms of ψ . This property is manifest in the usual presentation of the Lagrangian,

$$\mathcal{L} = \frac{i}{2} \psi_\mu \gamma^{\mu\nu\rho} \partial_\nu \psi_\rho + \text{h.c.}, \quad (2.98)$$

simply because the fully antisymmetric $\gamma^{\mu\nu\rho}$ vanishes in two dimensions. In the following we shall see that the whole class of one-column fully-antisymmetric fields exhibits a similar behavior in low enough space-time dimensions, but in generic mixed-symmetry cases the situation is more complicated, and in particular there are Weyl-invariant theories whose Lagrangians *do not* vanish identically.

Before starting our analysis of generic two-family fields, let us recall that the relevant $\mathcal{S}$ spinor-tensors are to satisfy the Bianchi identities, that reduce to

$$\mathcal{B}_i : \partial_i \mathcal{S} - \frac{1}{2} \not{\partial} \mathcal{S}_i - \frac{1}{2} \partial^j T_{ij} \mathcal{S} - \frac{1}{6} \partial^j \gamma_{ij} \mathcal{S} = 0 \quad (2.99)$$

after making use of the constraints (2.94), and are in addition to preserve the constraints

$$T_{(ij} \mathcal{S}_{k)} = 0. \quad (2.100)$$

As we have seen, these conditions are identically satisfied in the constrained case, while in the unconstrained case they hold after enforcing the field equations for the Lagrange multipliers. At any rate, out of the non-trivial solutions of eq. (2.93), or equivalently out of its shift symmetries, one must thus consider only those satisfying both (2.99) and (2.100). This key observation greatly simplifies the search for the field equations that do not reduce directly to the Labastida form, since non-trivial shifts of the Bianchi identities provide a most convenient starting point to attack the problem. The Bianchi identities really lie at the heart of the Rarita-Schwinger-like tensors, and we shall see shortly that all their shift symmetries also leave the field equations invariant.

In principle, eq. (2.93) can admit shift symmetries of the form ¹⁵

$$\delta \mathcal{S} = \gamma^i \Omega_i, \quad (2.101)$$

and now we would like to classify those preserving the Bianchi identities (2.99). In general, under shifts of the form (2.101) they vary according to ¹⁶

$$\begin{aligned} \delta \mathcal{B}_i = & -\frac{1}{2} \not{\partial} \left[(D-2) \Omega_i + 2 S^j{}_i \Omega_j \right] - \frac{1}{6} \partial^j \gamma_{[i|} \left[(D-2) \Omega_{|j]} + 2 S^k{}_{|j]} \Omega_k \right] \\ & + \gamma^j \mathcal{B}_i(\Omega_j), \end{aligned} \quad (2.102)$$

where $\mathcal{B}_i(\Omega_j)$ denotes the Bianchi identities for the parameters Ω_j . Thus, if the conditions

$$(D-2) \Omega_i + 2 S^j{}_i \Omega_j = 0 \quad (2.103)$$

are satisfied by non-trivial parameters Ω_i that are themselves subject to the Bianchi identities, the corresponding shift (2.101) preserves indeed eq. (2.99). Before presenting the solutions of this eigenvalue problem, we would like to stress that eq. (2.103) is clearly a *sufficient* condition to identify a symmetry of the Bianchi identities. In the general case, in fact, the contributions that appear in eq. (2.102) could in principle balance each other. However, one can proceed to look for solutions in an iterative manner, analyzing directly more complicated shifts of the Bianchi identities that involve more than one “naked” γ -matrix.

¹⁵Covariance justifies this assertion, although we cannot exclude degenerate cases where further cancelations take place. An example of this type is indeed the highly degenerate case of the $\{2,2\}$ field in two dimensions, that however is also captured by our analysis.

¹⁶With a slight abuse of notation, both in this section and in Section 3.4, with $\mathcal{B}_i$ and “Bianchi identities” we shall often denote the left-hand sides of eqs. (2.99).

At two families a more general shift involving pairs of γ -matrices is fully available, so that one can also consider

$$\delta \mathcal{S} = \eta^{ij} \Omega_{ij} + \gamma^{ij} \tilde{\Omega}_{ij}, \quad (2.104)$$

where the two parameters are symmetric and antisymmetric, respectively, in their family indices. This new type of shift gives rise to a more complicated variation of the Bianchi identities,

$$\begin{aligned} \delta \mathcal{B}_i = & -\frac{1}{2} \not{\partial} \gamma^j \left[\Omega_{ij} - 2 S^k_{(i} \tilde{\Omega}_{j)k} \right] - \not{\partial} \gamma^j \left[(D-3) \tilde{\Omega}_{ij} - S^k_{[i} \tilde{\Omega}_{j]k} \right] \\ & - \frac{1}{2} \partial^j \left[(D-2) \Omega_{ij} + S^k_{(i} \Omega_{j)k} + 2 S^k_{(i} \tilde{\Omega}_{j)k} \right] \\ & + \frac{1}{3} \partial^j \gamma^k_{[i} \left[(D-3) \tilde{\Omega}_{j]k} - 2 S^l_{[j} \tilde{\Omega}_{kl} + \frac{1}{2} \Omega_{j]k} \right] \\ & + \frac{D+2}{3} \partial^j \left[(D-3) \tilde{\Omega}_{ij} - S^k_{[i} \tilde{\Omega}_{j]k} \right] + \eta^{jk} \mathcal{B}_i(\Omega_{jk}) + \gamma^{jk} \mathcal{B}_i(\tilde{\Omega}_{jk}), \end{aligned} \quad (2.105)$$

where we have conveniently introduced the new quantities

$$\gamma^i{}_j = \gamma^i \gamma_j - S^i{}_j, \quad (2.106)$$

that belong to a class of operators described in Appendix B. One can thus identify some purely algebraic conditions granting that (2.105) vanish,

$$\begin{aligned} \Omega_{ij} &= 2 S^k_{(i} \tilde{\Omega}_{j)k}, \\ (D-3) \tilde{\Omega}_{ij} - S^k_{[i} \tilde{\Omega}_{j]k} &= 0. \end{aligned} \quad (2.107)$$

These eliminate all terms involving $\not{\partial}$ from eq. (2.105). All the rest then evidently vanishes, provided the Ω_{ij} and $\tilde{\Omega}_{ij}$ satisfy the Bianchi identities, with the single exception of the second line¹⁷. However, this can be turned into the second of eqs. (2.107) using the first and noticing that

$$S^k_{(i} S^l_{j)} \tilde{\Omega}_{kl} = \frac{1}{2} [S^k_{(i}, S^l_{j)}] \tilde{\Omega}_{kl} = -S^k_{(i} \tilde{\Omega}_{j)k}. \quad (2.108)$$

The present setting becomes degenerate at the next step, since a fully antisymmetric γ_{ijk} is clearly not available with only two families. As a result, the next shift in the iteration takes the form

$$\delta \mathcal{S} = \eta^{ij} \gamma^k \Omega_{ij;k}, \quad (2.109)$$

and gives rise to the variation

$$\begin{aligned} \delta \mathcal{B}_l = & -\frac{1}{2} \not{\partial} \eta^{ij} \left[(D-2) \Omega_{ij;l} + \Omega_{l(i;j)} + 2 S^m_l \Omega_{ij;m} \right] - \frac{1}{4} \not{\partial} \gamma^{ij} \Omega_{l[i;j]} \\ & - \frac{1}{6} \eta^{ij} \partial^m \gamma_{[l} \left[(D-2) \Omega_{ij;m]} + \Omega_{[m](i;j)} + 2 S^n_{[m} \Omega_{ij;n]} \right] - \frac{1}{12} \gamma^{ij} \partial^m \gamma_{[l} \Omega_{m][i;j]} \\ & - \frac{1}{2} \gamma^k \partial^m \left[(D-2) \Omega_{lm;k} + \Omega_{k(l;m)} + S^n_{(l} \Omega_{m)n;k} - \frac{1}{6} S^n_{[l} \Omega_{m](k;n)} \right] \\ & - \frac{1}{12} \gamma^k \partial^m \left[2(D-1) \Omega_{k[l;m]} - 3 S^n_{[l} \Omega_{m][k;n]} \right] + \eta^{ij} \gamma^k \mathcal{B}_l(\Omega_{ij;k}) \end{aligned} \quad (2.110)$$

¹⁷Notice that these new parameters *are not* special cases of the previous ones, as one might naively think, since they are selected by different conditions.

of the Bianchi identities. Again, the conditions that eliminate the terms with ∂ ,

$$\begin{aligned} (D-2)\Omega_{ij;k} + \Omega_{k(i;j)} + 2S^l_k \Omega_{ij;l} &= 0, \\ \Omega_{k[i;j]} &= 0, \end{aligned} \quad (2.111)$$

set to zero all other contributions if the parameters satisfy themselves the Bianchi identities. Indeed, the second condition in (2.111) implies that all combinations of parameters giving rise to a $\{2,1\}$ Young projection in family indices vanish, and consequently the first condition reduces to

$$\begin{aligned} 3D\widehat{\Omega}_{ijk} + 2S^l_{(i}\widehat{\Omega}_{jk)l} &= 0, \\ S^l_{[i}\widehat{\Omega}_{j]kl} &= 0, \end{aligned} \quad (2.112)$$

where

$$\widehat{\Omega}_{ijk} = Y_{\{3\}} \Omega_{ij;k} = \frac{1}{3} \Omega_{(ij;k)}. \quad (2.113)$$

We shall see shortly that eqs. (2.112) admit solutions at two families, while in Section 3.4 we shall also see that $\{2,1\}$ -projected combinations of parameters become available in the presence of more than two index families, when the second condition in eq. (2.111) acquires further contributions.

In principle one should move on and consider shifts of the form

$$\delta \mathcal{S} = \eta^{ij} \eta^{kl} \Omega_{ij,kl} + \eta^{ij} \gamma^{kl} \widetilde{\Omega}_{ij;kl}, \quad (2.114)$$

since only two types of parameters with four indices are available with only two families. The corresponding variation of the Bianchi identities reads

$$\begin{aligned} \delta \mathcal{B}_l &= \partial \eta^{ij} \gamma^k \left[(D-3) \widetilde{\Omega}_{ij;kl} - \widetilde{\Omega}_{l(i;j)k} - \Omega_{ij,kl} + 2S^m_l \widetilde{\Omega}_{ij;km} \right] \\ &\quad - \frac{1}{3} \partial^m \eta^{ij} \gamma^k_{[l} \left[(D-3) \widetilde{\Omega}_{ij;k|m]} - \widetilde{\Omega}_{|m|(i;j)k} - \Omega_{ij,k|m]} + 2S^n_{|m]} \widetilde{\Omega}_{ij;kn} \right] \\ &\quad - \eta^{ij} \partial^k \left[(D-2) \Omega_{ij,kl} + \Omega_{i(k,l)j} + S^m_{(k} \Omega_{l)m,ij} + S^m_{(k} \widetilde{\Omega}_{ij;|l)m} \right] \\ &\quad - \frac{1}{3} \eta^{ij} \partial^k \left[(D+2) \left((D-3) \widetilde{\Omega}_{ij;kl} - S^m_{[k} \widetilde{\Omega}_{ij;|l)m} \right) - 2(D-1) \widetilde{\Omega}_{i[k;l]j} \right. \\ &\quad \left. - S^m_{[k} \widetilde{\Omega}_{l](i;j)m} \right] - \frac{1}{2} \gamma^{ij} \partial^k \left[(D-2) \widetilde{\Omega}_{kl;ij} + 2\widetilde{\Omega}_{i(k;l)j} + S^m_{(k} \widetilde{\Omega}_{l)m;ij} \right] \\ &\quad + \frac{1}{6} \gamma^{ij} \partial^k \left[2(D-2) \widetilde{\Omega}_{i[k;l]j} + \Omega_{i[k,l]j} + S^m_{[k} \widetilde{\Omega}_{l]m;ij} + 2S^m_{[k} \widetilde{\Omega}_{l][i;j]m} \right] \\ &\quad + \eta^{ij} \eta^{mn} \mathcal{B}_l(\Omega_{ij,mn}) + \eta^{ij} \gamma^{mn} \mathcal{B}_l(\widetilde{\Omega}_{ij;mn}), \end{aligned} \quad (2.115)$$

so that now the conditions

$$(D-3) \widetilde{\Omega}_{ij;kl} - \widetilde{\Omega}_{l(i;j)k} - \Omega_{ij,kl} + 2S^m_l \widetilde{\Omega}_{ij;km} = 0 \quad (2.116)$$

eliminate the ∂ terms. However, at the end of the present section we shall prove that these conditions do not admit any solution for two-family fields, in contrast with the previous ones.

At two families the iteration stops here. Indeed, eqs. (2.100) imply that all combinations of more than four γ -traces acting on $\mathcal{S}$ vanish when only two index families are present, and therefore the two chains of shifts

$$\delta \mathcal{S} = \eta^{i_1 j_1} \dots \eta^{i_p j_p} \gamma^k \Omega_{i_1 j_1, \dots, i_p j_p; k}, \quad (2.117)$$

$$\delta \mathcal{S} = \eta^{i_1 j_1} \dots \eta^{i_p j_p} \Omega_{i_1 j_1, \dots, i_p j_p} + \eta^{i_1 j_1} \dots \eta^{i_{p-1} j_{p-1}} \gamma^{kl} \widetilde{\Omega}_{i_1 j_1, \dots, i_{p-1} j_{p-1}; kl}, \quad (2.118)$$

are expected to yield no further information, a fact that we verified using similar techniques.

We have thus classified the conditions characterizing the relevant symmetries of the Bianchi identities. While we shall discuss in some detail the corresponding eigenvalue problems in the second part of this section, we can now elaborate further upon the conditions imposed by eq. (2.100), before showing that all the resulting transformations actually leave eq. (2.93) invariant when they are combined with proper shifts of the $\mathcal{Y}_{ijk}$. The symmetrized triple γ -traces of eq. (2.101) read

$$T_{(ij}\gamma_k)\delta\mathcal{S} = T_{(ij|}\left[(D-2)\Omega_{|k)} + 2S^l{}_{|k)}\Omega_l\right] - \gamma^l T_{(ij}\gamma_k)\Omega_l, \quad (2.119)$$

and thus lead directly to Labastida-like constraints for the Ω_i satisfying the condition (2.103). In a similar fashion, the symmetrized triple γ -traces of eq. (2.104) read

$$\begin{aligned} T_{(ij}\gamma_k)\delta\mathcal{S} = & \gamma_{(i|}\left[D\Omega_{|jk)} + S^l{}_{|j}\Omega_{k)l} - 2S^l{}_{|j}\tilde{\Omega}_{k)l}\right] \\ & + \gamma^l T_{(ij|}\left[\Omega_{|k)l} - 2\left(S^m{}_{|k)}\tilde{\Omega}_{lm} + S^m{}_l\tilde{\Omega}_{|k)m}\right)\right] \\ & + 2\gamma^l T_{(ij|}\left[(D-3)\tilde{\Omega}_{|k)l} - \left(S^m{}_{|k)}\tilde{\Omega}_{lm} - S^m{}_l\tilde{\Omega}_{|k)m}\right)\right] \\ & + \eta^{lm}T_{(ij}\gamma_k)\Omega_{lm} + \gamma^{lm}T_{(ij}\gamma_k)\tilde{\Omega}_{lm}, \end{aligned} \quad (2.120)$$

and when eq. (2.107) holds they lead again to Labastida-like constraints for the Ω_{ij} and $\tilde{\Omega}_{ij}$ parameters. As we shall see, however, this particular set of conditions does not play a role at two families, since all solutions of eq. (2.107) correspond to fields bearing only a few space-time indices, for which the Labastida constraints are trivial. On the other hand, for the shift (2.109) the conditions (2.100) read

$$\begin{aligned} T_{(ij}\gamma_k)\delta\mathcal{S} = & \eta^{lm}T_{(ij|}\left[D\Omega_{lm;|k)} + 2S^n{}_{|k)}\Omega_{lm;n}\right] + \frac{1}{2}\gamma^{lm}T_{(ij}\Omega_k)[l;m] \\ & + \left[D(D+2)\Omega_{(ij;k)} + 2DS^l{}_{(i}\Omega_{jk);l} + (D+2)S^l{}_{(i|}\Omega_{l|j;k)} + 2S^l{}_{(i}S^m{}_j\Omega_{k)m;l}\right] \\ & - \gamma^l\gamma_{(i|}\left[D\Omega_{|jk);l} + S^m{}_{|j}\Omega_{k)m;l}\right] - \eta^{lm}\gamma^n T_{(ij}\gamma_k)\Omega_{lm;n}. \end{aligned} \quad (2.121)$$

They play a crucial role in the ensuing analysis, and reduce once more to Labastida-like constraints on these additional parameters when (2.111) holds. This becomes manifest after eliminating the $\{2,1\}$ component in the family indices from the parameters, as demanded by eq. (2.111), and making use of the rewriting in eq. (2.112).

We can now close the circle, showing that the shift symmetries with parameters satisfying Bianchi identities, Labastida-like triple γ -trace constraints and the algebraic conditions of eqs. (2.103), (2.107) and (2.111) leave the equation of motion (2.93) invariant, when they are combined with proper $\delta\mathcal{Y}_{ijk}$ shifts. Let us begin by considering the shift (2.101), that at two families affects the Rarita-Schwinger-like tensor of eq. (2.76) as follows:

$$\begin{aligned} \delta\mathcal{E} = & -\frac{1}{2}\gamma^i\left[(D-2)\Omega_i + 2S^j{}_i\Omega_j\right] - \frac{1}{12}\gamma^{ij}\gamma_{[i|}\left[(D-2)\Omega_{|j]} + 2S^k{}_{|j]} \Omega_k\right] \\ & + \frac{1}{6}\eta^{ij}\gamma^k Y_{\{2,1\}}T_{ij}\left[(D-2)\Omega_k + 2S^l{}_k\Omega_l\right] - \frac{1}{6}\eta^{ij}\gamma^k T_{(ij}\Omega_k) \\ & - \frac{1}{3}\eta^{ij}\eta^{kl}Y_{\{3,1\}}(Y_{\{2,1\}}T_{ij}\gamma_k)\Omega_l - \frac{1}{6}\eta^{ij}\gamma^{kl}Y_{\{3,1\}}(Y_{\{2,1\}}T_{ij}\gamma_k)\Omega_l \\ & + \frac{1}{12}\eta^{ij}\eta^{kl}\gamma^m Y_{\{3,2\}}(Y_{\{2,2\}}T_{ij}T_{kl})\Omega_m. \end{aligned} \quad (2.122)$$

The first three groups of terms vanish whenever eq. (2.103) holds, while at two families the others carry Young projections with more than two columns. As a result, in the variation of (2.93) they can be compensated by proper Ω -dependent shifts of the $\mathcal{Y}_{ijk}$ tensors. In particular, the contraction of all indices with invariant tensors makes it possible to factor out from all surviving terms the symmetric combinations $\eta^{(ij} \gamma^{k)}$ that accompany the $\mathcal{Y}_{ijk}$, so that one can indeed conclude that the equation of motion (2.93) is left invariant by the transformations

$$\begin{aligned}\delta \mathcal{S} &= \gamma^i \Omega_i, \\ \delta \mathcal{Y}_{ijk} &= -\frac{1}{3} \left\{ T_{(ij} \Omega_{k)} - \frac{1}{12} \gamma^l (2 \gamma_l T_{(ij} \Omega_{k)} - T_{l(i} \gamma_j \Omega_{k)}) \right. \\ &\quad \left. + \frac{1}{12} \eta^{lm} (2 T_{lm} T_{(ij} \Omega_{k)} - T_{l(i} T_{m|j} \Omega_{k)}) \right\},\end{aligned}\quad (2.123)$$

with the Ω_i satisfying the conditions (2.103). However, not all resulting solutions are relevant, since, as we repeatedly stressed, one is also to preserve eq. (2.94).

As we shall see, for the simple shifts (2.104) that solve eq. (2.107) the invariance of the field equation (2.93) can be verified almost by inspection. However, in order to better understand the behavior of the equation for $\bar{\psi}$ under Weyl-like transformations it is interesting to look at the variation of the Rarita-Schwinger-like tensor induced by generic shifts of the type (2.104):

$$\begin{aligned}\delta \mathcal{E} &= -\frac{1}{2} \eta^{ij} \left[D \Omega_{ij} + S^k_{(i} \Omega_{j)k} - 2 S^k_{(i} \tilde{\Omega}_{j)k} \right] \\ &+ \frac{1}{6} \gamma^{ij} \left[(D-4)(D-3) \tilde{\Omega}_{ij} - (2D-7) S^k_{[i} \tilde{\Omega}_{j]k} - \frac{1}{2} S^k_{[i} \Omega_{j]k} + 2 S^k_{[i} S^l_{|j]} \tilde{\Omega}_{kl} \right] \\ &- \frac{1}{6} \eta^{ij} \gamma^k Y_{\{2,1\}} \gamma_{(i} \left[D \Omega_{|j)k} + (S^l_{|j)} \Omega_{kl} + S^l_k \Omega_{|j)l} \right) - 2 (S^l_{|j)} \tilde{\Omega}_{kl} + S^l_k \tilde{\Omega}_{|j)l} \Big] \\ &+ \frac{1}{6} \eta^{ij} \eta^{kl} Y_{\{2,2\}} T_{ij} \left[D \Omega_{kl} + S^m_{(k} \Omega_{l)m} - 2 S^m_{(k} \tilde{\Omega}_{l)m} \right] \\ &- \frac{1}{6} \eta^{ij} \gamma^k \gamma_{(i} \Omega_{jk)} - \frac{1}{12} \eta^{ij} \eta^{kl} T_{(ij} \Omega_{kl)} - \frac{1}{2} \eta^{ij} \gamma^{kl} Y_{\{3,1\}} T_{ij} \tilde{\Omega}_{kl} - \frac{1}{12} \eta^{ij} \gamma^{kl} Y_{\{3,1\}} \gamma_{kl} \Omega_{ij} \\ &+ \frac{D}{9} \eta^{ij} \gamma^{kl} Y_{\{3,1\}} \left(2 T_{ij} \tilde{\Omega}_{kl} - T_{k(i} \Omega_{j)l} \right) - \frac{2}{3} \eta^{ij} \gamma^{kl} Y_{\{3,1\}} (Y_{\{2,1\}} T_{ij} S^m_k) \tilde{\Omega}_{lm} \\ &+ \frac{1}{9} \eta^{ij} \eta^{kl} \gamma^m Y_{\{3,2\}} \left[6 (Y_{\{2,1\}} T_{ij} \gamma_k) \tilde{\Omega}_{lm} - \left(2 T_{ij} \gamma_{(k} \tilde{\Omega}_{l)m} - \gamma_{(i} T_{j)(k} \tilde{\Omega}_{l)m} \right) \right].\end{aligned}\quad (2.124)$$

The last three lines contain terms that can be compensated by shifts of the $\mathcal{Y}_{ijk}$, while the two-column projected terms are to vanish by themselves. Although in this case the conditions are more complicated than those emerging from the Bianchi identities, one can verify that all these terms vanish whenever eq. (2.107) holds. In particular, the first, third and fourth groups of terms involve only rewritings of the second line of eq. (2.105), while the second group can be disposed of expressing everything in terms of $\tilde{\Omega}$. Although one cannot exclude that these complicated conditions admit other solutions aside from those of eq. (2.107), as we pointed out at the beginning of this section we are only interested in “physical” solutions, that as such also preserve the Bianchi identities (2.99) and the Labastida-like constraints (2.100).

The last class of relevant shift transformations appears in eq. (2.109). We already pointed out that the conditions (2.111) imply that the $\{2,1\}$ family-index component of the parameters vanishes, and this suffices to prove the invariance of the field equation (2.93). In fact, eq. (2.112) suffices to replace all terms containing the S^i_j operators with others where only the Ω parameters appear. As a result, the variation of the Rarita-Schwinger-like tensor can be

manifestly compensated by suitable $\delta \mathcal{Y}_{ijk}$ transformations, since all combinations of γ -traces of the parameters only admit projections in their family indices with at least three columns. This concludes the proof that the relevant shift symmetries of the Bianchi identities give rise to shift symmetries of the field equation (2.93). Furthermore, the direct variation of the field equation shows that the Bianchi identities greatly simplify the analysis of the relevant conditions, that could prove a non-trivial technical task otherwise.

2.3.1 WEYL-LIKE SYMMETRIES

In the first part of this section we identified the conditions that forbid the direct reduction of the field equations to the Labastida form, and in the following we would like to present a list of the types of two-family fields exhibiting this phenomenon. As a first step, however, we shall display Weyl-like transformations for the ψ fields that can gauge away the undetermined quantities. Let us begin from the simplest class of fields with pathological equations of motion, those admitting the shift symmetries (2.101) with parameters satisfying eq. (2.103). A Weyl-like shift of ψ of the form

$$\delta \psi = \gamma^i \Theta_i \quad (2.125)$$

gives rise to the corresponding variation of the Fang-Fronsdal-Labastida tensor,

$$\delta \mathcal{S} = -i \partial^k \left[(D-2) \Theta_k + 2 S^l{}_k \Theta_l \right] - \gamma^k \mathcal{S}(\Theta_k). \quad (2.126)$$

Eq. (2.103) is an algebraic condition that depends only on the space-time dimension D and on the tensorial structure of the parameters involved, and whenever it admits non-trivial solutions eq. (2.126) reduces to

$$\delta \mathcal{S} = -\gamma^k \mathcal{S}(\Theta_k), \quad (2.127)$$

while the variation of the Labastida constraints or, in the unconstrained case, the field equations (2.94) for the Lagrange multipliers, reduce to

$$T_{(ij} \gamma_k) \delta \psi = \gamma^l T_{(ij} \gamma_k) \Theta_l. \quad (2.128)$$

Notice that the $\mathcal{S}(\Theta_i)$ satisfy the conditions

$$(D-2) \mathcal{S}(\Theta_i) + 2 S^j{}_i \mathcal{S}(\Theta_j) = 0, \quad (2.129)$$

since the $S^i{}_j$ commute with the Fang-Fronsdal-Labastida $\mathcal{S}$ operators. Furthermore, they satisfy the constraints (2.100) provided the shifts (2.125) are supplemented by the conditions

$$T_{(ij} \gamma_k) \Theta_l = 0, \quad (2.130)$$

that are needed to preserve the Labastida constraints (2.18) on ψ . These conditions also imply that the $\mathcal{S}(\Theta_i)$ satisfy the Bianchi identities, so that one can use them to gauge away the corresponding undetermined quantities in the field equations.

In a similar fashion a shift of the form

$$\delta \psi = \eta^{ij} \Theta_{ij} + \gamma^{ij} \tilde{\Theta}_{ij} \quad (2.131)$$

gives rise to the following variation of the Labastida tensor:

$$\begin{aligned} \delta \mathcal{S} = & -i \partial^j \gamma^k \left[\Theta_{jk} - 2 S^l{}_{(j} \tilde{\Theta}_{k)l} \right] - 2i \partial^j \gamma^k \left[(D-3) \tilde{\Theta}_{jk} - S^l{}_{[j} \tilde{\Theta}_{k]l} \right] \\ & + \eta^{ij} \mathcal{S}(\Theta_{ij}) + \gamma^{ij} \mathcal{S}(\tilde{\Theta}_{ij}). \end{aligned} \quad (2.132)$$

Even in this case, whenever the Lorentz structure of the gauge field ψ guarantees that the conditions (2.107) be satisfied, one can properly choose the parameters Θ_{ij} and $\tilde{\Theta}_{ij}$ so that

$$\begin{aligned}\delta \mathcal{S} &= \eta^{ij} \mathcal{S}(\Theta_{ij}) + \gamma^{ij} \mathcal{S}(\tilde{\Theta}_{ij}), \\ T_{(ij} \gamma_k) \delta \psi &= \eta^{lm} T_{(ij} \gamma_k) \Theta_{lm} + \gamma^{lm} T_{(ij} \gamma_k) \tilde{\Theta}_{lm},\end{aligned}\tag{2.133}$$

with the $\mathcal{S}(\Theta_{ij})$ and $\mathcal{S}(\tilde{\Theta}_{ij})$ satisfying the conditions of eq. (2.107). As in the previous case, these quantities also satisfy the Bianchi identities once one requires that $\delta \psi$ preserve the Labastida constraints. Therefore, further undetermined quantities can be shifted away, as in the previous example, by the Weyl-like gauge transformation (2.131), with parameters satisfying the conditions (2.107) and the corresponding Labastida-like triple γ -trace constraints.

The last class of shift symmetries that we identified involves three γ -matrices, and one can dispose of it via the Weyl-like gauge transformations

$$\delta \psi = \eta^{ij} \gamma^k \Theta_{ij;k},\tag{2.134}$$

that affect the Fang-Fronsdal-Labastida tensor according to

$$\begin{aligned}\delta \mathcal{S} &= -i \eta^{ij} \partial^k \left[(D-2) \Theta_{ij;k} + \Theta_{k(i;j)} + 2 S^l_k \Theta_{ij;l} \right] - \frac{i}{2} \gamma^{ij} \partial^k \Theta_{k[i;j]} \\ &\quad - \eta^{ij} \gamma^k \mathcal{S}(\Theta_{ij;k}).\end{aligned}\tag{2.135}$$

Again, whenever the conditions (2.111) hold, one can factor out the invariant tensors, and a shift of $\mathcal{S}$ emerges that is capable of removing the undetermined quantities. Once more, the leftover Weyl parameters $\hat{\Theta}_{lmn}$ are fully symmetric in their family indices and must satisfy triple γ -trace constraints of the form

$$T_{(ij} \gamma_k) \hat{\Theta}_{lmn} = 0,\tag{2.136}$$

in order to leave eqs. (2.94) and the Bianchi identities invariant.

These observations actually suggest an alternative starting point to classify the fields whose equations of motion do not reduce directly to the Labastida form (2.92): one can classify the Weyl-like shifts of ψ , that by construction also leave the Bianchi identities invariant. We defer further comments to Section 3.4, where we shall adopt this convenient viewpoint to study the reduction procedure in the general N -family case.

We can now take a closer look at the previous conditions. As we anticipated, and as we are going to see in detail shortly, while eqs. (2.103), (2.107) and (2.111) admit non-trivial solutions, eqs. (2.116) do not. In all these cases one is facing eigenvalue problems for the S^i_j operators that can identify Weyl-like gauge transformations, but whose solutions are to be subjected by further conditions arising from triple γ -trace constraints. At two families the eigenvalue problems greatly simplify, since one can resort to the $gl(2)$ techniques already described in [2] for Bose fields. In fact, turning to the suggestive notation

$$L_+ = S^1_2, \quad L_- = S^2_1, \quad L_3 = \frac{1}{2} (S^1_1 - S^2_2),\tag{2.137}$$

one readily recovers the angular momentum algebra

$$[L_3, L_{\pm}] = \pm L_{\pm}, \quad [L_+, L_-] = 2L_3,\tag{2.138}$$

while looking at their definition in eq. (A.13) one can recognize that S^1_1 and S^2_2 act diagonally on generic spinor-tensors ψ of rank (s_1, s_2) , with

$$S^1_1 \psi = s_1 \psi, \quad S^2_2 \psi = s_2 \psi. \quad (2.139)$$

In order to proceed, it is thus convenient to split the problems into combinations of simpler ones, expanding in bases on which the “total angular momentum” $\mathbf{L}^2$ acts diagonally. For instance, a gauge field ψ of rank (s_1, s_2) can be decomposed according to

$$\psi = \sum_{n=0}^{s_2} Y_{\{s_1+s_2-n, n\}} \psi \equiv \sum_{n=0}^{s_2} \psi^{\{s_1+s_2-n, n\}}, \quad (2.140)$$

where the various components are characterized by the same L_3 eigenvalue, that is fixed by the rank of ψ to be

$$m = \frac{s_1 - s_2}{2}. \quad (2.141)$$

The terms of the sum (2.140) corresponding to Young projections of the type $Y_{\{s_1+s_2-n, n\}}$ can be related to null eigenvectors of L_+ via the standard descent relations

$$\psi^{\{s_1+s_2-n, n\}} = (L_-)^{s_2-n} \hat{\psi}^{\{s_1+s_2-n, n\}}, \quad L_+ \hat{\psi}^{\{s_1+s_2-n, n\}} = 0, \quad (2.142)$$

so that they are all eigenvectors of $\mathbf{L}^2$, but in general with different eigenvalues. The $\hat{\psi}$ lie at the tips of the chains and are clearly $gl(D)$ *irreducible* in their vector indices on account of eq. (A.13). In fact, they are annihilated by L_+ , so that a vanishing result obtains if one tries to extend the symmetrization beyond the first family. On the other hand, the lower members of the chains in the decomposition (2.140) are built acting on the $\hat{\psi}$ with powers of L_- , and as a result their symmetry is not manifest. However, they are still eigenvectors of $Y_{\{s_1+s_2-n, n\}}$, that commutes with L_- simply because the permutation group acts irreducibly within individual Y eigenspaces. All in all, the operator

$$L_- L_+ = \mathbf{L}^2 - L_3^2 - L_3 \quad (2.143)$$

is diagonal when acting on this basis, whose members $\psi^{\{s_1+s_2-n, n\}}$ are characterized by the *fixed* value (2.141) for m and by a *range* of values of ℓ ,

$$\ell = \frac{s_1 + s_2}{2} - n, \quad \frac{s_1 - s_2}{2} \leq \ell \leq \frac{s_1 + s_2}{2}, \quad (2.144)$$

so that

$$L_- L_+ \psi^{\{s_1+s_2-n, n\}} = (n - s_1 - 1)(n - s_2) \psi^{\{s_1+s_2-n, n\}}. \quad (2.145)$$

In the rest of this section we shall make extensive use of these tools to analyze the algebraic conditions that we previously identified, and we shall conclude with some comments on the solutions and with the identification of a special subclass of fields whose Lagrangians vanish identically, as is the case for the Rarita-Schwinger field in two dimensions. The discussion of the eigenvalue problems for the S^i_j operators is somewhat technical, and the reader may wish to skip the following pages and move directly to the tables that collect their outcome. Their two columns are labeled by $\mathcal{S}$ and $\mathcal{S}'$ and collect the types of fields whose equations of motion leave some combinations of single or double γ -traces of $\mathcal{S}$ undetermined. In fact, the solutions of the eigenvalue problem (2.111) only act on the symmetrized triple γ -traces of $\mathcal{S}$, and therefore will be discarded.

ANALYSIS OF EQ. (2.103)

Using the notation of eq. (2.137) and taking into account the previous considerations, for a two-family field ψ of rank (s_1, s_2) the conditions (2.103) become

$$\begin{aligned} (D + 2s_1 - 4)\Omega_1 + 2L_- \Omega_2 &= 0, \\ (D + 2s_2 - 4)\Omega_2 + 2L_+ \Omega_1 &= 0. \end{aligned} \quad (2.146)$$

Barring a couple of degenerate cases to be treated separately, this system can be solved by direct substitution. Thus, expressing Ω_2 in terms of Ω_1 and decomposing it into irreducible $gl(D)$ components, using the notation of eq. (2.140) leads to

$$(D + 2n - 4)(D + 2s_1 + 2s_2 - 2n - 4)(\Omega_1)^{\{s_1+s_2-n-1, n\}} = 0, \quad 0 \leq n \leq s_2. \quad (2.147)$$

The second factor vanishes only in the degenerate cases where the direct substitution is not possible. These correspond to the two-dimensional gravitino, that we have already discussed, and to another two-dimensional field, the $\{1, 1\}$ irreducible spinor-tensor, for which however $\mathcal{S}$ vanishes identically when expressed in terms of ψ . On the other hand, the first factor vanishes in $D = 2$ for $n = 1$ and in $D = 4$ for $n = 0$, but for all $s_1 \geq 2$ and $s_2 \geq 2$. These solutions are to be subjected to the double trace conditions

$$T_{(ij} \gamma_k) \Omega_l = 0, \quad (2.148)$$

that, as we shall see shortly, eliminate some of them. The final results are collected in the first column of Table 1.

Each value of n selects a particular irreducible component for the vector indices carried by the parameters, and consequently not all irreducible components of the kinetic spinor-tensor (or of the field, if one takes eq. (2.125) as a starting point) can be affected. In particular, for these two classes of solutions the resulting shifts of $\mathcal{S}$,

$$\delta \mathcal{S} = \gamma^1 \Omega_1 - \frac{2}{D + 2s_2 - 4} \gamma^2 L_+ \Omega_1, \quad (2.149)$$

satisfy the conditions

$$\begin{aligned} L_- L_+ \delta \mathcal{S} &= (s_1 - 1)(s_2 - 2) \delta \mathcal{S}, & D = 2, \quad n = 1, \\ L_- L_+ \delta \mathcal{S} &= s_1(s_2 - 1) \delta \mathcal{S}, & D = 4, \quad n = 0. \end{aligned} \quad (2.150)$$

Comparing these eigenvalues with eq. (2.145), one can see that eq. (2.149) affects only the $\{s_1 + s_2 - 2, 2\}$ component of $\mathcal{S}$ in $D = 2$ and its $\{s_1 + s_2 - 1, 1\}$ component in $D = 4$. However, the Weyl-like transformation (2.149) corresponding to the first solution in (2.150) disappears if the parameter is subject to the condition (2.148), but the corresponding undetermined quantity in the field equation $E_{\bar{\psi}}$ disappears as well on account of $E_{\bar{\zeta}}$. The $\{2, 2\}$ field in two dimensions is a notable exception, since it is not subject to any Labastida constraints, but this case is also highly degenerate, since the corresponding Lagrangian vanishes identically even before expressing $\mathcal{S}$ in terms of ψ . All these results can be obtained analyzing explicitly in components the various expressions, and the irreducible field components that are left undetermined are collected in the first column of Table 2.

ANALYSIS OF EQ. (2.107)

Moving on to eq. (2.107), one should notice that at two families the antisymmetric $\tilde{\Omega}_{ij}$ reduce to the single $\tilde{\Omega}_{12}$. For a rank- (s_1, s_2) field, the first set of conditions in eq. (2.107) then relates $\tilde{\Omega}_{12}$ to the symmetric parameters according to

$$\begin{aligned}\Omega_{11} &= 4 L_- \tilde{\Omega}_{12}, \\ \Omega_{12} &= -2 (s_1 - s_2) \tilde{\Omega}_{12}, \\ \Omega_{22} &= -4 L_+ \tilde{\Omega}_{12},\end{aligned}\tag{2.151}$$

while the second condition becomes

$$(D + s_1 + s_2 - 5) \tilde{\Omega}_{12} = 0.\tag{2.152}$$

Eq. (2.152) admits non-trivial solutions only for $s_1 = 2$ and $s_2 = 1$ in $D = 2$ and for $s_1 = 1$ and $s_2 = 1$ in $D = 3$, and these are indeed the only cases when non-trivial Ω_{ij} are admissible. In both cases only symmetric and hooked irreducible field components are available, and the pathological behavior clearly concerns the hooked components, since we know already that a gravitino in $D = 2$ is the only degenerate symmetric example. At any rate, it is also possible to confirm this expectation by a direct calculation similar to those of eq. (2.150). In this case eq. (2.100) does not impose any further conditions, since these fields are too simple and are not subject to any Labastida constraints. However, one is to discard the $\{2, 1\}$ case, that is degenerate: in this case the Weyl-like transformation (2.131) leaves $\mathcal{S}$ invariant, so that the γ -traces that are left undetermined by the Lagrangian field equation vanish identically. Again, this can be seen analyzing explicitly these expressions in components. The final results are collected in the second columns of Tables 1 and 2.

ANALYSIS OF EQ. (2.111)

As we already noticed, the second condition in eq. (2.111) annihilates the $\{2, 1\}$ combinations of the shift parameters, so that one can conveniently rewrite the relations directly in terms of the fully symmetric $\hat{\Omega}_{ijk}$. This was done in eq. (2.112), whose first line contains a square system of equations for the four $\hat{\Omega}_{ijk}$ that are available in the two-family case:

$$\begin{aligned}(D + 2s_1 - 6) \hat{\Omega}_{111} + 2L_- \hat{\Omega}_{112} &= 0, \\ 2L_+ \hat{\Omega}_{111} + (3D + 4s_1 + 2s_2 - 10) \hat{\Omega}_{112} + 4L_- \hat{\Omega}_{122} &= 0, \\ 4L_+ \hat{\Omega}_{112} + (3D + 2s_1 + 4s_2 - 10) \hat{\Omega}_{122} + 2L_- \hat{\Omega}_{222} &= 0, \\ 2L_+ \hat{\Omega}_{122} + (D + 2s_2 - 6) \hat{\Omega}_{222} &= 0.\end{aligned}\tag{2.153}$$

One can solve this system after multiplying the equations by suitable powers of L_- in order to convert them into others characterized by the same value of m . The operation corresponds to performing some symmetrizations, in order that all equations finally carry like sets of symmetric space-time indices. However, these sets can be displaced in different positions on the same parameter, so that for instance for a rank- $(4, 2)$ field ψ the two distinct combinations

$$\hat{\Omega}^{(112)}_{\mu_1 \mu_2, \nu} \quad \text{and} \quad \hat{\Omega}^{(112)}_{\nu (\mu_1, \mu_2)}\tag{2.154}$$

can appear. This problem can be solved projecting the result into irreducible $gl(D)$ components, since at two families the various combinations become proportional for any given component. In

our algebraic language, this means that the resulting expressions can be manipulated in order to leave only the diagonal L_-L_+ operator acting on the various components or on their L_- -shifted analogues. Suppressing for brevity the superscript introduced in eq. (2.145) to identify the irreducible components, when all parameters admit the $\{s_1 + s_2 - n - 3, n\}$ Young projection the system (2.153) reduces to the following chain of square systems of equations labeled by n :

$$\begin{aligned} (D + 2s_1 - 6) \widehat{\Omega}_{111} + 2L_- \widehat{\Omega}_{112} &= 0, \\ 2(n - s_1 + 2)(n - s_2) \widehat{\Omega}_{111} + (3D + 4s_1 + 2s_2 - 10) L_- \widehat{\Omega}_{112} + 4L_-^2 \widehat{\Omega}_{122} &= 0, \\ 4(n - s_1 + 1)(n - s_2 + 1) L_- \widehat{\Omega}_{112} + (3D + 2s_1 + 4s_2 - 10) L_-^2 \widehat{\Omega}_{122} + 2L_-^3 \widehat{\Omega}_{222} &= 0, \\ 2(n - s_1)(n - s_2 + 2) L_-^2 \widehat{\Omega}_{122} + (D + 2s_2 - 6) L_-^3 \widehat{\Omega}_{222} &= 0. \end{aligned} \quad (2.155)$$

Their determinants read

$$\begin{aligned} (D + 2n - 2)(3D + 4(s_1 + s_2) - 2n - 18)(D + 2(s_1 + s_2) - 2n - 6) \\ \times (3D + 2(s_1 + s_2) + 2n - 14), \end{aligned} \quad (2.156)$$

and vanish if and only if $D = 2$ and $n = 0$, since $0 \leq n \leq s_2$ and the whole system is available only if $s_1 \geq 3$ and $s_2 \geq 3$, and in the remaining degenerate cases no new solutions emerge. In $D = 2$, the $n = 0$ system has rank three and admits the solution

$$\begin{aligned} \widehat{\Omega}_{111} &= -\frac{1}{(s_1 - 2)(s_1 - 1)s_1} L_-^3 (\widehat{\Omega}_{222})^{\{s_1 + s_2 - 3, 0\}}, \\ \widehat{\Omega}_{112} &= \frac{1}{(s_1 - 1)s_1} L_-^2 (\widehat{\Omega}_{222})^{\{s_1 + s_2 - 3, 0\}}, \\ \widehat{\Omega}_{122} &= -\frac{1}{s_1} L_- (\widehat{\Omega}_{222})^{\{s_1 + s_2 - 3, 0\}}, \end{aligned} \quad (2.157)$$

where the fully symmetric $(\widehat{\Omega}_{222})^{\{s_1 + s_2 - 3, 0\}}$ is also the only non-vanishing component of $\widehat{\Omega}_{222}$. Actually, this solution also satisfies the second condition in eq. (2.112), but disappears if one imposes the triple γ -trace constraints

$$T_{(ij} \gamma_{k)} \widehat{\Omega}_{lmn} = 0, \quad (2.158)$$

so that it must be rejected.

More in detail, this can be seen identifying the single irreducible component of the fields that is affected by these transformations, since only $n = 0$ parameters are involved, that are fully symmetric in their space-time indices. Substituting the solution (2.157) in eq. (2.109) and applying L_-L_+ one finds

$$L_-L_+ \delta \mathcal{S} = (s_1 - 2)(s_2 - 3) \delta \mathcal{S}, \quad D = 2, \quad n = 0, \quad (2.159)$$

so that only the $\{s_1 + s_2 - 3, 3\}$ -component of $\mathcal{S}$ is affected. This type of terms can now be analyzed in components, showing that indeed the remaining independent parameter satisfying eq. (2.158) cannot lead to a non-trivial shift of the form (2.157). This concludes the classification of the $gl(D)$ irreducible fields whose equations of motion do not reduce to the Labastida form, and the results are summarized in Table 2.

	$\mathcal{S}$		$\mathcal{S}'$	
D	s_1	s_2	s_1	s_2
2	1	0		
2	2	2		
3			1	1
4	≥ 1	≥ 1		

Table 1: $gl(D)$ -reducible fields with Weyl-like symmetries

ANALYSIS OF EQ. (2.116)

We can finally show that eq. (2.116) admits no solutions at two families. Its decomposition into irreducible Young components for the family indices is more intricate than for the previous conditions, and in order to obtain the result it is convenient to proceed via some alternative manipulations. For instance, antisymmetrizing eq. (2.116) in the (k, l) indices leads to

$$2(D-3)\tilde{\Omega}_{ij;kl} + \tilde{\Omega}_{k(i;j)l} - \tilde{\Omega}_{l(i;j)k} - 2S^m_{[k|\tilde{\Omega}_{ij;|l]m}} = 0, \quad (2.160)$$

that at two families reduces to three simple conditions for the three available $\tilde{\Omega}_{ij;kl}$:

$$\begin{aligned} 2(D+s_1+s_2-6)\tilde{\Omega}_{11;12} &= 0, \\ 2(D+s_1+s_2-6)\tilde{\Omega}_{12;12} &= 0, \\ 2(D+s_1+s_2-6)\tilde{\Omega}_{22;12} &= 0. \end{aligned} \quad (2.161)$$

These immediately imply that no solutions exist for $D > 2$, since the $\tilde{\Omega}_{ij,kl}$ parameters are only available if $s_1 + s_2 \geq 4$ and the $\Omega_{ij,kl}$ are directly related to them via

$$\Omega_{ij;kl} = \frac{1}{2} \left(S^m_{(i|\tilde{\Omega}_{kl;|j)m}} + S^m_{(k|\tilde{\Omega}_{ij;|l)m}} \right), \quad (2.162)$$

that emerges symmetrizing eq. (2.116) in the (k, l) indices and also enforcing the symmetry under interchanges of the (i, j) and (k, l) pairs. Still, in principle a $\{3, 1\}$ or a $\{2, 2\}$ field in two dimensions could admit non-trivial Weyl-like symmetries, but in these cases all parameters are scalars and the equations obtained only symmetrizing (2.116) in (k, l) simply reduce to

$$\begin{aligned} \Omega_{11,12} &= \tilde{\Omega}_{11;12}, \\ \Omega_{11,12} &= -\tilde{\Omega}_{11;12}, \end{aligned} \quad (2.163)$$

for the $\{3, 1\}$ case and similarly to

$$\begin{aligned} \Omega_{11,22} &= \tilde{\Omega}_{12;12}, \\ \Omega_{11,22} &= -\tilde{\Omega}_{12;12}, \end{aligned} \quad (2.164)$$

for the $\{2, 2\}$ case. In conclusion, the conditions of eq. (2.116) do not admit any solution, as we anticipated.

	$\mathcal{S}$		$\mathcal{S}'$	
D	s_1	s_2	s_1	s_2
2	1	0		
2	2	2		
3			1	1
4	s	1		

Table 2: $gl(D)$ -irreducible fields with Weyl-like symmetries

2.3.2 LAGRANGIANS THAT ARE TOTAL DERIVATIVES

In the preceding pages we have classified the fields whose equations of motion do not directly reduce to the Labastida form (2.92). We have seen that this does not happen for $D > 4$, and that new Weyl-like symmetries emerge that can gauge away the undetermined quantities. We have also seen that some solutions of the eigenvalue equations, that identify symmetries of eq. (2.93), are not compatible with the triple γ -trace conditions (2.94), and therefore must be discarded¹⁸.

The paradigmatic example provided by the two-dimensional gravitino suggests that those symmetries could well signal the emergence of Lagrangians that are at most total derivatives, and indeed the cases listed in Table 2 correspond to representations carrying no degrees of freedom¹⁹. On the other hand, the example of the two-dimensional gravitino and its role in the dynamics of superstrings should make it clear that models of this type are nonetheless potentially quite rich.

The possibility that the pathological cases be characterized by vanishing actions can be verified directly at the level of the field equations, that should vanish identically for this type of models. We have seen how, after relating the $\mathcal{Y}_{ijk}$ tensors to the kinetic spinor-tensor $\mathcal{S}$, one ends up with a combination of $\mathcal{S}$ and its γ -traces. If $\mathcal{S}$ does not vanish identically, this combination can only vanish if it coincides with one of its vanishing $o(D)$ components, and when this is the case the remaining $o(D)$ components can be clearly shifted at will and a Weyl-like symmetry emerges. This is precisely what happens for the gravitino field in two dimensions, where the Rarita-Schwinger tensor coincides with the vanishing γ -traceless projection of $\mathcal{S}$, and in principle it can also happen for other fields in low space-time dimensions. It is thus interesting to try and understand whether all fields appearing in Table 2 behave exactly like a gravitino in $D = 2$ or, consistently with the expectation that one could build from the corresponding analysis performed in [2] for bosons, whether some Weyl-invariant models of a different type exist for mixed-symmetry fermions. First of all, one can observe that the example provided by the gravitino is not so exotic, since a whole class of similar theories actually exists. In fact, the

¹⁸For Bose fields the analogues of these considerations involve double trace constraints on Weyl-like parameters, that were correctly identified in [2]. However, we failed to notice that they also eliminate some of the solutions, namely the reducible $D = 2$ cases ($s_1 \geq 3, s_2 \geq 3$) of Table 1 of [2], or the corresponding irreducible cases $\{s, 3\}$ of Table 2 of [2].

¹⁹If an $o(n)$ bosonic representation disappears for a given n , the fermionic representation with the same tensorial structure disappears as well. The γ -tracelessness condition for $o(n)$ spinor-tensors is in fact stronger than the tracelessness condition for bosonic irreducible representations, so that the tensorial contribution to the dimension is always smaller in the fermionic case. One can verify this also looking at eqs. (A.38) and (A.39) for these dimensions. On the other hand, a general result of representation theory for $o(n)$ groups states that (see, for instance, the first reference in [33], §10-6), if the total number of boxes in the first two columns of a tableau exceeds n , the corresponding traceless tensor vanishes.

rewriting (2.98) of the Rarita-Schwinger Lagrangian suggests to consider the class of Lagrangians

$$\mathcal{L} = i \frac{(-1)^{\frac{q(q-1)}{2}}}{2q!} \bar{\psi}_{\mu_1 \dots \mu_q} \gamma^{\mu_1 \dots \mu_q \lambda \nu_1 \dots \nu_q} \partial_\lambda \psi_{\nu_1 \dots \nu_q} + \text{h.c.} \quad (2.165)$$

for one-column (*fully antisymmetric*) spinor-tensors²⁰. In Section 3.1 we shall derive the Lagrangians for arbitrary mixed-symmetry Fermi fields, and the reader can verify that the two presentations actually coincide, since the fields in this class are particularly simple and are not subject to any constraints. Moreover, in Section 2 we already saw that the structure of the Lagrangians is fixed by the requirement of gauge invariance, and (2.165) is indeed manifestly gauge invariant, simply because the gradients acting on the gauge parameters are antisymmetrized with the derivative that is already present. Therefore, spinor-tensor fields carrying q antisymmetric vector indices define in general a class of interesting analogs of the two-dimensional Rarita-Schwinger system for $q \leq D \leq 2q$. And indeed their only two-family representative, a $\{1, 1\}$ irreducible field, appears in Table 2 for $D = 3, 4$, while we left out the $D = 2$ case, that is somewhat degenerate since the corresponding $\mathcal{S}$ vanishes identically. As we anticipated, another interesting example is provided by an irreducible $\{2, 2\}$ field in two dimensions, whose Rarita-Schwinger-like tensor $\mathcal{E}$ vanishes identically even before expressing $\mathcal{S}$ in terms of ψ .

Not all models listed in Table 2, however, have Lagrangians that are total derivatives. A rather simple counterexample is provided by a $\{2, 1\}$ -irreducible $gl(D)$ spinor-tensor in $D = 4$: one can verify that the Rarita-Schwinger-like tensor (2.76) does not vanish in this case. This suffices to reach our conclusion, since the $\{2, 1\}$ fields are intrinsically unconstrained, simply because the combinations of γ -traces that identify the Labastida constraints are not available for them. Deferring to future work a more detailed analysis, we can already conclude that for mixed-symmetry Fermi fields the emergence of Weyl-like symmetries is not necessarily accompanied by vanishing actions, in analogy with the behavior of mixed-symmetry Bose fields discussed in [2].

2.4 AN EXAMPLE: REDUCIBLE SPINOR-TENSOR FIELDS OF RANK $(3, 2)$

In the previous sections we presented Lagrangians and field equations for general two-family spinor-tensor fields. Our concise notation made it possible to discuss simultaneously results that apply to all fields of this type, independently of their actual Lorentz labels, but the presentation is possibly less palatable for the reader than more conventional ones. We thus deem it appropriate, in analogy with what we did for Bose fields in [2], to dwell upon the explicit space-time structure of a relatively simple and yet very instructive example, a *reducible* spinor-tensor of rank $(3, 2)$, whose explicit space-time form is $\psi_{\mu_1 \mu_2 \mu_3, \nu_1 \nu_2}$.

2.4.1 LAGRANGIAN AND FIELD EQUATIONS

As for the bosonic examples presented in [2], in this section we shall resort to a shorthand notation to denote traces so that, for instance,

$$\begin{aligned} \eta^{\mu_1 \mu_2} \psi_{\mu_1 \mu_2 \mu_3, \nu_1 \nu_2} &\equiv \psi'^{\mu}_{\mu_3, \nu_1 \nu_2}, \\ \eta^{\mu_1 \nu_1} \psi_{\mu_1 \mu_2 \mu_3, \nu_1 \nu_2} &\equiv \psi^{\hat{\nu}}_{\mu_2 \mu_3, \nu_2}. \end{aligned} \quad (2.166)$$

²⁰Similar types of Lagrangians were recently discussed in [34], that appeared while this paper was being typed.

The (3, 2) example has the virtue of being relatively simple and yet well structured, since the two distinct gauge parameters that characterize generic two-family fields are already present in this case, so that the explicit gauge variation of the field reads

$$\delta \psi = \partial^1 \epsilon_1 + \partial^2 \epsilon_2 \equiv \partial_{(\mu_1} \epsilon^{(1)}_{\mu_2 \mu_3), \nu_1 \nu_2} + \partial_{(\nu_1} \epsilon^{(2)}_{\mu_1 \mu_2 \mu_3, | \nu_2)} . \quad (2.167)$$

The two Ψ_i compensators that accompany the two gauge parameters in the unconstrained theory are

$$\begin{aligned} \Psi_1 &\equiv \Psi^{(1)}_{\mu_1 \mu_2, \nu_1 \nu_2} , \\ \Psi_2 &\equiv \Psi^{(2)}_{\mu_1 \mu_2 \mu_3, \nu} , \end{aligned} \quad (2.168)$$

but, as we have seen in general, they enter the construction only via their symmetrized γ -traces. A generic two-family fermionic field makes use of three different combinations of this sort, a fact that is fully visible in the present example, where they take the form

$$\begin{aligned} \xi_{11} &= \gamma_1 \Psi_1 \equiv \xi^{(1)}_{\mu, \nu_1 \nu_2} , \\ \xi_{12} &= \frac{1}{2} (\gamma_1 \Psi_2 + \gamma_2 \Psi_1) \equiv \xi^{(2)}_{\mu_1 \mu_2, \nu} , \\ \xi_{22} &= \gamma_2 \Psi_2 \equiv \xi^{(3)}_{\mu_1 \mu_2 \mu_3} . \end{aligned} \quad (2.169)$$

On the other hand, the four Labastida constraints that annihilate the symmetrized triple γ -traces of generic two-family spinor-tensors reduce in the present (3, 2) example to only three conditions:

$$\begin{aligned} T_{11} \gamma_1 \psi &\equiv \gamma^\lambda \psi'^\mu_{\lambda, \nu_1 \nu_2} = 0 , \\ (T_{11} \gamma_2 + 2 T_{12} \gamma_1) \psi &\equiv \gamma^\lambda (\psi'^\mu_{\mu, \nu \lambda} + 2 \psi^{\hat{\mu}}_{\lambda \mu, \nu}) = 0 , \\ (T_{22} \gamma_1 + 2 T_{12} \gamma_2) \psi &\equiv \gamma^\lambda (\psi'^\nu_{\lambda \mu_1 \mu_2} + 2 \psi^{\hat{\nu}}_{\mu_1 \mu_2, \lambda}) = 0 . \end{aligned} \quad (2.170)$$

Notice that the Labastida constraints (2.170) also imply that the symmetrized double traces of the fermionic gauge field vanish,

$$\begin{aligned} T_{11} T_{12} \psi &\equiv \psi'^{\hat{\mu}}_{\hat{\nu}} = 0 , \\ (T_{11} T_{22} + 2 T_{12} T_{12}) \psi &\equiv \psi'^{\mu \nu}_{\mu} + 2 \psi^{\hat{\mu} \hat{\nu}}_{\hat{\mu}} = 0 , \end{aligned} \quad (2.171)$$

and these conditions, in their turn, imply that all expressions containing at least five γ -traces vanish. This is counterpart, for two-family Fermi fields, of the vanishing of all triple traces for two-family Bose fields, that was widely used in [2].

As we saw in eq. (2.29), the combinations of γ -traces of the kinetic tensor $\mathcal{W}$ that have the same structure as the Labastida constraints are proportional to the $\mathcal{Z}_{ijk}$ tensors. Two hooked triple γ -traces of $\mathcal{W}$, however, are not related to the constraints, and are the analogs in this setting of the non-trivial double trace $a_{\mu\nu}$ present in the (4, 2) bosonic example discussed in [2]. In the following these expressions will be often denoted as follows:

$$\begin{aligned} (T_{11} \gamma_2 - T_{12} \gamma_1) \mathcal{W} &\equiv \gamma^\lambda (\mathcal{W}'^\mu_{\mu, \nu \lambda} - \mathcal{W}^{\hat{\mu}}_{\lambda \mu, \nu}) \equiv v_{\mu, \nu} , \\ (T_{22} \gamma_1 - T_{12} \gamma_2) \mathcal{W} &\equiv \gamma^\lambda (\mathcal{W}'^\nu_{\lambda \mu_1 \mu_2} - \mathcal{W}^{\hat{\nu}}_{\mu_1 \mu_2, \lambda}) \equiv u_{\mu_1 \mu_2} . \end{aligned} \quad (2.172)$$

In addition, the $\{2, 2\}$ -projected double trace of the gauge field,

$$(T_{11} T_{22} - T_{12} T_{12}) \psi \equiv \psi'^{\mu\nu}_{\mu} - \psi^{\hat{\mu}\hat{\nu}}_{\mu} \equiv \omega_{\mu}, \quad (2.173)$$

is not related to the constraints, and can also contribute significantly to the Lagrangian. Finally, the structure of the Labastida constraints (2.170) reflects itself in the three Lagrange multipliers,

$$\lambda_{111} \equiv \lambda^{(1)}_{\nu_1 \nu_2}, \quad \lambda_{112} \equiv \lambda^{(2)}_{\mu, \nu}, \quad \lambda_{122} \equiv \lambda^{(3)}_{\mu_1 \mu_2}, \quad (2.174)$$

that complete the field content of this model.

In an explicit space-time notation, the unconstrained Lagrangian for $\psi_{\mu_1 \mu_2 \mu_3, \nu_1 \nu_2}$ reads

$$\begin{aligned} \mathcal{L} = & \frac{1}{2} \bar{\psi}^{\mu_1 \mu_2 \mu_3, \nu_1 \nu_2} (\mathcal{E}_{\psi})_{\mu_1 \mu_2 \mu_3, \nu_1 \nu_2} \\ & + \frac{3}{2} \bar{\xi}^{(1) \mu, \nu_1 \nu_2} \Xi^{(1)}_{\mu, \nu_1 \nu_2} + 3 \bar{\xi}^{(2) \mu_1 \mu_2, \nu} \Xi^{(2)}_{\mu_1 \mu_2, \nu} + \frac{1}{2} \bar{\xi}^{(3) \mu_1 \mu_2 \mu_3} \Xi^{(3)}_{\mu_1 \mu_2 \mu_3} \\ & + \frac{3}{2} \bar{\lambda}^{(1) \nu_1 \nu_2} \mathcal{Z}^{(1)}_{\nu_1 \nu_2} + 9 \bar{\lambda}^{(2) \mu, \nu} \mathcal{Z}^{(2)}_{\mu, \nu} + \frac{9}{2} \bar{\lambda}^{(3) \mu_1 \mu_2} \mathcal{Z}^{(3)}_{\mu_1 \mu_2} + \text{h.c.}, \end{aligned} \quad (2.175)$$

with the Rarita-Schwinger-like tensor

$$\begin{aligned} \mathcal{E}_{\psi} = & \mathcal{W}_{\mu_1 \mu_2 \mu_3, \nu_1 \nu_2} - \frac{1}{2} \gamma_{(\mu_1} \gamma^{\lambda} \mathcal{W}_{\mu_2 \mu_3) \lambda, \nu_1 \nu_2} - \frac{1}{2} \gamma_{(\nu_1} \gamma^{\lambda} \mathcal{W}_{\mu_1 \mu_2 \mu_3, | \nu_2) \lambda} \\ & - \frac{1}{2} \eta_{(\mu_1 \mu_2} \mathcal{W}'^{\mu}_{\mu_3), \nu_1 \nu_2} - \frac{1}{2} \eta_{(\nu_1 | (\mu_1} \mathcal{W}'^{\hat{\mu}}_{\mu_2 \mu_3), | \nu_2)} - \frac{1}{2} \eta_{\nu_1 \nu_2} \mathcal{W}'^{\nu}_{\mu_1 \mu_2 \mu_3} \\ & + \frac{1}{6} \gamma_{(\nu_1 | (\mu_1} \gamma^{\lambda \rho} \mathcal{W}_{\mu_2 \mu_3) \lambda, | \nu_2) \rho} + \frac{1}{18} (2 \eta_{(\mu_1 \mu_2 |} \gamma_{(\nu_1 |} - \eta_{(\nu_1 | (\mu_1} \gamma_{\mu_2 |}) v_{| \mu_3), | \nu_2)} \\ & + \frac{1}{18} (2 \eta_{\nu_1 \nu_2} \gamma_{(\mu_1 |} - \gamma_{(\nu_1} \eta_{\nu_2) (\mu_1 |}) u_{| \mu_2 \mu_3)} + \frac{1}{18} (2 \eta_{\nu_1 \nu_2} \eta_{(\mu_1 \mu_2 |} - \eta_{\nu_1 (\mu_1} \eta_{\mu_2 | \nu_2)} \omega_{| \mu_3)}, \end{aligned} \quad (2.176)$$

where the symbols u , v and ω were defined in eqs. (2.172) and (2.173) and where, as we saw in Section 2.1, the $\Xi^{(i)}$ tensors, that we do not display here for brevity, are built from suitable projections of the divergences of eq. (2.176).

The equations for the Lagrange multipliers are in this case

$$\mathcal{Z}^{(1)}_{\nu_1 \nu_2} = 0, \quad \mathcal{Z}^{(2)}_{\mu, \nu} = 0, \quad \mathcal{Z}^{(3)}_{\mu_1 \mu_2} = 0, \quad (2.177)$$

and after enforcing them the equation for an unconstrained gauge field $\psi_{\mu_1 \mu_2 \mu_3, \nu_1 \nu_2}$ becomes

$$\begin{aligned} E_{\bar{\psi}} : \mathcal{E}_{\psi} - \frac{1}{2} \eta_{(\mu_1 \mu_2} \gamma_{\mu_3)} \mathcal{Y}^{(1)}_{\nu_1 \nu_2} - \frac{1}{2} (\eta_{(\mu_1 \mu_2 |} \gamma_{(\nu_1 |} + \eta_{(\nu_1 | (\mu_1} \gamma_{\mu_2 |}) \mathcal{Y}^{(2)}_{| \mu_3), | \nu_2)} \\ - \frac{1}{2} (\eta_{\nu_1 \nu_2} \gamma_{(\mu_1 |} + \gamma_{(\nu_1} \eta_{\nu_2) (\mu_1 |}) \mathcal{Y}^{(3)}_{| \mu_2 \mu_3)} = 0, \end{aligned} \quad (2.178)$$

where the $\mathcal{Y}^{(i)}$ spinor-tensors denote suitable combinations of the fields ψ , $\xi^{(i)}$ and $\lambda^{(i)}$, that we also do not display here for brevity.

As we saw in Section 2.2, the equation for a constrained reducible field $\psi_{\mu_1 \mu_2 \mu_3, \nu_1 \nu_2}$ can also be cast in the form (2.178). However, as in the bosonic case, the field equation that originates from the constrained two-family Lagrangian (2.67) calls for some care if one wants to present it

without introducing a set of $\mathcal{Y}^{(i)}$ spinor-tensors. In fact, differently from the Fang-Fronsdal or one-family case, the symmetrized triple γ -traces of the Rarita-Schwinger tensor (2.176),

$$\begin{aligned}\gamma^\lambda \mathcal{E}'_{\psi\lambda, \nu_1 \nu_2} &= \frac{D+2}{9} v_{(\nu_1, \nu_2)} - \frac{4}{9} u_{\nu_1 \nu_2}, \\ \gamma^\lambda \left(\mathcal{E}'_{\psi\mu, \nu\lambda} + 2 \mathcal{E}'_{\psi\lambda\mu, \nu} \right) &= -\frac{2}{9} v_{(\mu, \nu)} - \frac{D}{9} u_{\mu\nu}, \\ \gamma^\lambda \left(\mathcal{E}'_{\psi\lambda\mu_1\mu_2} + 2 \mathcal{E}'_{\psi\mu_1\mu_2, \lambda} \right) &= -\frac{D+6}{9} v_{(\mu_1, \mu_2)} + \frac{2(D+2)}{9} u_{\mu_1\mu_2},\end{aligned}\tag{2.179}$$

do not vanish identically. At any rate, combining eq. (2.178) with its symmetrized triple γ -traces one can eliminate the $\mathcal{Y}^{(i)}$ and, after gauging away the Ψ_i compensators, one can finally obtain the constrained field equation

$$\begin{aligned}E_{\psi}^{constr} : \mathcal{E}_{\psi} &+ \frac{1}{18(3D+4)} \left\{ \frac{6}{3D-2} \eta_{(\mu_1\mu_2} \gamma_{\mu_3)} [4 u_{\nu_1\nu_2} - (3D+2) v_{(\nu_1, \nu_2)}] \right. \\ &+ \frac{2}{3D-2} (\eta_{(\mu_1\mu_2|} \gamma_{|\nu_1|} + \eta_{(\nu_1|} \mu_1 \gamma_{\mu_2|})} [3 D u_{|\mu_3| \nu_2)} - 2 v_{|\mu_3|, \nu_2)} - 2 v_{|\nu_2|, \mu_3)}] \\ &- \frac{2}{3D-2} (\eta_{\nu_1\nu_2} \gamma_{(\mu_1|} + \gamma_{(\nu_1} \eta_{\nu_2|} \mu_1) } [2 (3D+2) u_{|\mu_2\mu_3)} - 3 (D+2) v_{|\mu_2, \mu_3)}] \\ &\left. + \frac{1}{D^2-2} (2 \eta_{\nu_1\nu_2} \eta_{(\mu_1\mu_2} \omega_{\mu_3)} - \eta_{\nu_1(\mu_1} \eta_{\mu_2| \nu_1} \omega_{|\mu_3)}) \right\} = 0.\end{aligned}\tag{2.180}$$

This expression builds the symmetrized triply γ -traceless projection of $\mathcal{E}_{\psi}$ that, as we shall see in a moment, couples to currents that satisfy the same set of constraints as the gauge field, precisely as was the case for the symmetric spinor-tensors of [19]. As we already pointed out in Section 2.2, this presentation mimics the Fang-Fronsdal setting, and is quite convenient if one wants to work only with the gauge field ψ , but the resulting equations depend explicitly both on the space-time dimension and on the Lorentz labels carried by the gauge fields.

2.4.2 ON-SHELL REDUCTION

As we saw in Section 2.3, the reduction of the equation of motion (2.178) to the non-Lagrangian Labastida form

$$\mathcal{S}_{\mu_1\mu_2\mu_3, \nu_1\nu_2} = 0\tag{2.181}$$

involves new subtleties that are foreign to the symmetric case of [17, 19]. First of all, the $\mathcal{Y}^{(i)}$ tensors do not fully vanish on-shell, but their undetermined components do not contribute to the equations of motion, as a consequence of the shift symmetry (2.68) of the fermionic Lagrangians. The other difficulties can be traced to the anti-commutation relations (A.25), that introduce the S^i_j operators of Appendix A in the γ -traces of eq. (2.82), together with some additional technical complications related to the rather involved form of the fermionic Rarita-Schwinger-like tensors. In order to better illustrate the main steps of the reduction procedure, it is instructive to supplement the general discussion of Section 2.3 with a more direct derivation of the relevant results for the (3,2)-example, in an explicit space-time notation.

One can reduce eq. (2.178) to the Labastida form $\mathcal{S} = 0$ combining it with *all* its admissible

γ -traces. One can begin by taking five γ -traces of the field equation (2.178), which yields

$$\begin{aligned}\gamma \cdot E_{\tilde{\psi}}'^{\mu\hat{r}} &: \mathcal{Y}^{(1)\prime\nu} + \rho_5 \mathcal{Y}^{(2)\hat{r}} - 3\gamma^{\lambda\rho} \mathcal{Y}^{(2)}_{\lambda,\rho} + \rho_2 \mathcal{Y}^{(3)\prime\mu} = 0, \\ \gamma \cdot E_{\tilde{\psi}}'^{\mu\prime\nu} &: \rho_0 \mathcal{Y}^{(1)\prime\nu} + 6\mathcal{Y}^{(2)\hat{r}} + \gamma^{\lambda\rho} \mathcal{Y}^{(2)}_{\lambda,\rho} + \rho_2 \mathcal{Y}^{(3)\prime\mu} = 0, \\ \gamma \cdot E_{\tilde{\psi}}^{\hat{r}\hat{r}} &: \mathcal{Y}^{(1)\prime\nu} + 2\rho_2 \mathcal{Y}^{(2)\hat{r}} + 2\gamma^{\lambda\rho} \mathcal{Y}^{(2)}_{\lambda,\rho} + 6\mathcal{Y}^{(3)\prime\mu} = 0,\end{aligned}\tag{2.182}$$

where the coefficients

$$\rho_k \equiv D + k, \tag{2.183}$$

depend on the space-time dimension D . Notice that we obtained in this fashion *three* relations, those in eq. (2.182), that however involve the *four* available double γ -traces of the $\mathcal{Y}^{(i)}$ tensors. As a result, one can at most link three of these double γ -traces to the remaining one, and here for definiteness we make the choice of relating them to $\mathcal{Y}^{(1)}$, so that

$$\begin{aligned}\mathcal{Y}^{(2)\hat{r}} &= \frac{1}{6} \mathcal{Y}^{(1)\prime\nu}, \\ \gamma^{\lambda\rho} \mathcal{Y}^{(2)}_{\lambda,\rho} &= -\frac{\rho-1}{6} \mathcal{Y}^{(1)\prime\nu}, \\ \mathcal{Y}^{(3)\prime\mu} &= -\frac{2}{3} \mathcal{Y}^{(1)\prime\nu}.\end{aligned}\tag{2.184}$$

Still, decomposing the $\mathcal{Y}^{(i)}$ tensors into their doubly γ -traceless parts and corresponding remainders, according to

$$\begin{aligned}\mathcal{Y}^{(1)}_{\nu_1\nu_2} &= \mathcal{Y}_T^{(1)}_{\nu_1\nu_2} + \frac{1}{\rho_0} \eta_{\nu_1\nu_2} \mathcal{Y}^{(1)\prime\nu}, \\ \mathcal{Y}^{(2)}_{\mu,\nu} &= \mathcal{Y}_T^{(2)}_{\mu,\nu} + \frac{1}{\rho_0} \eta_{\mu\nu} \mathcal{Y}^{(2)\hat{r}} - \frac{1}{\rho_0 \rho - 1} \gamma_{\mu\nu} \gamma^{\lambda\rho} \mathcal{Y}^{(2)}_{\lambda,\rho}, \\ \mathcal{Y}^{(3)}_{\mu_1\mu_2} &= \mathcal{Y}_T^{(3)}_{\mu_1\mu_2} + \frac{1}{\rho_0} \eta_{\mu_1\mu_2} \mathcal{Y}^{(3)\prime\mu},\end{aligned}\tag{2.185}$$

and imposing the relations (2.184), one can verify that the remaining independent quantity disappears from (2.178). As in the bosonic case of [2], this fact simply reflects the shift symmetry of eq. (2.86), that for this example reduces to

$$\begin{aligned}\delta \mathcal{Y}_{111} &= \eta^{22} M_{111,22}, \\ \delta \mathcal{Y}_{112} &= 2\eta^{12} M_{112,12} + 2\gamma^{12} N_{112,12}, \\ \delta \mathcal{Y}_{122} &= \eta^{11} M_{122,11}.\end{aligned}\tag{2.186}$$

With only two index families, the parameter $N_{ijk,lm}$ admits only the $\{4,1\}$ component, and thus the projection does not impose any constraint on it, while $M_{ijk,lm}$ must satisfy the conditions

$$\begin{aligned}Y_{\{5\}} M_{ijk,lm} = 0 &\Rightarrow M_{111,22} + 6M_{112,12} + 3M_{122,11} = 0, \\ Y_{\{3,2\}} M_{ijk,lm} = 0 &\Rightarrow M_{111,22} - 2M_{112,12} + M_{122,11} = 0.\end{aligned}\tag{2.187}$$

However, $N_{ijk,lm}$ is linked to $M_{ijk,lm}$ via eq. (2.69), that in the $(3,2)$ case reduces to

$$N_{112,12} = \frac{1}{12} M_{111,22}, \tag{2.188}$$

so that the shift symmetry rests eventually on the single spinorial parameter $M = M_{111,22}$, and in the space-time notation of this section takes the explicit form

$$\begin{aligned}\delta \mathcal{Y}^{(1)}_{\nu_1 \nu_2} &= \eta_{\nu_1 \nu_2} M, \\ \delta \mathcal{Y}^{(2)}_{\mu, \nu} &= \frac{1}{6} \eta_{\mu \nu} M + \frac{1}{6} \gamma_{\mu \nu} M, \\ \delta \mathcal{Y}^{(3)}_{\mu_1 \mu_2} &= -\frac{2}{3} \eta_{\mu_1 \mu_2} M.\end{aligned}\tag{2.189}$$

Substituting (2.184) in (2.185), the possibility of gauging away the independent component left over by the reduction procedure via (2.189) becomes evident.

In the generic two-family setting we inferred that, after gauging away the undetermined γ -traces of the $\mathcal{Y}_{ijk}$ tensors, the remaining quantities are set to zero by the homogeneous field equations. In the present explicit space-time approach this means, in particular, that after using the shift symmetry (2.86) to reduce the number of independent γ -traces of the $\mathcal{Y}^{(i)}$ tensors, one is properly left with square systems of equations that are not singular provided $D \neq 2, 4$ (see Section 2.3.1). In contrast with the symmetric case of [17, 19], however, it is impossible to disentangle the equations that set to zero the γ -traces of $\mathcal{W}$ from those that do the same for the $\mathcal{Y}^{(i)}$ tensors. This is due to the existence of non-vanishing symmetrized triple γ -traces of $\mathcal{E}_\psi$. However, one can first express the $\mathcal{Y}^{(i)}$ in terms of the γ -traces of $\mathcal{W}$, using the combinations of γ -traces that annihilate ψ in the constrained theory, thus recovering an expression that has the same form as the constrained Lagrangian field equation (2.180). The γ -traces of $\mathcal{W}$ can then be set to zero using the remaining γ -traces of the field equation.

Continuing with the various γ -traces of eq. (2.178), one is to consider $E_{\bar{\psi}}^{\prime \mu \hat{I}}{}_{\nu}$, $(E_{\bar{\psi}}^{\prime \mu \nu} + 2 E_{\bar{\psi}}^{\hat{I} \hat{I}})_{\mu}$, $\gamma^{\lambda \rho} E_{\bar{\psi}}^{\prime \mu}{}_{\lambda, \rho \nu}$ and $\gamma^{\lambda \rho} E_{\bar{\psi}}^{\hat{I}}{}_{\lambda \mu, \rho}$, that can all be obtained combining γ -traces of the symmetrized triple γ -traces of $E_{\bar{\psi}}$ and read, respectively,

$$\begin{bmatrix} \rho_2 & 3D+8 & 2 & 4 \\ 2 & 4 & 3D+10 & 3D+8 \\ \rho_{-1}\rho_2 & -\frac{D^2+3D+4}{2} & 2D+3 & -\rho_0 \\ \rho_0 & -3\rho_2 & (D+2)^2 & -(D^2+2D-2) \end{bmatrix} \cdot \begin{bmatrix} \gamma^\lambda \mathcal{Y}_T^{(1)}{}_{\lambda \mu} \\ \gamma^\lambda \mathcal{Y}_T^{(2)}{}_{\lambda, \mu} \\ \gamma^\lambda \mathcal{Y}_T^{(2)}{}_{\mu, \lambda} \\ \gamma^\lambda \mathcal{Y}_T^{(3)}{}_{\lambda \mu} \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ \frac{\rho-2}{9} \\ -\frac{\rho-2}{9} \end{bmatrix} \omega_\mu, \tag{2.190}$$

together with the additional $\{2, 2\}$ -projected trace

$$(E_{\bar{\psi}}^{\prime \mu \nu} - E_{\bar{\psi}}^{\hat{I} \hat{I}})_{\mu} : \quad \frac{\rho-2}{6} \omega_\mu + \gamma^\lambda \left\{ \mathcal{Y}_T^{(1)}{}_{\lambda \mu} - \mathcal{Y}_T^{(2)}{}_{\lambda, \mu} - \mathcal{Y}_T^{(2)}{}_{\mu, \lambda} + \mathcal{Y}_T^{(3)}{}_{\lambda \mu} \right\} = 0. \tag{2.191}$$

Notice that only the doubly γ -traceless tensors $\mathcal{Y}_T^{(i)}$ appear in these equations, on account of the previous discussion and that, as expected, the number of unknowns now matches the number of equations. Eqs. (2.190) then suffice to obtain

$$\begin{aligned}\gamma^\lambda \mathcal{Y}_T^{(1)}{}_{\lambda \nu} &= \frac{\rho-2}{6(D^2-2)} \omega_\nu, \\ \gamma^\lambda \mathcal{Y}_T^{(2)}{}_{\lambda, \nu} &= -\frac{\rho-2}{18(D^2-2)} \omega_\nu, \\ \gamma^\lambda \mathcal{Y}_T^{(2)}{}_{\mu, \lambda} &= -\frac{\rho-2}{18(D^2-2)} \omega_\mu, \\ \gamma^\lambda \mathcal{Y}_T^{(3)}{}_{\lambda \mu} &= \frac{\rho-2}{18(D^2-2)} \omega_\mu,\end{aligned}\tag{2.192}$$

while eq. (2.191) will eventually set ω to zero. Considering the symmetrized triple γ -traces of the field equations and substituting into them the results collected in eq. (2.192) now gives

$$\begin{aligned} & \begin{bmatrix} \frac{\rho_0 \rho_2}{2} & 2\rho_2 & 4 \\ 2\rho_2 & \frac{3D^2+18D+40}{4} & 4\rho_3 \\ 4 & 4\rho_3 & \frac{\rho_2(3D+8)}{2} \end{bmatrix} \cdot \begin{bmatrix} \mathcal{Y}_T^{(1)}{}_{\mu_1\mu_2} \\ \mathcal{Y}_T^{(2)}{}_{(\mu_1, \mu_2)} \\ \mathcal{Y}_T^{(3)}{}_{\mu_1\mu_2} \end{bmatrix} \\ &= \frac{1}{9} \begin{bmatrix} -4 \\ -\rho_0 \\ 2\rho_2 \end{bmatrix} u_{\mu_1\mu_2} + \frac{1}{9} \begin{bmatrix} \rho_2 \\ 2 \\ -\rho_6 \end{bmatrix} v_{(\mu_1, \mu_2)} - \frac{\rho_0 - 2}{36(D^2 - 2)} \begin{bmatrix} \rho_0 \\ -\rho_0 \\ \rho_0 \end{bmatrix} \gamma_{(\mu_1} \omega_{\mu_2)}, \end{aligned} \quad (2.193)$$

together with the antisymmetric component of the second equation, that simply states that

$$\gamma^\lambda (E_{\bar{\psi}}{}^{\mu}{}_{[\mu, \nu] \lambda} + 2E_{\bar{\psi}}{}^{\dot{\lambda}}{}_{\lambda [\mu, \nu]}) : \mathcal{Y}_T^{(2)}{}_{[\mu, \nu]} = 0. \quad (2.194)$$

All together, these equations finally give

$$\begin{aligned} \mathcal{Y}_T^{(1)}{}_{\nu_1\nu_2} &= -\frac{1}{6(3D+4)} \left\{ \frac{4}{3D-2} [4u_{\nu_1\nu_2} - (3D+2)v_{(\nu_1, \nu_2)}] + \frac{\rho_0}{D^2-2} \gamma_{(\nu_1} \omega_{\nu_2)} \right\}, \\ \mathcal{Y}_T^{(2)}{}_{\mu, \nu} &= -\frac{1}{18(3D+4)} \left\{ \frac{4}{3D-2} [3\rho_0 u_{\mu\nu} - 2v_{(\mu, \nu)}] - \frac{\rho_0}{D^2-2} \gamma_{(\mu} \omega_{\nu)} \right\}, \\ \mathcal{Y}_T^{(3)}{}_{\mu_1\mu_2} &= \frac{1}{18(3D+4)} \left\{ \frac{8}{3D-2} \left[(3D+2)u_{\mu_1\mu_2} - \frac{3}{2}\rho_2 v_{(\mu_1, \mu_2)} \right] - \frac{\rho_0}{D^2-2} \gamma_{(\mu_1} \omega_{\mu_2)} \right\}. \end{aligned} \quad (2.195)$$

Substituting (2.195) in the unconstrained equation of motion (2.178), one can recover (2.180). Let us stress again that after eliminating the compensators Ψ_i this turns into a convenient presentation of the Lagrangian equations for the constrained theory. After these steps, the reduction to the Labastida form would proceed as in the constrained theory, and in fact much as in the bosonic case of [2], barring some inevitable technical complications that we shall not dwell upon here. Indeed, for the sake of brevity we shall confine ourselves here with these cursory remarks, since the actual procedure is straightforward if quite tedious. At any rate, in the next section we shall exhibit the explicit form of the propagator for a $(3, 2)$ field. This expression can be obtained with similar techniques, and actually embodies in a clearcut fashion all relevant information concerning the subtleties that can be met when performing the reduction in special (low numbers of) dimensions where Weyl-like symmetries emerge.

2.4.3 CURRENT EXCHANGES

Coupling a rank- $(3, 2)$ gauge field $\psi_{\mu_1\mu_2\mu_3, \nu_1\nu_2}$ to an external current leads to the inhomogeneous equation

$$\begin{aligned} E_{\bar{\psi}} : \mathcal{E}_\psi &= \frac{1}{2} \eta_{(\mu_1\mu_2} \gamma_{\mu_3)} \mathcal{Y}^{(1)}{}_{\nu_1\nu_2} - \frac{1}{2} (\eta_{(\mu_1\mu_2} \gamma_{\nu_1)} + \eta_{(\nu_1} \gamma_{\mu_2} \gamma_{\mu_3)}) \mathcal{Y}^{(2)}{}_{|\mu_3), |\nu_2)} \\ &- \frac{1}{2} (\eta_{\nu_1\nu_2} \gamma_{(\mu_1} + \gamma_{(\nu_1} \eta_{\nu_2)} \gamma_{\mu_1)}) \mathcal{Y}^{(3)}{}_{|\mu_2\mu_3)} = \mathcal{J}_{\mu_1\mu_2\mu_3, \nu_1\nu_2}, \end{aligned} \quad (2.196)$$

that can be manipulated using the approach followed for the homogeneous equation (2.178). In particular, the same steps of the preceding discussion can reduce eq. (2.196) to the form

$$E_{\bar{\psi}}^{constr} = \mathcal{K}_{\mu_1\mu_2\mu_3, \nu_1\nu_2}, \quad (2.197)$$

where $E_{\bar{\psi}}^{const}$ was defined in eq. (2.180) and $\mathcal{K}_{\mu_1\mu_2\mu_3, \nu_1\nu_2}$ is an effective current whose symmetrized triple γ -traces vanish. We would like to stress again that this form is not unique, since one can also adsorb different parts of the $\mathcal{Y}^{(i)}$ in the definition of the effective current $\mathcal{K}$, but it is anyway quite convenient, since it naturally extends the symmetric constrained setting, were the external current is subject to the same trace constraints as the gauge field.

One can compute further γ -traces of the equation of motion (2.196), and combining these results one can recast it in the form

$$\mathcal{W}_{\mu_1\mu_2\mu_3, \nu_1\nu_2} = \mathcal{J}_{\mu_1\mu_2\mu_3, \nu_1\nu_2} + \dots, \quad (2.198)$$

where the omitted terms are obtained expressing the γ -traces of $\mathcal{W}$ and $\mathcal{Y}^{(i)}$ in terms of $\mathcal{J}$. In order to give rise to the correct current exchanges, the combination on the right-hand side of eq. (2.198) must define a combination that is γ -traceless in $D - 2$ dimensions, and in the following we shall see that this is actually the case for the $(3, 2)$ -example. From the technical viewpoint, one can reach the form (2.198) dealing separately with the various irreducible Young projections of the γ -traces of $E_{\bar{\psi}}$, in order to extract square systems of equations from the results. As we have already pointed out in Section 2.3, this decomposition is the space-time counterpart of the independent treatment of terms corresponding to different values of ℓ , that we used to identify the degenerate cases where the gauge symmetry is enlarged to accommodate Weyl-like shifts.

As we repeatedly stressed, the γ -traces of the equation of motion contain the S^i_j operators. These can displace indices from one set to another, and as a result the propagators of mixed-symmetry fields involve *all* structures obtained displacing indices within the γ -traces of the current $\mathcal{J}$. In conclusion, the field equation (2.196) can be cast in the form

$$\begin{aligned} \mathcal{W}_{\mu_1\mu_2\mu_3, \nu_1\nu_2} = & \mathcal{J}_{\mu_1\mu_2\mu_3, \nu_1\nu_2} + \gamma_{(\mu_1|} K_{11} \cdot \underline{\mathcal{J}}_1^a |_{\mu_2\mu_3), \nu_1\nu_2} + \gamma_{(\nu_1|} K_{12} \cdot \underline{\mathcal{J}}_1^b |_{\mu_1\mu_2\mu_3, | \nu_2)} \\ & + \eta_{(\mu_1\mu_2|} K_{21} \cdot \underline{\mathcal{J}}_2^a |_{\nu_1\nu_2, | \mu_3)} + \eta_{(\nu_1|} (\mu_1 K_{22} \cdot \underline{\mathcal{J}}_2^a |_{\mu_2\mu_3), | \nu_2)} + \eta_{\nu_1\nu_2} K_{23} \cdot \underline{\mathcal{J}}_2^b |_{\mu_1\mu_2\mu_3} \\ & - \gamma_{(\nu_1|} (\mu_1 K_{24} \cdot \underline{\mathcal{J}}_2^a |_{\mu_2\mu_3), | \nu_2)} + \eta_{(\mu_1\mu_2} \gamma_{\mu_3)} K_{31} \cdot \underline{\mathcal{J}}_3 |_{\nu_1\nu_2)} + \eta_{(\mu_1\mu_2|} \gamma_{(\nu_1|} K_{32} \cdot \underline{\mathcal{J}}_3 |_{\mu_3), | \nu_2)} \\ & + \eta_{(\nu_1|} (\mu_1 \gamma_{\mu_2|} K_{33} \cdot \underline{\mathcal{J}}_3 |_{\mu_3), | \nu_2)} + \eta_{(\mu_1|} (\nu_1 \gamma_{\nu_2|} K_{34} \cdot \underline{\mathcal{J}}_3 |_{\mu_2, \mu_3)} + \eta_{\nu_1\nu_2} \gamma_{(\mu_1|} K_{35} \cdot \underline{\mathcal{J}}_3 |_{\mu_2, \mu_3)} \\ & + \eta_{(\mu_1\mu_2} \eta_{\mu_3)} (\nu_1| K_{41} \cdot \underline{\mathcal{J}}_4 |_{\nu_2)} + \eta_{\nu_1\nu_2} \eta_{(\mu_1\mu_2|} K_{42} \cdot \underline{\mathcal{J}}_4 |_{\mu_3)} + \eta_{(\mu_1|} (\nu_1 \eta_{\nu_2|} |_{\mu_2|} K_{43} \cdot \underline{\mathcal{J}}_4 |_{\mu_3)} \\ & + \eta_{(\mu_1\mu_2} \gamma_{\mu_3)} (\nu_1| K_{44} \cdot \underline{\mathcal{J}}_4 |_{\nu_2)} - \eta_{(\mu_1|} (\nu_1 \gamma_{\nu_2|} |_{\mu_2|} K_{45} \cdot \underline{\mathcal{J}}_4 |_{\mu_3)} + \eta_{(\mu_1\mu_2} \eta_{\mu_3)} (\nu_1 \gamma_{\nu_2|} K_{51} \cdot \underline{\mathcal{J}}_5 \\ & + \eta_{\nu_1\nu_2} \eta_{(\mu_1\mu_2} \gamma_{\mu_3)} K_{52} \cdot \underline{\mathcal{J}}_5 + \eta_{(\mu_1|} (\nu_1 \eta_{\nu_2|} |_{\mu_2|} \gamma_{\mu_3)} K_{53} \cdot \underline{\mathcal{J}}_5. \end{aligned} \quad (2.199)$$

The resulting propagator is rather complicated, and contains as many as 106 distinct contributions. Two types of vectors, denoted by $\underline{\mathcal{J}}_i$ and K_{ij} , actually build this expression, and their scalar products are denoted with a standard symbol, “ $\cdot$ ”. These vectors, however, have three, four, five or seven distinct components, depending on the case. The vectors of the first type collect essentially all possible constructs built from the current $\mathcal{J}$,

$$\begin{aligned} \underline{\mathcal{J}}_1^a |_{\mu_1\mu_2, \nu_1\nu_2} &= \gamma^\lambda (\mathcal{J}_{\lambda\mu_1\mu_2, \nu_1\nu_2}, \mathcal{J}_{\lambda(\mu_1|} (\nu_1, \nu_2) |_{\mu_2)}, \mathcal{J}_{\lambda\nu_1\nu_2, \mu_1\mu_2}, \mathcal{J}_{\mu_1\mu_2} (\nu_1, \nu_2)^\lambda, \mathcal{J}_{\nu_1\nu_2} (\mu_1, \mu_2)^\lambda), \\ \underline{\mathcal{J}}_1^b |_{\mu_1\mu_2\mu_3, \nu} &= \gamma^\lambda (\mathcal{J}_{\lambda(\mu_1\mu_2, \mu_3) \nu}, \mathcal{J}_{\lambda\nu} (\mu_1, \mu_2\mu_3), \mathcal{J}_{\mu_1\mu_2\mu_3, \nu\lambda}, \mathcal{J}_{\nu} (\mu_1\mu_2, \mu_3)^\lambda), \\ \underline{\mathcal{J}}_2^a |_{\mu_1\mu_2, \nu} &= (\mathcal{J}'^\mu_{\nu, \mu_1\mu_2}, \mathcal{J}'^\mu_{(\mu_1, \mu_2) \nu}, \mathcal{J}'^\mu_{\mu_1\mu_2, \nu}, \mathcal{J}'^\mu_{\nu} (\mu_1, \mu_2), \mathcal{J}'^\nu_{\mu_1\mu_2\nu}, \gamma^{\lambda\rho} \mathcal{J}_{\lambda\mu_1\mu_2, \nu\rho}, \\ &\quad \gamma^{\lambda\rho} \mathcal{J}_{\lambda\nu} (\mu_1, \mu_2)^\rho), \\ \underline{\mathcal{J}}_2^b |_{\mu_1\mu_2\mu_3} &= (\mathcal{J}'^\mu_{(\mu_1, \mu_2\mu_3)}, \mathcal{J}'^\mu_{(\mu_1\mu_2, \mu_3)}, \mathcal{J}'^\nu_{\mu_1\mu_2\mu_3}, \gamma^{\lambda\rho} \mathcal{J}_{\lambda(\mu_1\mu_2, \mu_3) \rho}), \end{aligned}$$

$$\begin{aligned}
\underline{\mathcal{J}}_{3\mu,\nu} &= \gamma^\lambda \left(\mathcal{J}'_{\mu\lambda,\mu\nu}, \frac{1}{2} \mathcal{J}'_{\mu(\mu,\nu)\lambda}, \frac{1}{2} \mathcal{J}'_{\mu[\mu,\nu]\lambda}, \frac{1}{2} \mathcal{J}'_{\lambda(\mu,\nu)}, \frac{1}{2} \mathcal{J}'_{\lambda[\mu,\nu]}, \mathcal{J}'_{\lambda\mu\nu}, \mathcal{J}'_{\mu\nu,\lambda} \right), \\
\underline{\mathcal{J}}_{4\mu} &= \left(\mathcal{J}'_{\mu\hat{\mu}}, \mathcal{J}'_{\mu'\nu\mu}, \mathcal{J}'_{\mu\hat{\mu}}, \gamma^{\lambda\rho} \mathcal{J}'_{\mu\lambda}, \gamma^{\lambda\rho} \mathcal{J}'_{\mu\lambda} \right), \\
\underline{\mathcal{J}}_5 &= \gamma \cdot \left(\mathcal{J}'_{\mu\hat{\mu}}, \mathcal{J}'_{\mu'\nu}, \mathcal{J}'_{\hat{\mu}\hat{\mu}} \right),
\end{aligned} \tag{2.200}$$

while their partners K_{ij} are vectors of coefficients that are defined below, and whose list closes the present section.

Let us begin by displaying the terms that in eq. (2.199) are accompanied by a single γ -matrix:

$$\begin{aligned}
K_{11} &= \frac{(-D^4 - 4D^3 + 20D^2 + 48D - 64, -4\rho_{-2}\rho_4, -64, 2\rho_2(D^2 + 2D - 16), 16\rho_2)}{\rho_{-4}\rho_{-2}\rho_2\rho_4\rho_6}, \\
K_{12} &= \frac{(2(D^2 + 2D - 16), 16, -\rho_2(D^2 + 2D - 20), -4\rho_2)}{\rho_{-4}\rho_{-2}\rho_4\rho_6}.
\end{aligned} \tag{2.201}$$

Notice that K_{11} and K_{12} have poles in $D = 2, 4$, precisely where the general considerations of Section 2.3 apply for a reducible $(3, 2)$ field. The same poles are also present in most of the other coefficients, that we collect in a few groups. The first group thus comprises the terms that in eq. (2.199) are accompanied by two γ -matrices

$$\begin{aligned}
K_{21} &= \frac{1}{\rho_{-4}\rho_{-2}\rho_0\rho_2\rho_4\rho_6} (-\rho_{-2}(D^4 + 6D^3 - 12D^2 - 96D - 96), -2D(D^2 - 2D - 16), \\
&\quad 16(D^2 - 3D - 6), 2(D^4 + D^3 - 22D^2 - 8D + 48), -4\rho_2(D^2 - 12), -16\rho_6, -2\rho_6(D^2 - 8)), \\
K_{22} &= \frac{1}{\rho_{-4}\rho_{-2}\rho_0\rho_2\rho_4\rho_6} (8(D^2 - 3D - 6), D^4 + D^3 - 22D^2 - 8D + 48, -D^5 - 3D^4 + 23D^3 \\
&\quad + 42D^2 - 68D - 72, -4(D^3 - D^2 - 11D - 6), \rho_2(D^3 + D^2 - 16D - 12), \rho_6(D^2 - 12), 4\rho_6), \\
K_{23} &= \frac{(-4(D^2 - 12), 2(D^3 + D^2 - 16D - 12), -D^4 - 4D^3 + 22D^2 + 64D - 72, 2\rho_{-2}\rho_6)}{\rho_{-4}\rho_{-2}\rho_0\rho_4\rho_6}, \\
K_{24} &= \frac{(8, D^2 - 8, 12 - D^2, -4, -\rho_{-2}\rho_2, -\rho_1(D^2 - 12), -4\rho_1)}{\rho_{-4}\rho_{-2}\rho_0\rho_2\rho_4},
\end{aligned} \tag{2.202}$$

and in a similar fashion the second group comprises the terms that are accompanied by three γ -matrices

$$\begin{aligned}
K_{31} &= \frac{1}{\rho_{-4}\rho_{-2}\rho_0\rho_2\rho_4\rho_6} (D(D^3 + 4D^2 - 16D - 56), -4\rho_2(D^2 - 12), 0, \\
&\quad -4D(D^2 - 2D - 16), 0, 4\rho_{-2}\rho_0, 8\rho_{-6}\rho_2), \\
K_{32} &= \frac{1}{\rho_{-4}\rho_{-2}\rho_0\rho_2\rho_4\rho_6} (-2\rho_2(D^2 - 12), \rho_2^2(D^2 - 12), \rho_{-4}\rho_0\rho_2\rho_6, -4\rho_{-2}\rho_0\rho_2, \\
&\quad -4\rho_{-4}\rho_0\rho_6, 4\rho_{-6}\rho_2, -2\rho_{-6}\rho_2^2), \\
K_{33} &= \frac{1}{\rho_{-4}\rho_{-2}\rho_2\rho_4\rho_6} (-D^2 + 2D + 16, -2\rho_{-2}\rho_2, -2\rho_{-4}\rho_6, D^3 + 2D^2 - 8D - 32, \\
&\quad \rho_{-4}\rho_0\rho_6, -D^2 + 2D + 16, -2\rho_{-2}\rho_2), \\
K_{34} &= \frac{(4\rho_{-6}, -2\rho_{-6}\rho_2, 0, -4\rho_{-2}\rho_0, 0, -2(D^2 - 12), \rho_2(D^2 - 12))}{\rho_{-4}\rho_{-2}\rho_0\rho_4\rho_6}, \\
K_{35} &= \frac{1}{\rho_{-4}\rho_{-2}\rho_0\rho_2\rho_4\rho_6} (4\rho_{-2}\rho_0, 8\rho_{-6}\rho_2, 0, -4D(D^2 - 2D - 16), 0, \\
&\quad D(D^3 + 4D^2 - 16D - 56), -4\rho_2(D^2 - 12)).
\end{aligned} \tag{2.203}$$

Finally, the last two groups comprise the terms that are accompanied by four γ -matrices,

$$\begin{aligned}
K_{41} &= \frac{(D^3 + D^2 - 17D - 18, -D^2 + 3D + 6, -2(D^2 - 3D - 6), \rho_6, 2\rho_6)}{\rho_{-4} \rho_0 \rho_2 \rho_4 \rho_6}, \\
K_{42} &= \frac{1}{\rho_{-4} \rho_{-2} \rho_{-1} \rho_0 \rho_2 \rho_4 \rho_6} (-2\rho_{-2} \rho_{-1} (D^2 - 3D - 6), D^5 + 3D^4 - 16D^3 - 62D^2 \\
&\quad + 24D + 120, -2(3D^4 - 2D^3 - 46D^2 + 8D + 72), -2\rho_6 (D^2 - D - 10), 4\rho_{-6} \rho_6), \\
K_{43} &= \frac{1}{\rho_{-4} \rho_{-2} \rho_{-1} \rho_0 \rho_2 \rho_4 \rho_6} (-2\rho_{-2} \rho_{-1} (D^2 - 3D - 6), -3D^4 + 2D^3 + 46D^2 - 8D - 72, \\
&\quad D^5 - 14D^3 - 16D^2 + 16D + 48, 2\rho_{-6} \rho_6, -2\rho_6 (D^2 - 2D - 4)), \\
K_{44} &= \frac{(-\rho_{-2} \rho_{-1}, D^2 - D - 10, -2\rho_{-6}, \rho_1 (D^2 - D - 10), -2\rho_{-6} \rho_1)}{\rho_{-4} \rho_{-2} \rho_{-1} \rho_0 \rho_2 \rho_4}, \\
K_{45} &= \frac{(-\rho_{-2} \rho_{-1}, -\rho_{-6}, D^2 - 2D - 4, -\rho_{-6} \rho_1, \rho_1 (D^2 - 2D - 4))}{\rho_{-4} \rho_{-2} \rho_{-1} \rho_0 \rho_2 \rho_4}, \tag{2.204}
\end{aligned}$$

or by five γ -matrices,

$$\begin{aligned}
K_{51} &= \frac{(-\rho_2, 2, 4)}{\rho_{-4} \rho_2 \rho_4 \rho_6}, \\
K_{52} &= \frac{(4\rho_{-2} \rho_{-1}, -D^3 - 3D^2 + 10D + 64, 2(3D^2 - 6D - 32))}{\rho_{-4} \rho_{-2} \rho_{-1} \rho_2 \rho_4 \rho_6}, \\
K_{53} &= \frac{(4\rho_{-2} \rho_{-1}, 3D^2 - 6D - 32, -D^3 + 4D + 32)}{\rho_{-4} \rho_{-2} \rho_{-1} \rho_2 \rho_4 \rho_6}. \tag{2.205}
\end{aligned}$$

The reader could verify, as we also did, that the right-hand side of eq. (2.199) defines an expression whose γ -traces in each of its two index sets vanish once the D -dependent factors that arise in this computation (but, of course, not those appearing in the original K_{ij} vector coefficients) are shifted letting $D \rightarrow D - 2$.

3 GENERAL FERMIONIC FIELDS

In this section we can finally derive Lagrangians and field equations for general unconstrained fermionic fields with arbitrary numbers of index families. We open our discussion by showing in detail how to complete the work of Labastida [30] with general constrained Lagrangians yielding the fermionic field equations that he introduced in the eighties. To this end, we follow rather closely the approach developed in [2] for Bose fields and based on the Bianchi identities, that as we shall see is particularly convenient for the extension to the unconstrained case. However, we also display an alternative derivation, based on the request that the Rarita-Schwinger-like kinetic operators be self-adjoint, that proceeds along the lines of the original bosonic work of Labastida [12], before closing the section with a discussion of the unconstrained Lagrangians, their field equations and their reduction to the form of [30].

3.1 THE CONSTRAINED LAGRANGIANS

The γ -traces of the Bianchi identities are a key tool to derive the Lagrangians for mixed-symmetry fermions with arbitrary numbers of index families. In the constrained setting they

take the form

$$\partial_i \mathcal{S} - \frac{1}{2} \not{\partial} \mathcal{S}_i - \frac{1}{2} \partial^j T_{ij} \mathcal{S} - \frac{1}{6} \partial^j \gamma_{ij} \mathcal{S} = 0, \quad (3.1)$$

where we have enforced the Labastida constraints

$$T_{(ij} \not{\psi}_{k)} = 0 \quad (3.2)$$

on the right-hand side of eq. (3.1). As we already saw in Section 2.1, the γ -trace of eq. (3.1),

$$\partial_{[i} \mathcal{S}_{j]} - \frac{2}{3} \not{\partial} \gamma_{ij} \mathcal{S} + \frac{1}{6} \partial^k \gamma_{ijk} \mathcal{S} - \frac{1}{6} \partial^k (3 T_{ik} \gamma_j + T_{j[i} \gamma_{k]}) \mathcal{S} = 0, \quad (3.3)$$

admits two independent projections. In analogy with the bosonic case, the symmetric one yields the gradient of

$$T_{(ij} \mathcal{S}_{k)} = 0, \quad (3.4)$$

an identity implied by the constraint (3.2), as one can verify by a direct computation. The antisymmetric projection, however, is more interesting, and reads

$$\partial_{[i} \mathcal{S}_{j]} - \frac{2}{3} \not{\partial} \gamma_{ij} \mathcal{S} + \frac{1}{6} \partial^k \gamma_{ijk} \mathcal{S} - \frac{1}{3} \partial^k T_{k[i} \gamma_{j]} \mathcal{S} = 0, \quad (3.5)$$

so that it relates in a non-trivial fashion different structures.

This suggests that the Lagrangians should rest on a special sub-chain of suitably projected consequences of the Bianchi identities, in analogy with the behavior of Bose fields described in [2]. In order to sort it out, it is convenient to first identify the non trivial (γ -)traces of the kinetic tensor $\mathcal{S}$, in order to build the most general Ansatz. In Appendix C we shall prove that, for any given number of traces and antisymmetric γ -traces of the kinetic tensor $\mathcal{S}$, *all Young projections in family indices with more than two columns vanish in the constrained theory* on account of eq. (3.4). This condition identifies the wide class of terms that can enter fermionic Lagrangians which, as a result, are far more involved than their bosonic counterparts. For N families the Lagrangians are in fact bound to rest on $\mathcal{O}(N^2)$ distinct Young projections, with a first column of length $l_1 \leq N$ and a second column of length $l_2 \leq l_1$, that in general is actually shorter, since antisymmetric γ -traces $\gamma_{i_1 \dots i_n}$ of $\mathcal{S}$ can accompany the ordinary T_{ij} traces that already enter the bosonic Lagrangians. Interestingly, *any* given combination of these operators admits *a single two-column projection*, and it is then convenient to adopt the short-hand notation

$$\begin{aligned} \mathcal{S}'_{ij} &= T_{ij} \mathcal{S}, \\ \mathcal{S}'_{ij;k} &= \frac{1}{3} (2 T_{ij} \gamma_k - T_{k(i} \gamma_{j)}) \mathcal{S}, \\ &\vdots \\ (\gamma^{[q]} \mathcal{S}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q} &= Y_{\{2p, 1^q\}} T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q} \mathcal{S}. \end{aligned} \quad (3.6)$$

The reader should appreciate that here colons separate again groups of symmetric indices, while a semicolon precedes the left-over group of antisymmetric ones.

In conclusion, the Lagrangian for generic N -family fermionic gauge fields ψ is of the form

$$\mathcal{L} = \frac{1}{2} \langle \bar{\psi}, \sum_{p,q=0}^N k_{p,q} \eta^p \gamma^q (\gamma^{[q]} \mathcal{S}^{[p]}) \rangle + \text{h.c.}, \quad (3.7)$$

where

$$\eta^p \gamma^q (\gamma^{[q]} \mathcal{S}^{[p]}) \equiv \eta^{i_1 j_1} \dots \eta^{i_p j_p} \gamma^{k_1 \dots k_q} (\gamma^{[q]} \mathcal{S}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q}. \quad (3.8)$$

As we shall see shortly, the $k_{p,q}$ coefficients in eq. (3.7) are uniquely determined by the condition that $\mathcal{L}$ be gauge invariant, up to the convenient choice $k_{0,0} = 1$.

In order to proceed, we can begin by noticing that the identities collected in Appendix B make it possible to recast the gauge variation of eq. (3.7), up to total derivatives, in the form

$$\begin{aligned} \delta \mathcal{L} = & - \sum_{p,q=0}^N \frac{1}{2^{p+1}} \langle T_{i_1 j_1} \dots T_{i_p j_p} \bar{\epsilon}_l \gamma_{k_1 \dots k_q}, k_{p,q} \partial_l (\gamma^{[q]} \mathcal{S}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q} \\ & + (q+1) k_{p,q+1} \not{\partial} (\gamma^{[q+1]} \mathcal{S}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q l} \\ & + (p+1) k_{p+1,q} \partial^m (\gamma^{[q]} \mathcal{S}^{[p+1]})_{i_1 j_1, \dots, i_p j_p, lm; k_1 \dots k_q} \\ & + (q+1)(q+2) k_{p,q+2} \partial^m (\gamma^{[q+2]} \mathcal{S}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q lm} \rangle + \text{h.c.} \end{aligned} \quad (3.9)$$

Notice that all $\gamma^{[q]} \mathcal{S}^{[p]}$ terms that are right entries of the scalar product are two-column projected by assumption. This statement applies to all free indices carried by the last three terms, while the first is slightly more complicated, since due to the l index carried by the divergence it also allows projections with three boxes in the first row. However, the left entry of the scalar product makes this type of projections proportional to the constraints on the gauge parameters

$$\gamma_{(i} \epsilon_{j)} = 0, \quad (3.10)$$

as expected, on account of the identity

$$\begin{aligned} & \langle Y_{\{3,2^{p-1},1^q\}} T_{i_1 j_1} \dots T_{i_p j_p} \bar{\epsilon}_l \gamma_{k_1 \dots k_q}, \partial_l (\gamma^{[q]} \mathcal{S}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q} \rangle \\ & = \langle (p+1) T_{i_2 j_2} \dots T_{i_p j_p} T_{(i_1 j_1} \bar{\epsilon}_{l)} \gamma_{k_1 \dots k_q} - T_{i_2 j_2} \dots T_{i_p j_p} T_{[k_1 | (l} \bar{\epsilon}_{i_1} \gamma_{j_1)} \gamma_{k_2 \dots k_q]}, \quad (3.11) \\ & \quad \frac{p}{3(p+1)(p+q+2)} \partial_{(l} (\gamma^{[q]} \mathcal{S}^{[p]})_{|i_1 j_1), \dots, i_p j_p; k_1 \dots k_q} \rangle, \end{aligned}$$

that we shall prove in Appendix C. In conclusion, up to the Labastida constraints one can effectively restrict the attention to two-column projections of the gauge variation (3.9).

As we saw in the two-family case, fermionic fields bring about a novel type of complication. Indeed, the γ -traces of the Bianchi identities do not suffice to set to zero the individual terms of eq. (3.9), but these, on the other hand, are not fully independent and can be partly combined using the constraints (3.10) on the gauge parameters. Even in the general case, the possible Young projections for the family indices are the guiding principle when trying to combine different terms in eq. (3.9). And indeed all terms of the type

$$T_{i_1 j_1} \dots T_{i_p j_p} \bar{\epsilon}_l \gamma_{k_1 \dots k_q} \quad (3.12)$$

that are left entries of the scalar products admit two distinct Young projections with two columns, since

$$\{2^p, 1^q\} \otimes \{1\} = \{2^p, 1^{q+1}\} \oplus \{2^{p+1}, 1^{q-1}\} \oplus \{3, 2^{p-1}, 1^q\}. \quad (3.13)$$

Leaving aside the last term, that is clearly related to the constraints, the first two projections can thus arise from *pairs* of neighboring contributions of the form (3.12), with values of p differing by one unit and with values of q differing by two units.

As in the two-family case, the key step is to relate these two types of terms, that are only apparently different, making use of the constraints (3.10) and of proper relabelings. To this end, it proves particularly convenient to manipulate the combinations $\bar{\epsilon}_l \gamma_{k_1 \dots k_q}$ according to

$$\bar{\epsilon}_l \gamma_{k_1 \dots k_q} = \frac{1}{q+1} \bar{\epsilon}_{[l} \gamma_{k_1 \dots k_q]} + \frac{1}{q+1} \sum_{n=1}^q (-1)^{(n+1)} \bar{\epsilon}_{(l} \gamma_{k_n) k_1 \dots k_{r \neq n} \dots k_q} . \quad (3.14)$$

The terms entering the sum can then be rewritten more conveniently as

$$\bar{\epsilon}_{(l} \gamma_{k_1) k_2 \dots k_q} = \bar{\epsilon}_{(l} \gamma_{k_1)} \gamma_{k_2 \dots k_q} - T_{[k_2 | (l} \bar{\epsilon}_{k_1)} \gamma_{| k_3 \dots k_q]} , \quad (3.15)$$

where the first contribution on the right-hand side clearly vanishes in the constrained setting, on account of eq. (3.10). Enforcing the constraints thus leads to

$$\sum_{n=1}^q (-1)^{(n+1)} \bar{\epsilon}_{(l} \gamma_{k_n) k_1 \dots k_{r \neq n} \dots k_q} \rightarrow T_{l[k_1} \bar{\epsilon}_{k_2} \gamma_{k_3 \dots k_q]} , \quad (3.16)$$

since the sum with oscillating signs induces a complete antisymmetry in the k_r indices, that as such forbids the simultaneous presence of two of them on the trace T . As a result, in the constrained theory eq. (3.14) can be finally cast in the form

$$T_{i_1 j_1} \dots T_{i_p j_p} \bar{\epsilon}_l \gamma_{k_1 \dots k_q} = \frac{1}{q+1} T_{i_1 j_1} \dots T_{i_p j_p} (\bar{\epsilon}_{[l} \gamma_{k_1 \dots k_q]} + T_{l[k_1} \bar{\epsilon}_{k_2} \gamma_{k_3 \dots k_q]}) , \quad (3.17)$$

where the first term is annihilated by the $\{2^{p+1}, 1^{q-1}\}$ projection while the second is similarly annihilated by the $\{2^p, 1^{q+1}\}$ projection, and in conclusion one is led to

$$Y_{\{2^p, 1^{q+1}\}} T_{i_1 j_1} \dots T_{i_p j_p} \bar{\epsilon}_l \gamma_{k_1 \dots k_q} = \frac{1}{q+1} Y_{\{2^p, 1^{q+1}\}} T_{i_1 j_1} \dots T_{i_p j_p} \bar{\epsilon}_{[l} \gamma_{k_1 \dots k_q]} \quad (3.18)$$

and

$$Y_{\{2^{p+1}, 1^{q-1}\}} T_{i_1 j_1} \dots T_{i_p j_p} \bar{\epsilon}_l \gamma_{k_1 \dots k_q} = \frac{1}{q+1} Y_{\{2^{p+1}, 1^{q-1}\}} T_{i_1 j_1} \dots T_{i_p j_p} T_{l[k_1} \bar{\epsilon}_{k_2} \gamma_{k_3 \dots k_q]} . \quad (3.19)$$

While the first of these relations is rather trivial, the second is quite interesting, since it connects a relatively complicated projection of p traces and a single q -fold antisymmetric γ -trace to a simpler one involving $p+1$ traces and a $(q-2)$ -fold antisymmetric γ -trace with the maximum possible number of antisymmetrizations, generalizing in this way the relation (2.40) that we used to build two-family Lagrangians. In this fashion, the current choice extends the $\kappa = 1$ construction of Section 2.1. Let us stress that one can also proceed in the opposite direction, decreasing the value of p . The relevant identity, that can be proved using the techniques of Appendix C and generalizes eq. (2.39) to the N -family setting, is then

$$Y_{\{2^p, 1^{q+1}\}} T_{i_1 j_1} \dots T_{i_p j_p} \bar{\epsilon}_l \gamma_{k_1 \dots k_q} = \frac{(-1)^{q+1}}{p(q+1)} Y_{\{2^p, 1^{q+1}\}} \sum_{n=1}^p \left(\prod_{r \neq n}^p T_{i_r j_r} \right) \bar{\epsilon}_{(i_n} \gamma_{j_n) k_1 \dots k_q l} , \quad (3.20)$$

but it is clearly simpler to work with terms whose projections involve the maximum number of antisymmetrizations compatible with their tensorial character.

As we anticipated, making use of eq. (3.11) one can eliminate all three column projections from the gauge variation (3.9), while the remaining two-column projected terms read

$$\begin{aligned}
\delta \mathcal{L} = & - \sum_{p,q=0}^N \frac{1}{2^{p+1}} \langle Y_{\{2^p, 1^{q+1}\}} T_{i_1 j_1} \dots T_{i_p j_p} \bar{\epsilon}_l \gamma_{k_1 \dots k_q}, k_{p,q} \partial_l (\gamma^{[q]} \mathcal{S}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q} \\
& + (q+1) k_{p,q+1} \not{\partial} (\gamma^{[q+1]} \mathcal{S}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q l} \\
& + (p+1) k_{p+1,q} \partial^m (\gamma^{[q]} \mathcal{S}^{[p+1]})_{i_1 j_1, \dots, i_p j_p, lm; k_1 \dots k_q} \\
& + (q+1)(q+2) k_{p,q+2} \partial^m (\gamma^{[q+2]} \mathcal{S}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q lm} \rangle \\
& - \sum_{p,q=0}^N \frac{1}{2^{p+1}} \langle Y_{\{2^{p+1}, 1^{q-1}\}} T_{i_1 j_1} \dots T_{i_p j_p} \bar{\epsilon}_l \gamma_{k_1 \dots k_q}, k_{p,q} \partial_l (\gamma^{[q]} \mathcal{S}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q} \\
& + (p+1) k_{p+1,q} \partial^m (\gamma^{[q]} \mathcal{S}^{[p+1]})_{i_1 j_1, \dots, i_p j_p, lm; k_1 \dots k_q} \rangle + \text{h.c.} .
\end{aligned} \tag{3.21}$$

Notice that the first term in eq. (3.9) gave contributions to both groups of projected terms above, while the second term gave a single contribution. The analysis of the gradient terms was slightly more subtle, but the end result was that one of them, the third term in eq. (3.9), contributed to both. Indeed,

$$\begin{aligned}
\{2^p, 1^{q+1}\} \otimes \{1\} &= \{2^p, 1^{q+2}\} \oplus \{2^{p+1}, 1^q\} \oplus \{3, 2^{p-1}, 1^{q+1}\}, \\
\{2^{p+1}, 1^{q-1}\} \otimes \{1\} &= \{2^{p+1}, 1^q\} \oplus \{2^{p+2}, 1^{q-2}\} \oplus \{3, 2^p, 1^{q-1}\},
\end{aligned} \tag{3.22}$$

so that both types of projections are compatible with the $\{2^{p+1}, 1^q\}$ projection carried by $(\gamma^{[q]} \mathcal{S}^{[p+1]})$. One can now manipulate the second group of terms in eq. (3.21) using the key relation (3.19), and after a proper relabeling eq. (3.21) can be finally turned into

$$\begin{aligned}
\delta \mathcal{L} = & - \sum_{p,q=0}^N \frac{1}{2^{p+1}} \langle Y_{\{2^p, 1^{q+1}\}} T_{i_1 j_1} \dots T_{i_p j_p} \bar{\epsilon}_l \gamma_{k_1 \dots k_q}, \\
& k_{p,q} \partial_l (\gamma^{[q]} \mathcal{S}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q} \\
& + \frac{(q+1)(q+2)}{p(q+3)} k_{p-1,q+2} \sum_{n=1}^p \partial_{(i_n |} (\gamma^{[q+2]} \mathcal{S}^{[p-1]})_{\dots, i_r \neq n j_r \neq n, \dots; | j_n) l k_1 \dots k_q} \\
& + (q+1) k_{p,q+1} \not{\partial} (\gamma^{[q+1]} \mathcal{S}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q l} \\
& + (p+1) k_{p+1,q} \partial^m (\gamma^{[q]} \mathcal{S}^{[p+1]})_{i_1 j_1, \dots, i_p j_p, lm; k_1 \dots k_q} \\
& + \frac{(q+1)(q+2)}{q+3} k_{p,q+2} \sum_{n=1}^p \partial^m (\gamma^{[q+2]} \mathcal{S}^{[p]})_{\dots, i_r \neq n j_r \neq n, \dots, m(i_n; j_n) l k_1 \dots k_q} \\
& + (q+1)(q+2) k_{p,q+2} \partial^m (\gamma^{[q+2]} \mathcal{S}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q lm} \rangle + \text{h.c.},
\end{aligned} \tag{3.23}$$

where we enforced the symmetry under interchanges of the (i_n, j_n) pairs brought about by the product of identical T tensors present on the right side of the scalar product. Furthermore, the contributions in the last two terms are proportional, since

$$(\gamma^{[q+2]} \mathcal{S}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q lm} = (\gamma^{[q+2]} \mathcal{S}^{[p]})_{i_1 j_1, \dots, i_{p-1} j_{p-1}, m(i_p; j_p) l k_1 \dots k_q}, \tag{3.24}$$

because a symmetrization in (i_p, j_p, m) vanishes on account of the two-column projection. Therefore, eq. (3.9) finally becomes

$$\begin{aligned}
\delta \mathcal{L} = & - \sum_{p,q=0}^N \frac{1}{2^{p+1}} \langle Y_{\{2^p, 1^{q+1}\}} T_{i_1 j_1} \dots T_{i_p j_p} \bar{e}_l \gamma_{k_1 \dots k_q} , \\
& k_{p,q} \partial_l (\gamma^{[q]} \mathcal{S}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q} \\
& + \frac{(q+1)(q+2)}{p(q+3)} k_{p-1, q+2} \sum_{n=1}^p \partial_{(i_n |} (\gamma^{[q+2]} \mathcal{S}^{[p-1]})_{\dots, i_r \neq n j_r \neq n, \dots; | j_n) l k_1 \dots k_q} \\
& + (q+1) k_{p, q+1} \not{\partial} (\gamma^{[q+1]} \mathcal{S}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q l} \\
& + (p+1) k_{p+1, q} \partial^m (\gamma^{[q]} \mathcal{S}^{[p+1]})_{i_1 j_1, \dots, i_p j_p, lm; k_1 \dots k_q} \\
& + \frac{(q+1)(q+2)(q+p+3)}{q+3} k_{p, q+2} \partial^m (\gamma^{[q+2]} \mathcal{S}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q lm} \rangle + \text{h.c.} .
\end{aligned} \tag{3.25}$$

In the fermionic case, the γ -traces of the Bianchi identities take the rather involved form presented in Appendix C, but it is possible to extract from them the far simpler chain of $\{2^p, 1^{q+1}\}$ -projected identities

$$\begin{aligned}
& (p+q+2) Y_{\{2^p, 1^{q+1}\}} \partial_l (\gamma^{[q]} \mathcal{S}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q} \\
& + \frac{1}{q+3} Y_{\{2^p, 1^{q+1}\}} \sum_{n=1}^p \partial_{(i_n |} (\gamma^{[q+2]} \mathcal{S}^{[p-1]})_{\dots, i_r \neq n j_r \neq n, \dots; | j_n) l k_1 \dots k_q} \\
& + (-1)^{q+1} \not{\partial} (\gamma^{[q+1]} \mathcal{S}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q l} - \partial^m (\gamma^{[q]} \mathcal{S}^{[p+1]})_{i_1 j_1, \dots, i_p j_p, lm; k_1 \dots k_q} \\
& - \frac{1}{q+3} \partial^m (\gamma^{[q+2]} \mathcal{S}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q lm} = 0 .
\end{aligned} \tag{3.26}$$

Comparing eqs. (3.25) and (3.26) one can now recognize that, in order to obtain a gauge invariant Lagrangian, the coefficients should satisfy the two recursion relations

$$\begin{aligned}
k_{p+1, q} &= - \frac{1}{(p+1)(p+q+2)} k_{p, q} , \\
k_{p, q+1} &= \frac{(-1)^{q+1}}{(q+1)(p+q+2)} k_{p, q} ,
\end{aligned} \tag{3.27}$$

whose solution is

$$k_{p, q} = \frac{(-1)^{p + \frac{q(q+1)}{2}}}{p! q! (p+q+1)!} \tag{3.28}$$

if $k_{0,0} = 1$, a result that is also fully consistent with the other terms in eq. (3.26). Notice that this expression includes as special cases the coefficients first obtained by Labastida for Bose fields: in fact, letting $q = 0$ in eq. (3.28) recovers the result presented in [2], that differs from the original one of [12] only due to a slight change of conventions and to the elimination of a typo in the relative signs.

The coefficients given in eq. (3.28) determine completely the constrained Lagrangians (3.7) for mixed-symmetry Fermi fields. Together with the corresponding unconstrained Lagrangians that we shall soon come across in eq. (3.42), these are the main results of this work.

Interestingly, eq. (3.28) can be also obtained demanding that the kinetic operator be self-adjoint, along the lines of what was done by Labastida for Bose fields in [12]. The idea is to build a Rarita-Schwinger-like tensor such that

$$\langle \bar{\psi}, \mathcal{E} \rangle + \text{h.c.} = 2 \langle \bar{\psi}, \mathcal{E} \rangle, \quad (3.29)$$

up to constraints and total derivatives, starting from a general Ansatz of the type

$$\mathcal{E} = \sum_{p,q=0}^N k_{p,q} \eta^{i_1 j_1} \dots \eta^{i_p j_p} \gamma^{k_1 \dots k_q} T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q} \mathcal{S}. \quad (3.30)$$

Whereas this sum apparently differs from the one contained in eq. (3.7), in the constrained case they actually coincide, since each term gives rise to a *unique* two-column projection. Hence, making use of the result for the general γ -trace of the $\mathcal{S}$ tensor given in eq. (B.22), up to total derivatives and constraints one is led to the condition

$$\begin{aligned} \langle \bar{\psi}, \mathcal{E} \rangle + \text{h.c.} &= 2 \langle \bar{\psi}, \mathcal{E} \rangle + \sum_{p,q=0}^N \frac{i}{2^{p+1}} \langle Y_{\{2^p, 1^q\}} T_{i_1 j_1} \dots T_{i_p j_p} \bar{\psi} \gamma_{k_1 \dots k_q} , \\ &- \left[(p+q+1) k_{p,q} - \frac{(-1)^q}{q} k_{p,q-1} \right] Y_{\{2^p, 1^q\}} \gamma_{[k_1 \dots k_{q-1}} \partial_{k_q]} \prod_{r=1}^p T_{i_r j_r} \psi \\ &+ \left[k_{p,q} - (-1)^q \frac{q+1}{p} k_{p-1,q+1} \right] Y_{\{2^p, 1^q\}} \sum_{n=1}^p \gamma_{k_1 \dots k_q} (i_n \partial_{j_n}) \prod_{r \neq n}^p T_{i_r j_r} \psi \\ &+ \left[k_{p,q} + (-1)^q \frac{p+1}{q} k_{p+1,q-1} \right] \partial^l Y_{\{2^{p+1}, 1^{q-1}\}} \gamma_{[k_1 \dots k_{q-1}} T_{k_q] l} \prod_{r=1}^p T_{i_r j_r} \psi \\ &+ \left[k_{p,q} + (-1)^q (q+1)(p+q+2) k_{p,q+1} \right] \partial^l Y_{\{2^p, 1^{q+1}\}} \gamma_{k_1 \dots k_q l} \prod_{r=1}^p T_{i_r j_r} \psi \rangle, \end{aligned} \quad (3.31)$$

so that the remainder vanishes precisely if the coefficients $k_{p,q}$ are those of eq. (3.28). To reiterate, the coefficients for the general fermionic Lagrangians can be derived in two distinct ways: enforcing the gauge symmetry via the Bianchi identities along the lines of what was done for the symmetric case in [17, 19], or alternatively requiring that the kinetic operator be self adjoint, as was done for Bose fields in [12]. We shall see again shortly that the first method appears more natural in the unconstrained case.

3.2 THE UNCONSTRAINED LAGRANGIANS

In order to build the unconstrained Lagrangians, barring some subtleties that we shall elaborate upon, one can follow the same steps that led to their constrained counterparts. The new ingredients are two kinds of terms that were previously discarded, those annihilated by the Labastida constraints on the gauge parameter (3.10) and those generated by the classical anomaly (3.2) that emerges on the right-hand side of the Bianchi identities (3.1) for unconstrained fields. In the unconstrained setting, however, the first class of terms is proportional to the gauge transformation of the $\xi_{ij}(\Psi)$ introduced in Section 2.1, and can thus be balanced by suitable compensator terms while, as we saw already in the two-family case, the Bianchi identities become

$$\mathcal{B}_i : \quad \partial_i \mathcal{W} - \frac{1}{2} \not{\partial} \mathcal{W}_i - \frac{1}{2} \partial^j T_{ij} \mathcal{W} - \frac{1}{6} \partial^j \gamma_{ij} \mathcal{W} = \frac{1}{2} \partial^j \partial^k \mathcal{Z}_{ijk}, \quad (3.32)$$

with $\mathcal{Z}_{ijk}$ the gauge-invariant constraint tensors introduced in eq. (2.16). Clearly, the left-hand sides of the successive γ -traces of eq. (3.32) take the same form as the γ -traces (C.20) of the constrained Bianchi identities while, as for two-family fields, the right-hand sides build combinations of the $\mathcal{Z}_{ijk}$ tensors to be eventually compensated by the gauge transformations of some Lagrange multipliers, that here we shall denote by λ_{ijk} .

As for the Bose fields discussed in [2], the natural starting point to build the unconstrained Lagrangians is the counterpart of eq. (3.7), namely a trial Lagrangian containing only the two-column projected combinations of γ -traces of the kinetic tensor $\mathcal{W}$ of eq. (2.13), that are not proportional to the constraint tensors $\mathcal{Z}_{ijk}$,

$$\mathcal{L}_0 = \frac{1}{2} \langle \bar{\psi}, \sum_{p,q=0}^N k_{p,q} \eta^p \gamma^q (\gamma^{[q]} \mathcal{W}^{[p]}) \rangle + \text{h.c.}, \quad (3.33)$$

where the $k_{p,q}$ are given in eq. (3.28). Starting from eq. (3.7), that has the same form as eq. (3.33), in the constrained case we had to impose the Labastida constraints (3.10) on the gauge parameters to annihilate all three-column projections in the gauge variation (3.9). In the unconstrained setting, this remainder can instead be compensated introducing a new term,

$$\begin{aligned} \mathcal{L}_1 = & -\frac{1}{4} \langle \bar{\xi}_{ij}, \sum_{p,q=0}^N \eta^{i_1 j_1} \dots \eta^{i_p j_p} \gamma^{k_1 \dots k_q} \times \\ & \times \left\{ \frac{(p+1)k_{p+1,q-1}}{p+q+2} \partial_{(k_1 |} (\gamma^{[q-1]} \mathcal{W}^{[p+1]})_{|ij), i_1 j_1, \dots, i_p j_p; k_2 \dots k_q} \right. \\ & \left. + \frac{2p(p-1)qk_{p,q+1}}{(p+1)(p+q+3)} \partial_{(i_1 |} (\gamma^{[q+1]} \mathcal{W}^{[p]})_{|ij), i_2 j_2, \dots, i_p j_p; j_1 k_1 \dots k_q} \right\} \rangle + \text{h.c.}, \end{aligned} \quad (3.34)$$

whose form is determined by eq. (3.11).

The total gauge variation then takes the same form as in the constrained case, barring the replacement of $\mathcal{S}$ with $\mathcal{W}$, and can not be directly manipulated using the γ -traces of the Bianchi identities. Rather, its terms should be rearranged and joined in pairs using the unconstrained version of eq. (3.19), that reads

$$\begin{aligned} Y_{\{2^{p+1}, 1^{q-1}\}} T_{i_1 j_1} \dots T_{i_p j_p} \bar{\epsilon}_l \gamma_{k_1 \dots k_q} &= \frac{1}{q+1} Y_{\{2^{p+1}, 1^{q-1}\}} T_{i_1 j_1} \dots T_{i_p j_p} T_l [k_1 \bar{\epsilon}_{k_2} \gamma_{k_3 \dots k_q}] \\ &- \frac{2}{q+1} Y_{\{2^{p+1}, 1^{q-1}\}} T_{i_1 j_1} \dots T_{i_p j_p} \delta \bar{\xi}_l [k_1 \gamma_{k_2 \dots k_q}], \end{aligned} \quad (3.35)$$

and gives rise to a new contribution proportional to the gauge variation of the ξ_{ij} . This, in its turn, can be compensated adding to the Lagrangian

$$\begin{aligned} \mathcal{L}_2 = & -\frac{1}{4} \langle \bar{\xi}_{ij}, \sum_{p,q=0}^N \eta^{i_1 j_1} \dots \eta^{i_p j_p} \gamma^{k_1 \dots k_q} \times \\ & \times \left\{ \frac{2(q+1)k_{p,q+1}}{q+2} Y_{\{2^{p+1}, 1^q\}} \partial_{(i |} (\gamma^{[q+1]} \mathcal{W}^{[p]})_{|j_1, \dots, i_p j_p; |j) k_1 \dots k_q} \right. \\ & \left. - \frac{2(p+1)(q+1)k_{p+1,q+1}}{q+2} \partial^m (\gamma^{[q+1]} \mathcal{W}^{[p+1]})_{i_1 j_1, \dots, i_p j_p, ij; m k_1 \dots k_q} \right\} \rangle + \text{h.c.}, \end{aligned} \quad (3.36)$$

which leads to the final gauge variation

$$\begin{aligned}
\delta(\mathcal{L}_0 + \mathcal{L}_1 + \mathcal{L}_2) = & - \sum_{p,q=0}^N \frac{1}{2^{p+1}} \frac{k_{p,q}}{(q+3)(p+q+2)} \langle T_{i_1 j_1} \dots T_{i_p j_p} \bar{\epsilon}_l \gamma_{k_1 \dots k_q} , \\
& (q+3)(p+q+2) Y_{\{2^p, 1^{q+1}\}} \partial_l (\gamma^{[q]} \mathcal{W}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q} \\
& + Y_{\{2^p, 1^{q+1}\}} \sum_{n=1}^p \partial_{(i_n |} (\gamma^{[q+2]} \mathcal{W}^{[p-1]})_{\dots, i_r \neq n j_r \neq n, \dots; | j_n) l k_1 \dots k_q} \\
& + (-1)^{q+1} (q+3) \not{\partial} (\gamma^{[q+1]} \mathcal{W}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q l} \\
& - (q+3) \partial^m (\gamma^{[q]} \mathcal{W}^{[p+1]})_{i_1 j_1, \dots, i_p j_p, lm; k_1 \dots k_q} \\
& - \partial^m (\gamma^{[q+2]} \mathcal{W}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q lm} \rangle + \text{h.c.} , \tag{3.37}
\end{aligned}$$

which is the counterpart of eq. (3.25), since as we already stated the $k_{p,q}$ are those of eq. (3.28).

The terms that appear in eq. (3.37) can be turned into others involving the constraint tensors $\mathcal{Z}_{ijk}$ by suitable consequences of the Bianchi identities. As we already saw for two-family fields, at this stage a new subtlety emerges with respect to the constrained setting. For a gradient term with the structure of the last contribution in eq. (3.37), a $\{2^p, 1^{q+1}\}$ projection in the free indices does not suffice to enforce a two-column projection extended to the additional index m . Hence, in order to reproduce the terms that enter the final gauge variation, one is to consider suitable linear combinations of the projected γ -traces of the Bianchi identities. The end result, that we shall prove in Appendix C, is

$$\begin{aligned}
& \frac{p+q+3}{3} \left\{ (q+3)(p+q+2) Y_{\{2^p, 1^{q+1}\}} \partial_l (\gamma^{[q]} \mathcal{W}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q} \right. \\
& + Y_{\{2^p, 1^{q+1}\}} \sum_{n=1}^p \partial_{(i_n |} (\gamma^{[q+2]} \mathcal{W}^{[p-1]})_{\dots, i_r \neq n j_r \neq n, \dots; | j_n) l k_1 \dots k_q} \\
& + (-1)^{q+1} (q+3) \not{\partial} (\gamma^{[q+1]} \mathcal{W}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q l} \\
& - (q+3) \partial^m (\gamma^{[q]} \mathcal{W}^{[p+1]})_{i_1 j_1, \dots, i_p j_p, lm; k_1 \dots k_q} \\
& \left. - \partial^m (\gamma^{[q+2]} \mathcal{W}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q lm} \right\} = \\
& = Y_{\{2^p, 1^{q+1}\}} \left\{ (p+2)(q+3) T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q} \partial^m \partial^n \mathcal{Z}_{lmn} \right. \\
& \left. - \sum_{n=1}^p \prod_{r \neq n}^p T_{i_r j_r} \gamma_{k_1 \dots k_q l (i_n |} \partial^m \partial^n \mathcal{Z}_{| j_n) mn} \right\} . \tag{3.38}
\end{aligned}$$

In conclusion, the gauge variation (3.37) can be cast in the form

$$\begin{aligned}
\delta(\mathcal{L}_0 + \mathcal{L}_1 + \mathcal{L}_2) = & - \sum_{p,q=0}^N \frac{1}{2^{p+2}} \frac{3 k_{p,q}}{(q+3)(p+q+2)(p+q+3)} \times \\
& \times \langle Y_{\{2^p, 1^{q+1}\}} T_{i_1 j_1} \dots T_{i_p j_p} \bar{\epsilon}_l \gamma_{k_1 \dots k_q} , (p+2)(q+3) T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q} \partial^m \partial^n \mathcal{Z}_{lmn} \tag{3.39} \\
& - \sum_{n=1}^p \prod_{r \neq n}^p T_{i_r j_r} \gamma_{k_1 \dots k_q l (i_n |} \partial^m \partial^n \mathcal{Z}_{| j_n) mn} \rangle + \text{h.c.} ,
\end{aligned}$$

and can be compensated by the Lagrange multiplier terms

$$\mathcal{L}_3 = \frac{1}{4} \langle \bar{\lambda}_{ijk}, \mathcal{Z}_{ijk} \rangle + \text{h.c.}, \quad (3.40)$$

where the λ_{ijk} transform as

$$\begin{aligned} \delta \lambda_{ijk} = & \sum_{p,q=0}^N \eta^{m_1 n_1} \dots \eta^{m_p n_p} \gamma^{l_1 \dots l_q} \times \\ & \times \partial_{(i} \partial_{j|} \left\{ \frac{(p+2) k_{p,q}}{(p+q+2)(p+q+3)} Y_{\{2^p, 1^{q+1}\}} T_{m_1 n_1} \dots T_{m_p n_p} \gamma_{l_1 \dots l_q} \epsilon_{|k)} \right. \\ & \left. + \frac{(-1)^{q+1} (p+1) k_{p+1, q-2}}{q(q+1)(p+q+1)(p+q+2)} Y_{\{2^{p+1}, 1^{q-1}\}} T_{m_1 n_1} \dots T_{m_p n_p} T_{|k} \gamma_{l_1 \dots l_{q-1}} \epsilon_{l_q} \right\}. \end{aligned} \quad (3.41)$$

For brevity, this expression is not presented here in its normal ordered form, but the reader can verify that for two families it reduces to eq. (2.65).

Summarizing, the final result for the *unconstrained* fermionic Lagrangians of fields carrying an arbitrary number of families is

$$\begin{aligned} \mathcal{L} = & \frac{1}{2} \langle \bar{\psi}, \sum_{p,q=0}^N \frac{(-1)^{p+\frac{q(q+1)}{2}}}{p! q! (p+q+1)!} \eta^p \gamma^q (\gamma^{[q]} \mathcal{W}^{[p]}) \rangle + \frac{1}{4} \langle \bar{\xi}_{ij}, \Xi_{ij} \rangle \\ & + \frac{1}{4} \langle \bar{\lambda}_{ijk}, \mathcal{Z}_{ijk} \rangle + \text{h.c.}, \end{aligned} \quad (3.42)$$

with

$$\begin{aligned} \Xi_{ij} = & - \sum_{p,q=0}^N \eta^{i_1 j_1} \dots \eta^{i_p j_p} \gamma^{k_1 \dots k_q} \times \\ & \times \left\{ \frac{(-1)^{(p+1)+\frac{q(q-1)}{2}}}{p!(q-1)!(p+q+2)!} \partial_{(k_1 |} (\gamma^{[q-1]} \mathcal{W}^{[p+1]})_{|ij), i_1 j_1, \dots, i_p j_p; k_2 \dots k_q} \right. \\ & + \frac{2(p-1)pq(-1)^{p+\frac{(q+1)(q+2)}{2}}}{(p+1)!(q+1)!(p+q+3)!} \partial_{(i_1 |} (\gamma^{[q+1]} \mathcal{W}^{[p]})_{|ij), i_2 j_2, \dots, i_p j_p; j_1 k_1 \dots k_q} \\ & + \frac{2(q+1)(-1)^{p+\frac{(q+1)(q+2)}{2}}}{p!(q+2)!(p+q+2)!} Y_{\{2^{p+1}, 1^q\}} \partial_{(i |} (\gamma^{[q+1]} \mathcal{W}^{[p]})_{i_1 j_1, \dots, i_p j_p; |j) k_1 \dots k_q} \\ & \left. - \frac{2(q+1)(-1)^{(p+1)+\frac{(q+1)(q+2)}{2}}}{p!(q+2)!(p+q+3)!} \partial^m (\gamma^{[q+1]} \mathcal{W}^{[p+1]})_{i_1 j_1, \dots, i_p j_p, ij; m k_1 \dots k_q} \right\}, \end{aligned} \quad (3.43)$$

and where the gauge transformations of the Lagrange multipliers are given in eq. (3.41).

As in the two-family case, the projected Bianchi identities (3.38) and the compensator terms share a portion of their Young projections, and can thus be combined. The end result is the option, analyzed in detail for two-family fields, of modifying at the same time compensator terms and gauge transformations of the multipliers without altering the form of the Rarita-Schwinger-like tensor and the types of Young projections appearing in the Lagrangians. All this has no analogue in the bosonic case.

Furthermore, as we already saw in Section 2.1, the unconstrained Lagrangians (3.42) can be also presented in an alternative form that proves convenient when deriving the field equations. To this end, one can make a first step in the direction of the unconstrained theory considering Lagrangians for fields ψ *not* subject to the Labastida constraints (3.2) but *only* invariant under constrained gauge transformations. These can be obtained adding to the constrained Lagrangians a set of multipliers enforcing on-shell the Labastida constraints, and read

$$\mathcal{L}_C = \frac{1}{2} \langle \bar{\psi}, \sum_{p,q=0}^N \frac{(-1)^{p+\frac{q(q+1)}{2}}}{p!q!(p+q+1)!} \eta^p \gamma^q (\gamma^{[q]} \mathcal{S}^{[p]}) \rangle + \frac{i}{12} \langle \bar{\zeta}_{ijk}, T_{(ij} \psi_{k)} \rangle + \text{h.c.} \quad (3.44)$$

In this way the two classes of Lagrangians (3.7) and (3.44) give rise to two sets of equations of motion that are completely equivalent while, in order to guarantee the gauge invariance of (3.44), the ζ_{ijk} Lagrange multipliers are to transform as the λ_{ijk} did in eq. (3.41). Therefore, the unconstrained Lagrangians of eq. (3.42) can be recovered from eq. (3.44) via the Stueckelberg-like shifts

$$\begin{aligned} \psi &\rightarrow \psi - \partial^i \Psi_i, \\ \zeta_{ijk} &\rightarrow \lambda_{ijk} - \Delta_{ijk}(\Psi), \end{aligned} \quad (3.45)$$

where the Ψ_i are the compensator fields that we had introduced in Section 2.1 and the $\Delta_{ijk}(\Psi)$ that appear in the shift of the Lagrange multipliers ζ_{ijk} can be formally obtained replacing ϵ_i with Ψ_i in the gauge variation of the Lagrange multipliers λ_{ijk} , so that the ζ_{ijk} are mapped into a gauge invariant combination by eq. (3.45). A detailed discussion of the subtleties entailed by this step can be found in Section 2.2.

Both derivations of the unconstrained Lagrangians (3.42) rest on a suitable class of successive traces of the Bianchi identities but, as was already stressed for Bose fields in [2], there is further freedom to redefine simultaneously the Rarita-Schwinger-like tensor and the λ_{ijk} or ζ_{ijk} Lagrange multipliers. For instance, the Lagrangian for an unconstrained field ψ that only allows constrained gauge transformations can be also presented in the form

$$\tilde{\mathcal{L}}_C = \frac{1}{2} \langle \bar{\psi}, \sum_{p,q=0}^N \frac{(-1)^{p+\frac{q(q+1)}{2}}}{p!q!(p+q+1)!} \eta^p \gamma^q \hat{\mathcal{S}}^{[p,q]} \rangle + \frac{i}{12} \langle \bar{\mathcal{Y}}_{ijk}, T_{(ij} \psi_{k)} \rangle + \text{h.c.}, \quad (3.46)$$

where the $\mathcal{Y}_{ijk}$ are *gauge invariant* Lagrange multipliers and the $\hat{\mathcal{S}}^{[p,q]}$ are two-column projected γ -traces of $\mathcal{S}$ deprived of the $\{3, 2^{p-1}, 1^q\}$ component originally present in the last term of eq. (B.22)²¹,

$$\begin{aligned} \hat{\mathcal{S}}^{[p,q]}_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q} &\equiv i (-1)^q Y_{\{2^p, 1^q\}} \left\{ (q+1) \not{\partial} T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q} \psi \right. \\ &\quad - (p+q+1) \partial_{[k_1} T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_2 \dots k_q]} \psi - \sum_{n=1}^p \partial_{(i_n} \prod_{r \neq n}^p T_{i_r j_r} \gamma_{|j_n} k_1 \dots k_q \psi \\ &\quad \left. + \partial^m T_{i_1 j_1} \dots T_{i_p j_p} T_{m[k_1} \gamma_{k_2 \dots k_q]} \psi - \partial^m Y_{\{2^p, 1^{q+1}\}} T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{m k_1 \dots k_q} \psi \right\}. \end{aligned} \quad (3.47)$$

In complete analogy with the bosonic case of [2], the $\hat{\mathcal{S}}^{[p,q]}$ satisfy two-column projected “anomaly free” Bianchi identities that have the form of eq. (3.26) and, as a result, do not

²¹Using the techniques of Appendix C, the reader can verify that the other gradient term is already two-column projected, thanks to the projection induced in its free indices.

contain any terms related to the constraints. Looking back at the derivation of the Lagrangians presented in the previous pages, the reader can easily recognize that the Lagrangians (3.46) are gauge invariant, and indeed the Rarita-Schwinger-like tensor in eq. (3.46) is manifestly self-adjoint. One can then recover Lagrangians for fully unconstrained gauge fields performing the Stueckelberg-like shift

$$\psi \rightarrow \psi - \partial^i \Psi_i. \quad (3.48)$$

The result, however, differs from eq. (3.42) by a field redefinition of the Lagrange multipliers λ_{ijk} , while the constrained gauge invariance of (3.46) guarantees that the Ψ_i compensators still enter only via the combinations $\xi_{ij}(\Psi)$.

Let us notice, to conclude, that the terms involving the Lagrange multipliers have precisely the type of structure identified in the two-family case. The arguments presented there still apply, so that when working with the independent Ψ_i compensators all forms of the N -family Lagrangians are invariant provided the Lagrange multipliers transform like

$$\delta \lambda_{ijk} = \eta^{lm} M_{ijk,lm} + \gamma^{lm} N_{ijk;lm}, \quad (3.49)$$

where $M_{ijk,lm}$ and $N_{ijk;lm}$ are $\{4,1\}$ projected in their family indices and are mutually related according to

$$N_{ij(k;l)m} + N_{kl(i;j)m} = -\frac{1}{2} (M_{ijm,kl} + M_{klm,ij}). \quad (3.50)$$

3.3 THE FIELD EQUATIONS

As was the case for generic Bose fields [2], and as we saw for two-family Fermi fields in Section 2.2, recovering the Lagrangian (3.42) from eq. (3.44) via Stueckelberg-like shifts proves particularly convenient when deriving the field equations. In fact, retracing the derivation of the constrained Lagrangian via the request of self adjointness and keeping track of the constraint terms, one can conveniently obtain the equations of motion following from eq. (3.44):

$$\begin{aligned} E_{\bar{\psi}} : & \sum_{p,q=0}^N k_{p,q} \eta^{i_1 j_1} \dots \eta^{i_p j_p} \gamma^{k_1 \dots k_q} (\gamma^{[q]} \mathcal{S}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q} \\ & - \frac{i}{2} \sum_{p,q=0}^N (-1)^q k_{p,q} \eta^{i_1 j_1} \dots \eta^{i_p j_p} \gamma^{k_1 \dots k_q} Y_{\{2^p, 1^q\}} \partial^m Y_{\{3, 2^{p-1}, 1^q\}} T_{i_1 j_1} \dots T_{i_p j_p} \gamma_m k_1 \dots k_q \psi \\ & - \frac{1}{2} \eta^{ij} \gamma^k \mathcal{Y}_{ijk} = 0, \end{aligned} \quad (3.51)$$

$$E_{\bar{\zeta}} : \frac{i}{12} T_{(ij} \psi_{k)} = 0, \quad (3.52)$$

where the ψ terms contained in the $\mathcal{Y}_{ijk}$ originate from the hermitian conjugates of those present in the second line of eq. (3.51). After some algebra, one can recast the $\mathcal{Y}_{ijk}$ in the form

$$\begin{aligned} \mathcal{Y}_{ijk} \equiv & i \zeta_{ijk} + i \sum_{p,q=0}^N \frac{(-1)^{p+\frac{q(q+1)}{2}+1}}{p!q!(p+q+3)!} \eta^{m_1 n_1} \dots \eta^{m_p n_p} \gamma^{l_1 \dots l_q} \times \\ & \times \partial_{(i} Y_{\{2^{p+1}, 1^q\}} T_{|jk)} T_{m_1 n_1} \dots T_{m_p n_p} \gamma_{l_1 \dots l_q} \psi, \end{aligned} \quad (3.53)$$

and the arguments summarized in Appendix C then show that, due to the $\{3, 2^{p-1}, 1^q\}$ projection, the whole second line in eq. (3.51) vanishes on-shell on account of eq. (3.52). As a consequence, the equation of motion obtained from the constrained Lagrangian (3.7) only contains the first line of eq. (3.51) while, performing the Stueckelberg-like shift (3.45) as we did in Section 2.2, one can recover the equations of motion for the Lagrangian (3.42), that can be finally cast in the form

$$E_{\bar{\psi}} : \sum_{p,q=0}^N \frac{(-1)^{p+\frac{q(q+1)}{2}}}{p!q!(p+q+1)!} \eta^{i_1 j_1} \dots \eta^{i_p j_p} \gamma^{k_1 \dots k_q} (\gamma^{[q]} \mathcal{W}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q} - \frac{1}{2} \eta^{ij} \gamma^k \mathcal{Y}_{ijk} = 0, \quad (3.54)$$

$$E_{\bar{\Psi}} : \partial_l E_{\bar{\psi}} - \sum_{p,q=0}^N \frac{3(q+1)(q+2)(-1)^{p+\frac{q(q+1)}{2}}}{p!(q+3)!(p+q+3)!} \eta^{i_1 j_1} \dots \eta^{i_p j_p} \gamma^{k_1 \dots k_q} Y_{\{2p, 1^{q+1}\}} \times \\ \times \left\{ (p+2)(q+3) T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q} \partial^m \partial^n (E_{\bar{\lambda}})_{lmn} - \sum_{n=1}^p \prod_{r \neq n}^p T_{i_r j_r} \gamma_{k_1 \dots k_q l(i_n)} \partial^m \partial^n (E_{\bar{\lambda}})_{|j_n\rangle mn} \right\} = 0, \quad (3.55)$$

$$E_{\bar{\lambda}} : \frac{1}{4} \mathcal{Z}_{ijk} = 0, \quad (3.56)$$

where we have already enforced the last of these, $E_{\bar{\lambda}}$, in the field equation for $\bar{\psi}$. As usual, the equations of motion for the compensator fields $\bar{\Psi}_i$ simply guarantee the conservation of external currents coupled to the field ψ , while the $\mathcal{Y}_{ijk}$ tensors can be built performing the Stueckelberg shift (3.45) in eq. (3.53), barring the subtleties discussed in detail in Section 2.2.

3.4 COMMENTS ON THE ON-SHELL REDUCTION TO $\mathcal{S} = 0$

The next step would be the reduction of the various choices for the field equations presented in the previous section to the Labastida form $\mathcal{S} = 0$. For N -family fields, the operators on which the procedure rests are $gl(N)$ counterparts of those exhibited in Section 2.3, and the experience developed with two-family fields suggests that, in low enough dimensions, similar types of phenomena related to Weyl-like symmetries should emerge.

In the following we shall propose a set of conditions identifying the fields with this type of behavior, working directly at the level of the field equations, but we shall content ourselves with solving them explicitly in Section 3.4.2 for simple but instructive classes of examples. On the other hand, in analogy with the bosonic case discussed in [2], one can argue rather simply that in the $D \rightarrow \infty$ limit N -family gauge fields exhibit a universal behavior, with current exchanges that are free of special poles, that can be captured ignoring altogether the intricacies brought about by the S^i_j operators. In this limit the reduction proceeds along the lines of the symmetric case, and the field equations can be directly turned into $\mathcal{S} = 0$. Moreover, there are all reasons to expect that, even for higher values of N , the special behavior is confined to dimensions $D \leq 2N + 1$, where the corresponding models should be at most dual to other representations characterized by lower values of N , although we do not have a complete argument to this effect. The wider class of solutions that will be exhibited for N -family fields in Section 4.1 vindicate these expectations.

Before coming to these points, however, we would like to dwell briefly upon the very meaning of the Labastida equation $\mathcal{S} = 0$, showing that it propagates in general representation of the little group $O(D-2)$, and thus the expected numbers of degrees of freedom. Combining this step with the results of the reduction procedure one can thus conclude that the Lagrangian systems that we are proposing describe indeed the correct free dynamics of mixed-symmetry Fermi fields.

3.4.1 THE DEGREES OF FREEDOM DESCRIBED BY $\mathcal{S} = 0$

In order to proceed, it is convenient to resort momentarily to an explicit oscillator realization. This is attained introducing a set of commuting vectors $u^{i\mu}$ to cast a generic multi-symmetric gauge field ψ in the form

$$\psi \equiv \frac{1}{s_1! \dots s_N!} u^{1\mu_1} \dots u^{1\mu_{s_1}} u^{2\nu_1} \dots u^{2\nu_{s_2}} \dots \psi_{\mu_1 \dots \mu_{s_1}, \nu_1 \dots \nu_{s_2}, \dots} \quad (3.57)$$

As a result, divergences, gradients and γ 's with family indices take the form

$$\begin{aligned} \partial_i &\equiv \partial^\mu \frac{\partial}{\partial u^{i\mu}}, & \partial^i &\equiv \partial_\mu u^{i\mu}, \\ \gamma_i &\equiv \gamma^\mu \frac{\partial}{\partial u^{i\mu}}, & \gamma^i &\equiv \gamma_\mu u^{i\mu}. \end{aligned} \quad (3.58)$$

In studying the plane-wave solutions of the Labastida equation, that in momentum space takes the form

$$\not{p} \psi - (p \cdot u^i) \gamma_i \psi = 0, \quad (3.59)$$

it is convenient to decompose ψ according to

$$\psi = (p \cdot u^i) \chi_i + \hat{\psi}, \quad (3.60)$$

sorting out its portion $\hat{\psi}$ that is independent of $(p \cdot u^i)$. Substituting in (3.59) then gives

$$\not{p} \hat{\psi} - (p \cdot u^i) \gamma_i \hat{\psi} - \frac{1}{2} (p \cdot u^i) (p \cdot u^j) \gamma_{(i} \chi_{j)} = 0. \quad (3.61)$$

Now the three terms must vanish independently, since they carry different powers of $(p \cdot u^i)$, so that (3.61) is equivalent to the three conditions

$$\not{p} \hat{\psi} = 0, \quad (3.62)$$

$$\gamma_i \hat{\psi} = 0, \quad (3.63)$$

$$\gamma_{(i} \chi_{j)} = 0. \quad (3.64)$$

First of all, one can notice that (3.64) subjects χ_i to the Labastida constraints, so that the first contribution in eq. (3.60) can be gauged away altogether. One is then left with $\hat{\psi}$, that is subject to the Dirac equation (3.62). This implies the mass-shell condition $p^2 = 0$, so that one can take a momentum p with a single non-vanishing light-cone component, p^+ . As a result, the lack of dependence of $\hat{\psi}$ on $(p \cdot u^i)$ translates into the conditions

$$\frac{\partial}{\partial u^{i-}} \hat{\psi} = 0, \quad (3.65)$$

while the Dirac equation (3.62) turns into the standard projection

$$\gamma_+ \hat{\psi} = 0 , \quad (3.66)$$

that halves the on-shell components of $\hat{\psi}$. Finally, eq. (3.63) becomes

$$\left(\gamma^+ \frac{\partial}{\partial u^{i+}} + \gamma^{\underline{m}} \frac{\partial}{\partial u^{i\underline{m}}} \right) \hat{\psi} = 0 , \quad (3.67)$$

where we have used $\underline{m}$ to denote the spatial Lorentz index in order to distinguish it from the family indices. Multiplying eq. (3.67) by γ_+ and making use of eq. (3.66) one is finally led to the conditions

$$\frac{\partial}{\partial u^{i+}} \hat{\psi} = 0 , \quad (3.68)$$

$$\gamma^{\underline{m}} \frac{\partial}{\partial u^{i\underline{m}}} \hat{\psi} = 0 . \quad (3.69)$$

All components of $\hat{\psi}$ along the two light-cone directions thus are absent on account of eqs. (3.65) and (3.68), while eq. (3.69) forces the remaining transverse components to be γ -traceless. In conclusion, the Labastida equation propagates in general representations of the little group $O(D-2)$, as expected on account of its uniqueness. Let us stress that at this stage these are reducible representations, since we did not enforce any projection on ψ . In Section 4 we shall explain how to project fields and gauge parameters in order to describe irreducible representations.

We can conclude this short section by displaying a similar argument, not included in [2], that proves that the bosonic Labastida equation propagates in general representations of the little group $O(D-2)$. In the same notation used above for Fermi fields, the starting point is now the decomposition

$$\varphi = (p \cdot u^i) \zeta_i + \hat{\varphi} , \quad (3.70)$$

with $\hat{\varphi}$ independent of $(p \cdot u^i)$. Substituting in the momentum-space field equation

$$p^2 \varphi - (p \cdot u^i) \partial_i \varphi + \frac{1}{2} (p \cdot u^i) (p \cdot u^j) T_{ij} \varphi = 0 \quad (3.71)$$

one obtains

$$p^2 \hat{\varphi} - (p \cdot u^i) \partial_i \hat{\varphi} + \frac{1}{2} (p \cdot u^i) (p \cdot u^j) T_{ij} \hat{\varphi} + \frac{1}{6} (p \cdot u^i) (p \cdot u^j) (p \cdot u^k) T_{(ij} \zeta_{k)} = 0 , \quad (3.72)$$

where again the terms with different powers of $(p \cdot u^i)$ must vanish independently. As a result one is led to the conditions

$$p^2 \hat{\varphi} = 0 , \quad (3.73)$$

$$\partial_i \hat{\varphi} = 0 , \quad (3.74)$$

$$T_{ij} \hat{\varphi} = 0 , \quad (3.75)$$

$$T_{(ij} \zeta_{k)} = 0 . \quad (3.76)$$

Eq. (3.76) now implies that one can gauge away the ζ_i , since they are subject to the Labastida constraints, and eq. (3.73) then leads to the mass-shell condition, so that one can let again

$p = p^+$. $\hat{\varphi}$ is by construction independent of $(p \cdot u^i)$, which now translates into its independence from u^{i-} . When combined with eq. (3.74), this implies that $\hat{\varphi}$ has only transverse components, while finally eq. (3.75) implies that all traces of $\hat{\varphi}$ vanish. In conclusion, one is led once more to representations of the little group $O(D-2)$.

Notice that neither the triple γ -trace constraints for Fermi fields, nor the double trace constraints for Bose fields, were needed to come to this point directly from the non-Lagrangian equations. This is in contrast with what we shall see in Section 4, where they will have a direct bearing on a different, *off-shell* counting of degrees of freedom.

3.4.2 WEYL-LIKE SYMMETRIES

In Section 2.3 we classified the two-family fields whose equations of motion do not reduce directly to the Labastida form $\mathcal{S} = 0$. We saw that this can happen only if the $\mathcal{S}$ spinor-tensors satisfy special conditions, and in order to pinpoint their nature we looked for solutions preserving both the Bianchi identities and the triple γ -trace constraints induced on $\mathcal{S}$ by the corresponding constraints on the ψ fields. Finally, in Section 2.3.1 we showed explicitly that in *all* these cases new Weyl-like gauge transformations emerge, that can be used to complete the reduction to $\mathcal{S} = 0$ by suitable gauge-fixings.

This suggests another convenient starting point to classify the sporadic cases exhibiting this type of behavior. To begin with, in complete analogy with the discussion presented for two-family fields, all forms of the equations of motion can be reduced to

$$E_{\bar{\psi}} : \sum_{p,q=0}^N \frac{(-1)^{p+\frac{q(q+1)}{2}}}{p!q!(p+q+1)!} \eta^{i_1 j_1} \dots \eta^{i_p j_p} \gamma^{k_1 \dots k_q} (\gamma^{[q]} \mathcal{S}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q} - \frac{1}{2} \eta^{ij} \gamma^k \mathcal{Y}_{ijk} = 0, \quad (3.77)$$

$$E_{\bar{\zeta}} : \frac{i}{12} T_{(ij} \psi_{k)} = 0 \implies T_{(ij} \mathcal{S}_{k)} = 0, \quad (3.78)$$

possibly after a partial gauge fixing eliminating the compensators. Even in the N -family case eq. (3.77) generally sets to zero the γ -traceless part of $\mathcal{S}$. As a result, one can reduce the problem of classifying the fields whose equations of motion are not invertible to that of classifying the Weyl-like gauge transformations of ψ that leave invariant the γ -traceless part of $\mathcal{S}$. In fact, by construction these transformations leave invariant both the Bianchi identities and the constraints induced on $\mathcal{S}$ provided they also preserve the Labastida constraints on ψ , so that the fields allowing them are good candidates to look for cases where the direct reduction of the field equations is not possible. In the next few pages, following this approach, we shall propose a set of conditions identifying these Weyl-like shifts, and we shall also argue that they exhaust all relevant shift symmetries of the Bianchi identities. Finally, we shall argue that, as we saw explicitly for two-family fields, the resulting transformations leave invariant the equation of motion (3.77) when they are combined with proper shifts of the $\mathcal{Y}_{ijk}$, and a fortiori that the conditions leading to them classify the fields whose equations of motion do not reduce directly to the Labastida form.

The discussion in Section 2.3 suggests to consider generic Weyl-like gauge transformations of the form

$$\delta \psi = \eta^{i_1 j_1} \dots \eta^{i_p j_p} \gamma^{k_1 \dots k_q} \Theta_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q}. \quad (3.79)$$

They affect the kinetic spinor-tensor $\mathcal{S}$ according to

$$\begin{aligned}
\delta \mathcal{S} = & -i q \eta^{i_1 j_1} \dots \eta^{i_p j_p} \gamma^{k_1 \dots k_{q-1}} \partial^l \left\{ (D - q - 1) \Theta_{i_1 j_1, \dots, i_p j_p; l k_1 \dots k_{q-1}} \right. \\
& + \sum_{n=1}^p \Theta_{\dots, i_r \neq n j_r \neq n, \dots, l(i_n; j_n) k_1 \dots k_{q-1}} + 2 S^m_l \Theta_{i_1 j_1, \dots, i_p j_p; m k_1 \dots k_{q-1}} \left. \right\} \\
& - i \frac{p}{q+1} \eta^{i_1 j_1} \dots \eta^{i_{p-1} j_{p-1}} \gamma^{k_1 \dots k_{q+1}} \partial^l \Theta_{i_1 j_1, \dots, i_{p-1} j_{p-1}, l [k_1; k_2 \dots k_{q+1}]} \\
& + (-1)^q \eta^{i_1 j_1} \dots \eta^{i_p j_p} \gamma^{k_1 \dots k_q} \mathcal{S} \left(\Theta_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q} \right),
\end{aligned} \tag{3.80}$$

while in view of the detailed analysis of Section 2.3 the relevant conditions should reduce eq. (3.80) to its last term. Moreover, we saw in eq. (2.104) that, for two family fields, one ought to combine different types of parameters carrying identical numbers of family indices. In the N -family case, one is thus led to consider Weyl-like shifts of the form

$$\delta \psi = \sum_{p=0}^{\left[\frac{q}{2}\right]} \eta^{i_1 j_1} \dots \eta^{i_p j_p} \gamma^{k_1 \dots k_{q-2p}} \Theta^{[p, q-2p]}_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_{q-2p}}, \tag{3.81}$$

involving q family indices distributed in all possible ways between η 's and γ 's. These affect the kinetic spinor-tensor $\mathcal{S}$ according to

$$\delta \mathcal{S} = \sum_{p=0}^{\left[\frac{q}{2}\right]} \eta^{i_1 j_1} \dots \eta^{i_p j_p} \gamma^{k_1 \dots k_{q-2p}} \mathcal{S} \left(\Theta^{[p, q-2p]}_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_{q-2p}} \right), \tag{3.82}$$

provided the chains of p -dependent conditions

$$\begin{aligned}
& \frac{p+1}{(q-2p-1)(q-2p)} \Theta^{[p+1, q-2(p+1)]}_{i_1 j_1, \dots, i_p j_p, l [k_1; k_2 \dots k_{q-2p-1}]} \\
& + (D - q + 2p - 1) \Theta^{[p, q-2p]}_{i_1 j_1, \dots, i_p j_p; l k_1 \dots k_{q-2p-1}} \\
& + \sum_{n=1}^p \Theta^{[p, q-2p]}_{\dots, i_r \neq n j_r \neq n, \dots, l(i_n; j_n) k_1 \dots k_{q-2p-1}} \\
& + 2 S^m_l \Theta^{[p, q-2p]}_{i_1 j_1, \dots, i_p j_p; m k_1 \dots k_{q-2p-1}} = 0
\end{aligned} \tag{3.83}$$

hold for the allowed values of q . Actually, when N index families are present p cannot take in general all values between zero and the integer part of $q/2$, so that the relevant range for eq. (3.83) is

$$\max \left(0, \left\lceil \frac{q-N}{2} \right\rceil \right) \leq p \leq \left\lfloor \frac{q-1}{2} \right\rfloor, \tag{3.84}$$

simply because for a given N there is an upper limit for the number of indices that can be carried by an antisymmetric product of γ -matrices. Similar considerations apply to the lower ends of the sums in eqs. (3.81) and (3.82), while their upper limits are explicitly displayed. Moreover, although in principle one should take into account all values of q not exceeding the rank $\sum_{i=1}^N s_i$ of the spinor-tensor ψ , the two-family example suggests to confine the attention to $q \leq 2N$, since the (on-shell) Labastida constraints (3.2) annihilate all combinations of more than $2N$ γ -traces acting on ψ .

An admissible shift symmetry of ψ must also preserve the Labastida constraints (3.2), that hold off-shell in the constrained theories and are enforced on-shell in the unconstrained theories by the equations of motion (3.78) for the Lagrange multipliers. Our experience with two-family fields gives us confidence that the conditions (3.83), together with the Labastida-like constraints

$$T_{(lm} \gamma_n) \Theta^{[p, q-2p]}_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_{q-2p}} = 0, \quad (3.85)$$

suffice to preserve the Labastida constraints on ψ in eq. (3.78), although we have not carried out in full detail the type of analysis leading to eqs. (2.119), (2.120) and (2.121) for the general shifts of eq. (3.81). Once the Labastida constraints on ψ are satisfied, similar constraints on $\mathcal{S}$ hold by construction, and moreover the shifts (3.82) are bound to preserve the Bianchi identities, simply because they originate from ψ shifts. Furthermore, the $\mathcal{S}(\Theta)$ satisfy the same set of conditions (3.83) as the Θ parameters, since the S^i_j operators commute with $\mathcal{S}$.

On the other hand, one can sharpen the correspondence with the analysis of Section 2.3 considering directly the shifts

$$\delta \mathcal{S} = \eta^{i_1 j_1} \dots \eta^{i_p j_p} \gamma^{k_1 \dots k_q} \Omega_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q} \quad (3.86)$$

of the Labastida spinor-tensors $\mathcal{S}$, where for simplicity we consider a single term of the sum corresponding to eq. (3.82). They lead to the variations

$$\begin{aligned} \delta \mathcal{B}_l = & -\frac{q}{2} \not{\partial} \eta^{i_1 j_1} \dots \eta^{i_p j_p} \gamma^{k_1 \dots k_{q-1}} \left\{ (D - q - 1) \Omega_{i_1 j_1, \dots, i_p j_p; l k_1 \dots k_{q-1}} \right. \\ & + \sum_{n=1}^p \Omega_{\dots, i_r \neq n j_r \neq n, \dots, l(i_n; j_n) k_1 \dots k_{q-1}} + 2 S^m_l \Omega_{i_1 j_1, \dots, i_p j_p; m k_1 \dots k_{q-1}} \left. \right\} \\ & - \frac{1}{2} \frac{p}{q+1} \not{\partial} \eta^{i_1 j_1} \dots \eta^{i_{p-1} j_{p-1}} \gamma^{k_1 \dots k_{q+1}} \Omega_{i_1 j_1, \dots, i_{p-1} j_{p-1}, l[k_1; k_2 \dots k_{q+1}]} + \dots \\ & + \eta^{i_1 j_1} \dots \eta^{i_p j_p} \gamma^{k_1 \dots k_q} \mathcal{B}_l(\Omega_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q}) \end{aligned} \quad (3.87)$$

of the Bianchi identities, where the omitted terms contain overall gradients. Notice that the $\not{\partial}$ terms have the same structure as the terms within brackets in eq. (3.80), and are thus eliminated by the conditions (3.83), that are hence plausibly not only sufficient but also necessary to preserve the Bianchi identities. All gradient terms, that we have not displayed for brevity in (3.87), must in fact disappear if eq. (3.83) holds, since they cannot cancel against the last term, that vanishes if $\Omega \sim \mathcal{S}(\Theta)$ with a Θ -parameter satisfying eq. (3.85), so that *a fortiori* the Ω -parameters are also to satisfy the Bianchi identities. In conclusion, if as we expect the conditions (3.83) preserve the Labastida constraints when they are combined with eq. (3.85), the transformations (3.82) share all the key features of the Ansatz that in Section 2.3 made it possible to classify the sporadic two-family cases with pathological reductions. This should not surprise the reader: looking for shift symmetries that preserve the Bianchi identities as we did in Section 2.3 is equivalent to looking for shift symmetries of the field equations that can be realized directly in terms of ψ .

The next issue is to understand whether the shifts of $\mathcal{S}$ that we have identified lead to symmetries of the field equation (3.77), possibly when combined with proper shifts of the $\mathcal{Y}_{ijk}$ spinor-tensors. Since the structure of the Rarita-Schwinger-like tensor is fully determined by the Bianchi identities and their γ -traces, the promotion of the symmetries of the Bianchi identities to symmetries of the equation of motion for $\bar{\psi}$ is really expected. For instance, one can neatly recover it for the simplest Weyl-like transformation of the Labastida tensor,

$$\delta \mathcal{S} = \gamma^i \Omega_i, \quad (3.88)$$

since our previous analysis selects Ω_i parameters satisfying

$$(D - 2) \Omega_i + 2 S^j{}_i \Omega_j = 0, \quad (3.89)$$

while eq. (3.88) induces the variation

$$\begin{aligned} \delta \mathcal{E} = & \sum_{p,q=0}^N k_{p,q} \eta^{i_1 j_1} \dots \eta^{i_p j_p} \gamma^{k_1 \dots k_q} [Y_{\{2^p, 1^q\}} T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q}, \gamma^l]_{(-1)^{q+1}} \Omega_l \\ & + \sum_{p,q=0}^N \frac{(-1)^q}{q+1} k_{p,q} \eta^{i_1 j_1} \dots \eta^{i_p j_p} \gamma^{k_1 \dots k_{q+1}} Y_{\{2^p, 1^{q+1}\}} T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{[k_1 \dots k_q} \Omega_{k_{q+1}]} \\ & + \sum_{p,q=0}^N \frac{(-1)^q q}{p+1} k_{p,q} \eta^{i_1 j_1} \dots \eta^{i_{p+1} j_{p+1}} \gamma^{k_1 \dots k_{q-1}} Y_{\{2^{p+1}, 1^{q-1}\}} \sum_{n=1}^{p+1} \prod_{r \neq n}^p T_{i_r j_r} \gamma_{k_1 \dots k_{q-1}} (i_n \Omega_{j_n}) \\ & + \sum_{p,q=0}^N (-1)^q k_{p,q} \eta^{i_1 j_1} \dots \eta^{i_p j_p} \gamma^{k_1 \dots k_q} \gamma^l Y_{\{3, 2^{p-1}, 1^q\}} (Y_{\{2^p, 1^q\}} T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q}) \Omega_l \quad (3.90) \end{aligned}$$

of the Rarita-Schwinger-like tensor $\mathcal{E}$ of eq. (3.30). In the variation of eq. (3.77), the last line in eq. (3.90) can be compensated by Ω -dependent shifts of the $\mathcal{Y}_{ijk}$ tensors, on account of its three-column projection. On the other hand, the first three lines should vanish directly, and they indeed do provided they combine to rebuild eq. (3.89). Starting from the rules collected in Appendix A, one can indeed prove that

$$\begin{aligned} [T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q}, \gamma^l]_{(-1)^{q+1}} \Omega_l = & (-1)^q \sum_{n=1}^p \prod_{r \neq n}^p T_{i_r j_r} T_{(i_n | [k_1 \gamma_{k_2 \dots k_q}]} \Omega_{|j_n)} \quad (3.91) \\ & + T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{[k_1 \dots k_{q-1}]} \left\{ (D + q - 1) \Omega_{|k_q]} + 2 S^l{}_{|k_q]} \Omega_l \right\} + \sum_{n=1}^p \prod_{r \neq n}^p T_{i_r j_r} \gamma_{k_1 \dots k_q} (i_n \Omega_{j_n}), \end{aligned}$$

so that, restricting the attention to two-column Young projections, one can finally show that

$$\begin{aligned} [Y_{\{2^p, 1^q\}} T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q}, \gamma^l]_{(-1)^{q+1}} \Omega_l = & Y_{\{2^p, 1^q\}} \sum_{n=1}^p \prod_{r \neq n}^p T_{i_r j_r} \gamma_{k_1 \dots k_q} (i_n \Omega_{j_n}) \quad (3.92) \\ & + Y_{\{2^p, 1^q\}} T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{[k_1 \dots k_{q-1}]} \left\{ (D + p + q - 1) \Omega_{|k_q]} + 2 S^l{}_{|k_q]} \Omega_l \right\}. \end{aligned}$$

Substituting this identity in eq. (3.90) and combining the terms containing the same invariant tensors finally gives

$$\begin{aligned} \delta \mathcal{E} = & \sum_{p,q=0}^N k_{p,q} \eta^{i_1 j_1} \dots \eta^{i_p j_p} \gamma^{k_1 \dots k_q} \times \quad (3.93) \\ & \times Y_{\{2^p, 1^q\}} T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{[k_1 \dots k_{q-1}]} \left\{ (D - 2) \Omega_{|k_q]} + 2 S^l{}_{|k_q]} \Omega_l \right\} + \dots, \end{aligned}$$

where we have omitted the last line of eq. (3.90), since we have already stressed that proper shifts of the $\mathcal{Y}_{ijk}$ tensors can cancel it. This expression extends the two-family result of eq. (2.122) and shows clearly that the field equation (3.77) is left invariant by the transformation (3.88) with parameters subject to the conditions (3.89). It is reasonable to expect that similar results hold

for the generic Weyl-like shifts (3.82) whose parameters are subject to the conditions (3.83), but we do not have a complete proof of this fact, although concrete indications to this effect can be found in Appendix D.

Let us conclude by stressing that the emergence of Weyl-like symmetries could be also investigated directly at the level of the action, and a brief digression on this matter is perhaps instructive. First of all, let us stress that one should also take into account the degenerate cases in which the tensor $\mathcal{S}$ (or, more generally, one of its γ -traces $T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q} \mathcal{S}$) vanishes identically. These are not captured by our previous analysis that was just meant to identify the conditions under which the equation $\mathcal{E} = 0$ *does not* reduce to $\mathcal{S} = 0$. Actually, we already came across an example of this sort in Section 2.3, an irreducible $\{1, 1\}$ field in two dimensions. And indeed, as we shall explain in some detail in Appendix D, $\mathcal{S} \equiv 0$ for all fully antisymmetric spinor-tensors carrying N space-time indices in $D = N$. Barring these special cases, we can now discuss how to restate our previous results from this viewpoint. For brevity, we shall however content ourselves with rephrasing in these terms the simplest Weyl-like symmetry involving a single γ^i , already analyzed for two-family fields in Section 2.3 and for N -family fields here. Let us therefore consider a transformation of the field ψ of the form

$$\delta \psi = \gamma^i \Theta_i. \quad (3.94)$$

The corresponding variation of the constrained Lagrangian (3.7), to which we restrict our attention for simplicity, reads then

$$\delta \mathcal{L} = -2 \langle \bar{\Theta}_i, \gamma_i \mathcal{E} \rangle, \quad (3.95)$$

thus stressing again that the emergence of Weyl-like symmetries is actually related to conditions on γ -traces of $\mathcal{E}$. Following steps similar to those leading to eq. (3.93), one can show that

$$\begin{aligned} \gamma_l \mathcal{E} &= [(D-2)\delta^m_l + 2S^m_l] \times \\ &\times \sum_{p,q=0}^N (q+1) k_{p,q+1} \eta^{i_1 j_1} \dots \eta^{i_p j_p} \gamma^{k_1 \dots k_q} Y_{\{2^p, 1^{q+1}\}} T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{m k_1 \dots k_q} \mathcal{S}, \end{aligned} \quad (3.96)$$

since the three-column projected terms are annihilated by the Labastida-like constraints (3.4) on $\mathcal{S}$. All in all, the invariance of $\mathcal{L}$ therefore translates into the conditions (3.89), that we already met for two-family fields in eq. (2.103) and that can be recovered in this fashion moving the S^i_j -dependent operator to the left in the scalar product.

4 IRREDUCIBLE FIELDS OF MIXED SYMMETRY

The gauge fields ψ considered in the previous sections were only symmetric under interchanges of pairs of vector indices belonging to the same set, but did not possess any symmetry relating different sets. In other words, they were *reducible* $gl(D)$ spinor-tensors. Fields of this type naturally emerge in String Theory, where they are associated to products of bosonic oscillators, together with more general reducible $gl(D)$ tensors, some sets of whose indices are actually antisymmetrized rather than symmetrized. We shall explain how to adapt our construction to this other class of fields in Section 5, but in the following we would like to concentrate briefly on *irreducible* $gl(D)$ spinor-tensors, still with manifestly symmetric index sets. Fields of this type can be associated to Young diagrams, and are more akin to the familiar low-spin examples.

In contrast with the more familiar (Majorana-)Weyl projections, these irreducibility conditions only affect in a non-trivial fashion the vector indices carried by the ψ fields, so that the changes induced in the present case are actually along the lines of what we already described in [2] for Bose fields. For completeness, however, we now briefly adapt the irreducibility constraints to the present fermionic setting and display their solutions. We also present an instructive counting argument for the propagating degrees of freedom for irreducible two-family spinor-tensors.

Irreducibility requires that a vanishing result obtain if one tries to symmetrize all vector indices of ψ belonging to a given family with any additional index of the lower lines of its Young diagram. In terms of the S^i_j operators introduced in Appendix A, this condition reads

$$S^i_j \psi = 0, \quad \forall i < j, \quad 2 \leq j \leq N, \quad (4.1)$$

where N is the number of families. The Fang-Fronsdal-Labastida kinetic operator (2.5) *commutes with the S^i_j operators*, as was the case for its bosonic counterpart [2], and in a similar fashion the Rarita-Schwinger-like tensors of eq. (3.30) also commute with the S^i_j operators, on account of the relations

$$[S^m_n, \eta^{i_1 j_1} \dots \eta^{i_p j_p} \gamma^{k_1 \dots k_q} T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q}] = 0, \quad (4.2)$$

or more simply because they are “family” scalars. As a consequence, in the constrained theory the Lagrangians can be presented in *exactly* the same form (3.7) and (3.28) for both reducible and irreducible gauge fields. For irreducible fields, however, not all γ -traces of ψ remain independent, and therefore it is also possible to reduce the number of terms appearing in them, although the redundant description is anyway correct and has the further advantage of being universal. An illustration of this fact may be found in Section 4 of [2].

At any rate, in general irreducible ψ fields involve fewer independent gauge parameters and constraints with respect to their reducible counterparts, and it is interesting to characterize them. The gauge transformations (2.6) take the same form as their bosonic analogues, and therefore the considerations of [2] apply verbatim. In particular, the requirement that the constraints (4.1) be preserved by gauge transformations leads to

$$\partial^k (S^i_j \epsilon_k + \delta_k^i \epsilon_j) = 0, \quad \forall i < j, \quad 2 \leq j \leq N, \quad (4.3)$$

and thus, up to gauge-for-gauge transformations, to

$$S^i_j \epsilon_k + \delta_k^i \epsilon_j = 0, \quad \forall i < j, \quad 2 \leq j \leq N. \quad (4.4)$$

As for Bose fields, the ϵ_i that solve this system can be built from the available Young-projected $\tilde{\epsilon}_i$, that exist only when the corresponding Young diagrams, obtained stripping one box from the original diagram of ψ , are admissible. With this proviso, one can solve eqs. (4.4), obtaining

$$\epsilon_{N-k} = \tilde{\epsilon}_{N-k} + \sum_{l=1}^k \sum_{i_1 > \dots > i_l \geq 0}^{k-1} [l_k]_{i_1 \dots i_l} S^{N-i_1}_{N-k} S^{N-i_2}_{N-i_1} \dots S^{N-i_l}_{N-i_{l-1}} \tilde{\epsilon}_{N-i_l}, \quad (4.5)$$

where

$$[l_k]_{i_1 \dots i_l} = \frac{1}{s_{N-i_l} - s_{N-k} + i_l - k} \prod_{h=1}^l \frac{1}{s_{N-i_l} - s_{N-i_h} + i_l - i_h}. \quad (4.6)$$

For each of the original gauge parameters ϵ_i , the expansion (4.5) in powers of the S^i_j operators thus starts with the corresponding irreducible $\tilde{\epsilon}_i$, if this exists, but in higher orders it also

receives contributions from other irreducible $\tilde{\epsilon}_{N-k}$ parameters corresponding to lower values of k . Let us stress again that the irreducible parameters are in one-to-one correspondence with the admissible Young diagrams obtained stripping *one* box from the original diagram of the spinor-tensor ψ . As a result, they are in general fewer than the number of families. For instance, a set of k lines of identical length gives rise to k distinct gauge parameters in the reducible case but to a single gauge parameter in the irreducible case. This was first pointed out in [11, 30], and here we are thus providing an explicit derivation of the result.

Eqs. (4.4) induce an identical set of conditions on the Ψ_i compensators,

$$S^m_n \Psi_i + \delta^m_i \Psi_n = 0, \quad \forall m < n, \quad 2 \leq n \leq N, \quad (4.7)$$

and computing a γ -trace of eq. (4.7) then leads to the relations satisfied by the composite compensators $\xi_{ij}(\Psi)$,

$$S^m_n \xi_{ij}(\Psi) + \delta^m_{(i} \xi_{j)n}(\Psi) = 0, \quad \forall m < n, \quad 2 \leq n \leq N. \quad (4.8)$$

In a similar fashion, applying S^m_n to the symmetrized triple γ -trace of the gauge field ψ , one can identify the conditions satisfied by the constraint tensors $\mathcal{Z}_{ijk}$, and consequently by the Lagrange multipliers λ_{ijk} ,

$$S^m_n \lambda_{ijk} + \delta^m_{(i} \lambda_{jk)n} = 0, \quad \forall m < n, \quad 2 \leq n \leq N. \quad (4.9)$$

In analogy with the result described for the gauge parameters, one can see that eqs. (4.9) associate an independent irreducible Lagrange multiplier to each admissible Young diagram obtained stripping *three* boxes from that of the original spinor-tensor gauge field. Actually, the S^i_j operators satisfy Leibnitz's rule, and as a result explicit solutions of eqs. (4.8) and (4.9) can be obtained tensoring the solutions (4.5) of eq. (4.4).

This concludes our cursory look at the restrictions on the field content in the irreducible case. As we already stated, similar conditions can be used to relate the irreducible Young projections of the various γ -traces appearing in the Rarita-Schwinger-like tensors, in complete analogy with what was illustrated in [2] for some types of Bose fields.

An important, related issue, has to do with the propagating modes of these systems. While the argument presented in Section 3.4.1 can be adapted to this irreducible case by simply enforcing on ψ the irreducibility conditions (4.1), possibly together with the available (Majorana-)Weyl conditions, it is perhaps instructive to count in detail the propagating modes, taking into account all Labastida constraints on the independent components of the field ψ and of the parameters selected by the procedure that we have just illustrated. We shall content ourselves with the case of generic irreducible two-family gauge fields ²² $\psi_{\mu_1 \dots \mu_{s_1}, \nu_1 \dots \nu_{s_2}}$, since the combinatorics behind this argument becomes readily clumsy as the number of index families increases.

Following the procedure proposed in [35], the number of physical degrees of freedom propagated by an $\{s_1, s_2\}$ Young projected two-family gauge field ψ is

$$N_{\text{d.o.f.}} = \frac{1}{2} \left(N_\psi - 3 N_{\tilde{\epsilon}_i} + 5 N_{\tilde{\epsilon}_{[ij]}} \right), \quad (4.10)$$

where N_ψ , $N_{\tilde{\epsilon}_i}$ and $N_{\tilde{\epsilon}_{[ij]}}$ denote the numbers of independent components of the constrained irreducible gauge field ψ , of the two constrained irreducible gauge parameters $\tilde{\epsilon}_i$ and of the

²²The fields could be also subject to (Majorana-)Weyl projections, but these do not play a direct role in our arguments. For brevity, we assume that $s_1 > s_2 + 1 > 2$ in order to leave out special degenerate cases, and that $D > 5$ so that the corresponding $\{s_1, s_2\}$ representations of $o(D-2)$ exist.

single constrained irreducible gauge-for-gauge parameters $\tilde{\epsilon}_{[ij]}$. In order to identify these contributions, let us begin by decomposing the spinor-tensors of interest in $o(D)$ representations. This procedure differs slightly from the one used in the examples of [12], since it focusses directly on quantities that survive the Labastida constraints (3.2) and (3.10), bypassing somehow the potential difficulties introduced by the linear dependence of the constraints.

The decompositions of $gl(D)$ unconstrained two-family gauge fields ψ and gauge parameters $\tilde{\epsilon}_i$ in terms of $o(D)$ representations would read

$$\begin{aligned} \psi : \{s_1, s_2\}_{gl} = & \{s_1, s_2\}_o \oplus \{s_1 - 1, s_2\}_o \oplus \{s_1, s_2 - 1\}_o \oplus \{s_1 - 2, s_2\}_o \oplus \{s_1, s_2 - 2\}_o \\ & \oplus 2\{s_1 - 1, s_2 - 1\}_o \oplus 2\{s_1 - 2, s_2 - 1\}_o \oplus 2\{s_1 - 1, s_2 - 2\}_o \\ & \oplus 3\{s_1 - 2, s_2 - 2\}_o \oplus \dots, \end{aligned} \quad (4.11)$$

where the gl and o subscripts are meant to stress that on the left-hand side we refer to $gl(D)$ representations, that as such include all (γ) -traces, while on the right-hand side we refer to $o(D)$ representations, where all (γ) -traces have been removed. In a similar fashion

$$\begin{aligned} \tilde{\epsilon}_1 : \{s_1 - 1, s_2\}_{gl} = & \{s_1 - 1, s_2\}_o \oplus \{s_1 - 2, s_2\}_o \oplus \{s_1 - 1, s_2 - 1\}_o \oplus \{s_1 - 1, s_2 - 2\}_o \\ & \oplus 2\{s_1 - 2, s_2 - 1\}_o \oplus \dots, \\ \tilde{\epsilon}_2 : \{s_1, s_2 - 1\}_{gl} = & \{s_1, s_2 - 1\}_o \oplus \{s_1 - 1, s_2 - 1\}_o \oplus \{s_1, s_2 - 2\}_o \oplus \{s_1 - 2, s_2 - 1\}_o \\ & \oplus 2\{s_1 - 1, s_2 - 2\}_o \oplus \dots, \end{aligned} \quad (4.12)$$

where we have refrained from displaying terms that are manifestly related to the Labastida constraints. These include all contributions to ψ obtained removing at least three boxes from the same row and all contributions to the $\tilde{\epsilon}_i$ obtained removing at least two boxes from the same row. For two-family fields it is then relatively simple to sort out in eqs. (4.11) and (4.12) the $o(D)$ components that survive the Labastida constraints. Denoting by $[s_1, s_2]$ the dimension of the $o(D)$ representation with s_1 boxes in its first row and s_2 boxes in its second row, one thus obtains

$$\begin{aligned} N_\psi = & [s_1, s_2] + [s_1 - 1, s_2] + [s_1, s_2 - 1] + [s_1 - 2, s_2] + 2[s_1 - 1, s_2 - 1] \\ & + [s_1, s_2 - 2] + [s_1 - 2, s_2 - 1] + [s_1 - 1, s_2 - 2] + [s_1 - 2, s_2 - 2] \end{aligned} \quad (4.13)$$

and

$$N_{\tilde{\epsilon}_i} = [s_1 - 1, s_2] + [s_1, s_2 - 1] + [s_1 - 1, s_2 - 1] + [s_1 - 2, s_2 - 1] + [s_1 - 1, s_2 - 2]. \quad (4.14)$$

Furthermore, the same constraints imply that the only available $\tilde{\epsilon}_{[ij]}$ is γ -traceless, so that

$$N_{\tilde{\epsilon}_{[ij]}} = [s_1 - 1, s_2 - 1]. \quad (4.15)$$

Taking into account all these contributions and making use of eq. (A.38), one is finally led to

$$N_{\text{d.o.f.}} = 2^{\lfloor \frac{D-2}{2} \rfloor} \frac{s_1 - s_2 + 1}{(s_1 + 1)! s_2!} \frac{(D + s_1 - 5)!}{(D - 4)!} \frac{(D + s_2 - 6)!}{(D - 6)!} (D + s_1 + s_2 - 4), \quad (4.16)$$

which is precisely the dimension of the $\{s_1, s_2\}$ irreducible representation of $o(D - 2)$, up to a possible overall factor reflecting a (Majorana-)Weyl projection that we have not considered for brevity. Similar results can be presented for Bose fields, and in principle one could also adapt these arguments to fields with larger numbers of index families, taking into account larger numbers of leftover $o(D)$ components and the presence of further gauge-for-gauge parameters.

4.1 AN APPLICATION: SOME LAGRANGIANS WITH WEYL-LIKE SYMMETRIES

The search for Weyl-invariant Lagrangians can be performed considering directly irreducible fields and, as we shall see shortly, the relations displayed in the previous section greatly simplify the analysis of the conditions identifying those symmetries. We thus conclude this section by displaying some non-trivial solutions of eqs. (3.83). To begin with, we can show how to solve *all* these conditions for the special case of single-column spinor-tensors carrying s vector indices. In Section 2.3.2 we presented concise forms for their Lagrangians, that vanish manifestly for $s \leq D \leq 2s$, and we noticed that the formal expressions for their field equations in terms of $\mathcal{S}$ should possess Weyl-like symmetries. We can now recover this result solving the conditions (3.83) for spinor-tensor forms. First of all, in this case the η^{ij} tensors do not enter Lagrangians and field equations, so that the chains of conditions (3.83) reduce to the far simpler independent equations

$$(D - q - 1) \Theta^{[q]}_{k_1 \dots k_{q-1} l} + 2 S^m_l \Theta^{[q]}_{k_1 \dots k_{q-1} m} = 0, \quad (4.17)$$

whose solutions identify Weyl-like shifts of the form

$$\delta \psi = \gamma^{k_1 \dots k_q} \Theta^{[q]}_{k_1 \dots k_q}. \quad (4.18)$$

Let us also stress that no other conditions are to be imposed on these parameters, because all spinor-tensor forms are *unconstrained*. Furthermore, for this type of fields $S^m_l \psi = 0$ for all $l \neq m$, since it is clearly impossible to symmetrize their space-time indices. As a result

$$0 = S^m_l \delta \psi = \gamma^{k_1 \dots k_q} \left(S^m_l \Theta^{[q]}_{k_1 \dots k_q} + (-1)^{q+1} \delta^m_{[k_1} \Theta^{[q]}_{k_2 \dots k_q] l} \right), \quad l \neq m, \quad (4.19)$$

so that eq. (4.17) becomes

$$(D - 2N + q - 1) \Theta^{[q]}_{k_1 \dots k_{q-1} l} = 0. \quad (4.20)$$

Fully antisymmetric fields can admit a single independent q -th γ -trace, that as we now see is left undetermined in $D = 2N - q + 1$, since it can be shifted by non-vanishing $\Theta^{[q]}$ parameters. For one-column fields the number N of families appearing in eq. (4.20) is equivalent to the total number s of space-time indices carried by the field ψ . Therefore, as expected, we are recovering the existence of Weyl-like symmetries for these fields for $s + 1 \leq D \leq 2s$. Notice, finally, that the Lagrangians (2.165) also vanish manifestly for $D = s$, but in these cases the $\mathcal{S}$ spinor-tensors vanish as well, as one can see computing explicitly their only available component. This explains why these special space-time dimensions do not emerge from the previous discussion. Further comments on this case can be found in Appendix D.

We can now supplement the classification of Section 2.3 of the Weyl-like symmetries emerging in the two-family case by displaying some interesting classes of irreducible N -family fields that allow Weyl-like shifts of the form

$$\delta \psi = \gamma^i \Theta_i. \quad (4.21)$$

In this case the irreducibility of ψ translates into the conditions

$$\delta^i_k \Theta_j + S^i_j \Theta_k = 0 \quad (i < j) \quad (4.22)$$

on the Θ_i , to be combined with the conditions allowing (4.21) to define a Weyl symmetry, that were given in eq. (3.89) and that we can rewrite in the convenient form²³

$$(D + 2s_i - 4) \Theta_i + 2 \sum_{j < i} S^j_i \Theta_j + 2 \sum_{j > i} S^j_i \Theta_j = 0 \quad (i = 1, \dots, N), \quad (4.23)$$

²³Here and in the rest of this section no summations are left implicit.

where s_i denotes the number of Lorentz labels in the i -th family for the spinor-tensor ψ . Eqs. (4.22) finally reduce (4.23) to

$$[D + 2s_i - 2(i + 1)] \Theta_i - 2 \sum_{j>i} S^j_i S^i_j \Theta_i = 0 \quad (i = 1, \dots, N). \quad (4.24)$$

The sum is absent in the last of these equations, that is simply

$$[D + 2s_N - 2(N + 1)] \Theta_N = 0, \quad (4.25)$$

and selects the condition

$$D + 2s_N = 2(N + 1) \quad (4.26)$$

to recover a Weyl-like symmetry with a non-vanishing Θ_N parameter. As a result, one can conclude right away that no solutions of this kind exist in odd dimensions, or if $D > 2N$, consistently with the tables of Section 2.3 and with the considerations made at the beginning of Section 3.4. Inverting the diagonal equations in (4.22) would give

$$\Theta_i = -\frac{1}{s_i - s_j + 1} S^j_i \Theta_j \quad (i < j), \quad (4.27)$$

up to possible homogeneous solutions. At any rate, if Θ_N is non-trivial, all the Θ_i for $i < N$ are also non trivial, since in our conventions $s_i \geq s_j$.

Turning to the other equations with $i < N$, one can notice that the irreducibility conditions (4.22) also imply that the operators $S^j_i S^i_j$ act *diagonally* on Θ_i . In fact, S^j_i and S^i_j are but the $L_-^{(ij)}$ and $L_+^{(ij)}$ operators for the $su(2)$ subalgebra of $gl(N)$ connecting the two rows i and j , for which

$$L_3^{(ij)} = \frac{1}{2} (S^i_i - S^j_j) \quad (4.28)$$

while eq. (4.22) implies that

$$L_+^{(ij)} \Theta_j = 0, \quad \Theta_j = -L_+^{(ij)} \Theta_i. \quad (4.29)$$

As a result, Θ_i and Θ_j have the same “total angular momentum” quantum number

$$\ell_{ij}(\Theta_i) = \ell_{ij}(\Theta_j) = \frac{s_i - s_j + 1}{2}, \quad (4.30)$$

that is also the “magnetic” quantum number of Θ_j , that lies at the tip of the $su(2)$ chain and lacks one space-time index of the j -th row when compared to the gauge field ψ . At the same time, the “magnetic” quantum number of Θ_i , that lacks an index in the i -th row compared to ψ , is

$$m_{ij}(\Theta_i) = \frac{s_i - s_j - 1}{2}, \quad (4.31)$$

so that finally

$$S^j_i S^i_j \Theta_i = (s_i - s_j + 1) \Theta_i. \quad (4.32)$$

In conclusion, the remaining equations (4.24) with $i < N$ reduce to the chain of conditions

$$\left[(N - i - 1) s_i - \sum_{j=i+1}^{N-1} s_j \right] \Theta_i = 0 \quad (i = 1, \dots, N - 2), \quad (4.33)$$

while the $(N - 1)$ -th condition is also identically satisfied on account of eq. (4.26). Notice that, as we anticipated, on account of the irreducibility all the Θ_i must be non-vanishing to define a non-trivial solution when eq. (4.26) holds. One thus identifies the $\{s, \dots, s, s_N\}$ irreducible fields with s arbitrary and $s_N \leq N$ in the space-time dimensions determined by eq. (4.26). We shall content ourselves with this general class of solutions, although others with vanishing Θ_N exist in low-enough dimensions.

The solutions that we identified are also to preserve the Labastida constraints (3.2), and in eq. (2.128) we have seen that, whenever eqs. (4.23) hold, for the Weyl-like shifts that we are considering this is guaranteed by the additional conditions

$$T_{(ij} \gamma_k) \Theta_l = 0. \quad (4.34)$$

On the other hand, the conditions (4.22) imply that all Θ_l parameters have the same $gl(D)$ structure as the irreducible Θ_N , that in this sense represents the most convenient choice for the single independent quantity left over in eq. (4.23). Therefore, the transformations of the $\{s, \dots, s, 1\}$ fields in $D = 2N$ rest on this $\{s, \dots, s, 0\}$ conformal Weyl parameter, that in this space-time dimension admits a γ -traceless component that as such cannot be affected by the triple γ -trace constraints (4.34). This is consistent with the fact that for two-family fields we did find explicitly that $\{s, 1\}$ fields admit Weyl-like symmetries of the type (4.21) compatible with the Labastida constraints. In this fashion, we have shown that Weyl-like symmetries emerge in a class of examples that are richer than fully antisymmetric spinor-tensors, and already display the key features of generic mixed-symmetry fields. On the other hand, for higher values of s_N a standard theorem already quoted in Section 2.3.2 (for which we refer the reader to the first reference in [33], §10-6) implies that a γ -traceless component does not exist for $\{s, \dots, s, s_N - 1\}$ -projected parameters in $D = 2(N - s_N + 1)$, so that the previous considerations protecting this type of shift are no more available. While we do not have a complete argument, our explicit analysis of two-family fields makes us expect that the triple γ -trace conditions eliminate at least some of these solutions (4.26) in low enough space-time dimensions. Let us close this section by stressing that even these new classes of solutions emerge in space-time dimensions $D \leq 2N + 1$, in agreement with the considerations made at the beginning of Section 3.4. Moreover, the standard theorem just recalled shows that *all* these fields do not propagate any local degrees of freedom. We do not know whether all these $\{s, \dots, s, s_N\}$ models have vanishing actions, but in Section 2.3.2 we have already seen that the simplest member of this class, a $\{2, 1\}$ field in two dimensions, does not, so that it is reasonable to expect that they all behave in the same fashion.

5 MULTI-FORM SPINOR-TENSORS

In the previous sections we have described the properties of multi-symmetric gauge fields of the type $\psi_{\mu_1 \dots \mu_{s_1}, \nu_1 \dots \nu_{s_2}, \dots}$, that in general propagate the degrees of freedom of reducible representations of the Poincaré group. As we have seen, this “redundant” reducible description is particularly convenient to manifest the structure of the algebraic constraints on fields and gauge parameters, but it is also tailored for the comparison with massive string spectra, where fields of this type are the natural counterparts of products of bosonic oscillators. Superstring spectra, however, also involve fermionic oscillators, and it is thus interesting to describe how the previous results can be adapted to the case of multi-form fermionic gauge fields, for which the indices belonging to a given family are fully antisymmetric. As we shall see shortly, the

algebraic properties of the basic operators are altered to some extent, but the structure of the whole construction remains pretty much unaffected in this variant of the formalism.

The first ingredient for moving from multi-symmetric fields to multi-forms is a proper definition of the operators that we used in the preceding sections. As we have seen in Section 3.4.1, one can describe multi-symmetric spinor-tensors in an explicit oscillator realization contracting their vector indices with commuting vectors $u^{i\mu}$, as in eq. (3.57). One can then adapt the basic operations of eq. (3.58) to the case of multi-form gauge fields replacing the *commuting* $u^{i\mu}$ with *anticommuting* $\theta^{i\mu}$ vectors, so that

$$\begin{aligned}\partial_i &\equiv \partial^\mu \frac{\partial}{\partial \theta^{i\mu}}, & \partial^i &\equiv \partial_\mu \theta^{i\mu}, \\ \gamma_i &\equiv \gamma^\mu \frac{\partial}{\partial \theta^{i\mu}}, & \gamma^i &\equiv -\gamma_\mu \theta^{i\mu},\end{aligned}\tag{5.1}$$

where for convenience we have also flipped the sign in the definition of the γ^i ²⁴. Moreover, it is convenient to adopt the conventions

$$\{\gamma_\mu, \theta^{i\nu}\} = 0, \quad \{\gamma_\mu, \frac{\partial}{\partial \theta^{i\nu}}\} = 0.\tag{5.2}$$

The key ingredients of our construction are the algebraic properties of the operators that we have introduced, and with the definitions (5.1) divergences and gradients become anticommuting objects satisfying

$$\{\partial_i, \partial^j\} = \square \delta_i^j,\tag{5.3}$$

while the γ 's now satisfy the algebra

$$\begin{aligned}[\gamma_i, \gamma_j] &= 2T_{ij}, \\ [\gamma^i, \gamma^j] &= -4\eta^{ij},\end{aligned}\tag{5.4}$$

that defines the *antisymmetric* T_{ij} and η^{ij} operators of the present setting.

Furthermore, multi-form gauge fields ψ have the gauge transformations

$$\delta \psi = \partial^i \epsilon_i,\tag{5.5}$$

that maintain the form of eq. (2.6) but now admit the gauge-for-gauge transformations

$$\delta \epsilon_i = \partial^j \epsilon_{(ij)}, \quad \delta \epsilon_{(ij)} = \partial^k \epsilon_{i(jk)}, \quad \dots\tag{5.6}$$

with parameters that are *symmetric* under the interchange of their family indices. This property reflects the well-known fact that gauge-for-gauge transformations are already present for one-family fields, that correspond to spinorial forms in the antisymmetric basis. Taking into account that

$$\{\partial_i, \partial\} = 0\tag{5.7}$$

and following almost verbatim the reasoning that led Labastida [30] to the $\mathcal{S}$ tensor of eq. (2.5), one can now identify the basic kinetic tensor of the theory,

$$\mathcal{S} = i(\not{\partial}\psi + \partial^i \not{\psi}_i),\tag{5.8}$$

²⁴With the definitions of eq. (5.1), the terms in the Lagrangians that already show up for spin-3/2 fields retain the usual coefficients.

as the gauge invariant completion of $\not\partial \psi$. As in the multi-symmetric setting, the gauge invariance of $\mathcal{S}$ can be attained at the price of suitable constraints on the gauge parameters,

$$\gamma_{[i} \epsilon_{j]} = 0, \quad (5.9)$$

since

$$\delta \mathcal{S} = \frac{i}{2} \partial^i \partial^j \gamma_{[i} \epsilon_{j]}. \quad (5.10)$$

In analogy with the presentation of Section 2.1, unconstrained equations of motion obtain performing the Stueckelberg-like shift

$$\psi \rightarrow \psi - \partial^i \Psi_i \quad \text{with} \quad \delta \Psi_i = \epsilon_i, \quad (5.11)$$

that builds the unconstrained gauge invariant kinetic tensor

$$\mathcal{W} = \mathcal{S} - i \partial^i \partial^j \xi_{ij}(\Psi). \quad (5.12)$$

As in the multi-symmetric case, only the composite compensator fields

$$\xi_{ij}(\Psi) = \frac{1}{2} \gamma_{[i} \Psi_{j]}, \quad (5.13)$$

that are now *antisymmetric*, appear in $\mathcal{W}$, due to the constrained gauge invariance of $\mathcal{S}$.

The other key ingredients of the construction developed in the previous sections were the Bianchi identities satisfied by $\mathcal{S}$, that now become

$$\partial_i \mathcal{S} - \frac{1}{2} \not\partial \gamma_i \mathcal{S} - \frac{1}{2} \partial^j T_{ij} \mathcal{S} - \frac{1}{6} \partial^j \gamma_{ij} \mathcal{S} = -\frac{i}{6} \partial^i \partial^j T_{[ij} \gamma_{k]} \psi, \quad (5.14)$$

where one can recognize the first two members of the family of *symmetric* operators

$$\gamma_{i_1 \dots i_n} \equiv \frac{1}{n!} \gamma_{(i_1} \gamma_{i_2} \dots \gamma_{i_n)}. \quad (5.15)$$

The Bianchi identities (5.14) identify the analogues of the Labastida constraints (2.18) for multi-form fields,

$$T_{[ij} \gamma_{k]} \psi = 0, \quad (5.16)$$

that can be enforced off-shell, from the very beginning, or equivalently only on-shell, via Lagrange multipliers λ_{ijk} , that are now *totally antisymmetric* in their family indices. For brevity, in the following we shall confine ourselves to showing how the details that led to the constrained Lagrangians are modified in the present multi-form setting, since the unconstrained Lagrangians behave similarly.

First of all, one can verify by a direct computation that for multi-forms the constraints (5.16) induce the relations

$$T_{[ij} \gamma_{k]} \mathcal{S} = 0, \quad (5.17)$$

so that the only non vanishing combinations of γ -traces of $\mathcal{S}$ are now *two-row* projected,

$$(\gamma^{[q} \mathcal{S}^{p]})_{k_1 \dots k_q; i_1 j_1; \dots; i_p j_p} \equiv Y_{\{p+q, p\}} T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q} \mathcal{S}. \quad (5.18)$$

This property can be easily proved adapting the arguments of Appendix C. Working in the *antisymmetric basis* for Young tableaux is particularly convenient in this context, while the

relevant consequences of the Bianchi identities are now their maximally symmetric *two-row* projected γ -traces. Starting from

$$[T_{i_1 j_1} \dots T_{i_p j_p}, \partial^l] = - \sum_{n=1}^p \partial_{[i_n} \delta_{j_n]}^l \prod_{r \neq n}^p T_{i_r j_r}, \quad (5.19)$$

$$[\gamma_{k_1 \dots k_q}, \not{\partial}] = -2 \partial_{(k_1} \gamma_{k_2 \dots k_q)}, \quad (5.20)$$

$$[\gamma_{k_1 \dots k_q}, \partial^l] = \not{\partial} \gamma_{(k_1 \dots k_{q-1}} \delta_{k_q)}^l - \partial_{(k_1} \gamma_{k_2 \dots k_{q-1}} \delta_{k_q)}^l, \quad (5.21)$$

and considering the composition rules for the $\gamma_{k_1 \dots k_q}$ operators,

$$\gamma_{k_1 \dots k_q} \gamma_l = \gamma_{k_1 \dots k_q l} + \gamma_{(k_1 \dots k_{q-1}} T_{k_q) l}, \quad (5.22)$$

$$\begin{aligned} \gamma_{k_1 \dots k_q} \gamma_{lm} &= \gamma_{k_1 \dots k_q lm} + (\gamma_{(k_1 \dots k_{q-1}} |l T_{|k_q) m} + \gamma_{(k_1 \dots k_{q-1}} |m T_{|k_q) l}) \\ &\quad - \gamma_{(k_1 \dots k_{q-2}} T_{k_{q-1}} |l T_{|k_q) m}, \end{aligned} \quad (5.23)$$

one can then compute the $\{p+q+1, p\}$ projection of the generic $T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q}$ γ -trace of the Bianchi identities (5.14):

$$\begin{aligned} &(p+q+2) Y_{\{p+q+1, p\}} \partial_l (\gamma^{[q]} \mathcal{S}^{[p]})_{k_1 \dots k_q; i_1 j_1; \dots; i_p j_p} \\ &+ \frac{1}{q+3} Y_{\{p+q+1, p\}} \sum_{n=1}^p \partial_{[i_n} (\gamma^{[q+2]} \mathcal{S}^{[p-1]})_{|j_n]} k_1 \dots k_q l; \dots; i_r \neq n j_r \neq n; \dots \\ &- \not{\partial} (\gamma^{[q+1]} \mathcal{S}^{[p]})_{k_1 \dots k_q l; i_1 j_1; \dots; i_p j_p} - \partial^m (\gamma^{[q]} \mathcal{S}^{[p+1]})_{k_1 \dots k_q; i_1 j_1; \dots; i_p j_p; lm} \\ &- \frac{1}{q+3} \partial^m (\gamma^{[q+2]} \mathcal{S}^{[p]})_{k_1 \dots k_q lm; i_1 j_1; \dots; i_p j_p} = 0. \end{aligned} \quad (5.24)$$

As was the case for the Bianchi identities (5.14), these results maintain the same form as their analogues presented in eq. (3.26), barring some sign differences and the replacement of symmetric sets of family indices with antisymmetric ones and vice versa.

In a similar fashion, the counterpart of the Ansatz (3.7) is

$$\mathcal{L} = \frac{1}{2} \langle \bar{\psi}, \sum_{p,q=0}^N \widehat{k}_{p,q} \eta^{i_1 j_1} \dots \eta^{i_p j_p} \gamma^{k_1 \dots k_q} (\gamma^{[q]} \mathcal{S}^{[p]})_{k_1 \dots k_q; i_1 j_1; \dots; i_p j_p} \rangle + \text{h.c.}, \quad (5.25)$$

where we would like to stress again that only *two-row* projected quantities appear here, to be contrasted with the two-column projected ones entering eq. (3.7). Finally, using

$$[\partial_l, \eta^{i_1 j_1} \dots \eta^{i_p j_p}] = \frac{1}{2} \sum_{n=1}^p \delta_l^{[i_n} \partial^{j_n]} \prod_{r \neq n}^p \eta^{i_r j_r}, \quad (5.26)$$

$$[\partial_l, \gamma^{k_1 \dots k_q}] = \delta_l^{(k_1} \gamma^{k_2 \dots k_q)} \not{\partial} - \delta_l^{(k_1} \gamma^{k_2 \dots k_{q-1}} \partial^{k_q)}, \quad (5.27)$$

and the relation

$$Y_{\{2p+1, 1q-1\}} T_{i_1 j_1} \dots T_{i_p j_p} \bar{\epsilon}_l \gamma_{k_1 \dots k_q} = \frac{1}{q+1} Y_{\{2p+1, 1q-1\}} T_{i_1 j_1} \dots T_{i_p j_p} T_l (k_1 \bar{\epsilon}_{k_2} \gamma_{k_3 \dots k_q}), \quad (5.28)$$

that holds in the constrained theory, one can recognize that, up to the convenient choice of an overall normalization $\widehat{k}_{0,0} = 1$, the constrained Lagrangians (5.25) are gauge invariant only if

$$\widehat{k}_{p,q} = \frac{(-1)^{p+q}}{p! q! (p+q+1)!}. \quad (5.29)$$

Aside from a slight difference in the oscillating signs, that are simpler for multi-forms, the coefficients are thus the same as in eq. (3.28).

It should be clear that, in this presentation of the formalism, the need for constraints is determined solely by the number of columns of the Young tableaux associated to generic representations of the Poincaré group. In fact, the antisymmetry in the family indices of eqs. (5.9) and (5.16) shows that all single-column fields, like the gravitino, are not subject to any constraints, while two-column fields are only subject to constraints on the gauge parameters. This is the case, for instance, for a spin 5/2 field. Furthermore, in this setting the Fang-Fronsdal-Labastida operator (5.8) can be recast in the form

$$\mathcal{S} = i \gamma_k \partial^k \psi, \quad (5.30)$$

and for one-family spinorial forms it is manifestly related to the corresponding curvature via a single γ -trace.

6 ELIMINATION OF HIGHER DERIVATIVES

Different types of Lagrangian descriptions for unconstrained irreducible higher-spin fields were explored in the last years. Still, until recently, they all displayed some unusual features with respect to their lower-spin counterparts: as recalled in the Introduction, they typically involve many auxiliary fields, in numbers that increase linearly with the spin [14], or alternatively they contain non-local terms [15, 18, 19, 21], or finally local descriptions with minimal (and fixed) sets of auxiliary fields contain some higher-derivative terms. This is certainly the case for the constructions of [17, 19, 2], but it is equally true for the unconstrained formulations proposed in the previous sections for mixed-symmetry fermions, as can be seen for instance in eqs. (2.13) and (3.42).

A more conventional solution to the same problem was first obtained in [20] for symmetric bosons and fermions. There it was shown how to complement with suitable constraints the Lagrangians given in [16] for the “triplets” of String Field Theory [7], so as to avoid the propagation of the lower spin components contained in the symmetric gauge potentials. The end result of this analysis is an elegant off-shell extension of the constraint relating triplets and compensators equations that was first discussed in [16]. A detailed analysis of the triplet systems in the “frame-like” formalism of [23] was also recently presented in [29].

An alternative construction, that is more in the spirit of the minimal Lagrangians of [17, 19], was proposed in [21] for symmetric Bose fields, and was then extended to mixed-symmetry Bose fields in [2]. In this section we would like to further extend those results to Fermi fields of mixed symmetry. The logic behind the construction will mirror the procedure that we followed to build the minimal local Lagrangians: as a first step, we shall identify a candidate kinetic tensor that is gauge invariant under unconstrained gauge transformations, but is now devoid of higher derivatives. Several choices are indeed possible, and for each of them the second step will be determined by the corresponding form of the Bianchi identities, and will be aimed at building the gauge invariant completion of a trial action suggested by the analogy with the constrained case. One is then to make sure that the resulting equations of motion propagate only the desired degrees of freedom, taking a closer look at the candidate kinetic tensors, in particular in relation with the structure of their Bianchi identities, in order to identify the proper choices. While the result of [21, 2] was more involved than its minimal counterparts involving higher derivatives,

the clearcut logic behind the construction makes it possible to extend the program to Fermi fields in a systematic way.

We shall describe the procedure and the results in the next two sections. First, in Section 6.1 we discuss in some detail the case of symmetric Fermi fields, for which the general construction is presented here for the first time. Having introduced all basic ideas and problems in a relatively simple setting, in Section 6.2 we then turn to the general solution for Fermi fields of mixed symmetry.

6.1 SYMMETRIC FERMIONS

Let us begin by analyzing the behavior of the Fang-Fronsdal theory when the constraints are not enforced, with the aim of building eventually gauge invariant kinetic tensors that are free from higher derivatives. Indeed, in order to compensate the variation of the Fang-Fronsdal tensor ²⁵

$$\delta \mathcal{S} = -i \partial \partial \not{\phi} \quad (6.1)$$

without introducing higher derivatives, the simplest options would be provided by the following alternative definitions for the compensator fields:

$$\begin{aligned} \sigma_{(1)} : \quad \delta \sigma_{(1)} &= \partial \not{\phi} & \rightarrow & \quad \mathcal{W}_{\sigma_{(1)}} = \mathcal{S} + i \partial \sigma_{(1)}, \\ \sigma_{(2)} : \quad \delta \sigma_{(2)} &= \partial \partial \not{\phi} & \rightarrow & \quad \mathcal{W}_{\sigma_{(2)}} = \mathcal{S} + i \sigma_{(2)}. \end{aligned} \quad (6.2)$$

Actually, in contrast with the choice made in [16, 17, 19], corresponding to the one-family case of eq. (2.10), the fields in (6.2) are *not* pure gauge contributions. On the other hand, this observation does not represent a true obstruction, since in principle one could add a Lagrange multiplier term relating these fields to gradients of the usual compensators in order to reduce them to the status of “effective” Stueckelberg-like fields. A closer look at the corresponding Bianchi identities

$$\begin{aligned} \sigma_{(1)} : \quad \partial \cdot \mathcal{W}_{\sigma_{(1)}} - \frac{1}{2} \partial \mathcal{W}'_{\sigma_{(1)}} - \frac{1}{2} \not{\partial} \mathcal{W}_{\sigma_{(1)}} &= \frac{i}{2} (\partial \partial \psi' + \square \sigma_{(1)} - \not{\partial} \not{\phi}_{(1)} - \partial \partial \sigma'_{(1)}), \\ \sigma_{(2)} : \quad \partial \cdot \mathcal{W}_{\sigma_{(2)}} - \frac{1}{2} \partial \mathcal{W}'_{\sigma_{(2)}} - \frac{1}{2} \not{\partial} \mathcal{W}_{\sigma_{(2)}} &= \frac{i}{2} (\partial \partial \psi' + 2 \partial \cdot \sigma_{(2)} - \not{\partial} \not{\phi}_{(2)} - \partial \sigma'_{(2)}), \end{aligned} \quad (6.3)$$

can clearly exhibit the difficulties potentially entailed by these simple choices. Indeed, if one were to pursue one of the two options in (6.2), starting from trial Lagrangians built from either $\mathcal{W}_{\sigma_{(1)}}$ or $\mathcal{W}_{\sigma_{(2)}}$ the gauge variations would give rise to the remainders in eq. (6.3). In order to compensate them, proceeding along the lines that led to (3.42), one would be tempted to introduce Lagrange multipliers $\lambda_{(1)}$ and $\lambda_{(2)}$, but here they would transform *proportionally to the bare gauge parameter* ϵ , since the double gradient acting on ψ' in these cases cannot be factored out of the other terms. A different manifestation of these difficulties is that, in contrast with what happened in the minimal theory of [17, 19], the multipliers would appear in the equations of motion for $\bar{\psi}$ only via their double divergence, so that they would not be expressible algebraically,

²⁵Consistently with the choice made in [2], here we largely abide to the conventions followed in the rest of the present paper, even though for one-family fields the notation of [15, 16, 17, 19, 21] would allow further simplifications, in particular for what concerns products of mutually commuting tensors, obtained at the minor price of defining some new operators like the double gradient in (6.1). For the sake of brevity, however, we use “primes” to denote traces and we still resort to that notation in eq. (6.6).

on shell, in terms of the gauge potential itself, which would have guaranteed against the danger that they carry additional degrees of freedom.

This approach works nicely for symmetric bosons, since in that case one of the multipliers transforms proportionally to the divergence of the gauge parameter. This makes it possible to replace it with a combination of the gauge potential and the ordinary compensator that possesses the same kind of gauge transformation, as shown in [21]. On the other hand, this alternative is clearly not viable in the present case, since no single combination of the fields ψ and $\sigma_{(i)}$ can be defined that transforms like the *bare* gauge parameter ϵ . In the same spirit of the solution that we gave in [2] for mixed-symmetry bosons, we are thus led to look for more general definitions for the gauge invariant kinetic tensors, where a certain amount of additional technical complications can still be acceptable provided the corresponding Bianchi identities are devoid of the previous difficulties. To this end, mimicking what we did in [2] for bosons, let us reconsider the gauge transformation of ψ , separating in the gauge parameter ϵ its γ -traceless part $\epsilon^{(t)}$ from the rest, letting

$$\begin{aligned}\delta\psi &= \partial\epsilon, \\ \epsilon &= \epsilon^{(t)} + \gamma\epsilon^{(g)},\end{aligned}\tag{6.4}$$

so that

$$\not\epsilon = \gamma \cdot (\gamma\epsilon^{(g)}).\tag{6.5}$$

The solution for $\epsilon^{(g)}$ is then

$$\epsilon^{(g)} = \sum_{n=1}^{\lfloor \frac{s}{2} \rfloor} \rho_n(D, s) \eta^{n-1} \left\{ \not\epsilon^{[n-1]} + \frac{1}{2n} \gamma\epsilon^{[n]} \right\},\tag{6.6}$$

with

$$\rho_n(D, s) = (-1)^{n+1} \prod_{k=1}^n \frac{1}{D + 2(s - k - 1)}.\tag{6.7}$$

Here s is the tensorial rank of ψ and $\epsilon^{[n]}$ denotes the n -th trace of ϵ , while η^n denotes a combination of products of n Minkowski metric tensors defined with unit overall normalization and with the minimal number of terms needed in order to obtain a totally symmetric expression. The corresponding decomposition of the unconstrained variation of ψ ,

$$\delta\psi = \partial\epsilon^{(t)} + \gamma\partial\epsilon^{(g)},\tag{6.8}$$

allows one to express the gauge variation of $\mathcal{S}$ in terms of $\epsilon^{(g)}$ alone,

$$\delta\mathcal{S} = -i(D + 2s - 4)\partial\partial\epsilon^{(g)} + i\gamma\partial\partial\not\epsilon^{(g)}.\tag{6.9}$$

Introducing a compensator $\sigma_{(g)}$ such that

$$\delta\sigma_{(g)} = \partial\epsilon^{(g)},\tag{6.10}$$

one can now define a new unconstrained tensor, $\mathcal{W}_{\sigma_{(g)}}$, as

$$\mathcal{W}_{\sigma_{(g)}} = \mathcal{S} + i(D + 2s - 4)\partial\sigma_{(g)} + \gamma\mathcal{S}(\sigma),\tag{6.11}$$

where $\mathcal{S}(\sigma)$ is the Fang-Fronsdal tensor for $\sigma_{(g)}$,

$$\mathcal{S}(\sigma) = i(\not\partial\sigma_{(g)} - \partial\not\sigma_{(g)}).\tag{6.12}$$

Once more, the somewhat involved structure of $\mathcal{W}_{\sigma_{(g)}}$ can well be justified if it avoids the complications that emerged for the simpler choices of eq. (6.2). In fact, in particular thanks to the presence of the tensor $\mathcal{S}(\sigma)$ in $\mathcal{W}_{\sigma_{(g)}}$, the Bianchi identity satisfied by the latter tensor reduces to

$$\partial \cdot \mathcal{W}_{\sigma_{(g)}} - \frac{1}{2} \partial \mathcal{W}'_{\sigma_{(g)}} - \frac{1}{2} \not{\partial} \mathcal{W}_{\sigma_{(g)}} = \frac{i}{2} \partial \partial \left\{ \psi' + \gamma \not{\sigma}_{(g)}' - (D + 2s - 4) \sigma'_{(g)} \right\}, \quad (6.13)$$

so that its structure makes it manifest that the Lagrange multiplier to be introduced does not carry any additional degrees of freedom.

For this reason, one can now consider the variation of a new trial Lagrangian built out of $\mathcal{W}_{\sigma_{(g)}}$ and eventually one can complete the construction in the usual way. However, in order to better understand the rationale behind the kinetic tensor (6.11), it is useful to trace its origin to the substitution

$$\psi \rightarrow \psi - \gamma \sigma_{(g)}, \quad (6.14)$$

to be performed in the Fang-Fronsdal tensor $\mathcal{S}$.

Eq. (6.14) should be compared to the Stueckelberg shift (2.14), that in the symmetric case would simply read

$$\psi \rightarrow \psi - \partial \Psi, \quad (6.15)$$

and would give rise to the higher-derivative tensor (2.13). In this case, one defines an *identically* gauge invariant combination of fields that, as such, would make *any* tensor built out of it gauge invariant as well. The price to pay, however, is to introduce a compensator field whose dimension is different from that of ψ . This choice is then ultimately responsible for the appearance of higher derivatives after the substitution (6.15) is performed in a Lagrangian that initially involved only ψ and one-derivative terms. On the other hand, eq. (6.14) defines a combination of fields displaying *the same gauge transformation as the Fang-Fronsdal field*. As a result, that substitution would guarantee the *unconstrained* gauge invariance of any expression that is gauge invariant, to begin with, under *constrained* gauge transformations. This property is all one really needs. Moreover, it is manifest from (6.14) that the compensator $\sigma_{(g)}$ has the same dimension as ψ , so that all kinetic operators in the Fang-Fronsdal Lagrangian maintain by construction the same number of derivatives even after the substitution (6.14) is implemented.

Thus, enforcing eq. (6.14) in the constrained Lagrangian and adding a Lagrange multiplier for the triple γ -trace of the field ψ results in a new Lagrangian with only one-derivative terms that is invariant under *unconstrained* gauge transformations. Adding to it a further constraint to guarantee the equivalence between $\sigma_{(g)}$ and a suitable linear combination of γ -traces of the “old” compensator ξ finally leads to

$$\begin{aligned} \mathcal{L} = & \frac{1}{2} \langle \bar{\psi} + \bar{\sigma}_{(g)} \gamma, \mathcal{W}_{\sigma_{(g)}} - \frac{1}{2} \gamma \mathcal{W}_{\sigma_{(g)}} - \frac{1}{2} \eta \mathcal{W}'_{\sigma_{(g)}} \rangle \\ & + \langle \bar{\lambda}, \gamma \cdot (\psi - \gamma \sigma_{(g)})' \rangle + \langle \bar{\chi}, \sigma - \partial \Pi \xi \rangle + h.c., \end{aligned} \quad (6.16)$$

that does not contain any higher-derivative contributions. The projector Π entering this expression, whose explicit form can be obtained from eq. (6.6), is such that $\Pi \delta \xi = \epsilon^{(g)}$, while the full

set of gauge transformations is

$$\begin{aligned}
\delta \psi &= \partial \epsilon^{(t)} + \gamma \partial \epsilon^{(g)}, \\
\delta \sigma_{(g)} &= \partial \epsilon^{(g)}, \\
\delta \xi &= \not{\epsilon} = \gamma \cdot (\gamma \epsilon^{(g)}), \\
\delta \lambda &= \partial \cdot \partial \cdot \epsilon^{(t)}, \\
\delta \chi &= 0.
\end{aligned} \tag{6.17}$$

The consistency of the system can be easily verified considering the equations of motion obtained after removing ξ by a partial gauge fixing. In this case the equation for $\bar{\psi}$ reduces to

$$E_{\bar{\psi}} : \mathcal{S} - \frac{1}{2} \gamma \not{\mathcal{S}} - \frac{1}{2} \eta \mathcal{S}' - \frac{i}{2} \eta \gamma \left(\lambda - \frac{1}{2} \partial \cdot \psi' \right) = 0, \tag{6.18}$$

that can be shown to imply $\lambda = \frac{1}{2} \partial \cdot \psi'$ and $\mathcal{S} = 0$, as usual, while the equations for the multipliers $\bar{\chi}$ and $\bar{\lambda}$, in their turn, set to zero $\sigma_{(g)}$ and $\not{\psi}'$. Finally, the equation for $\bar{\sigma}_{(g)}$ determines the multiplier χ as a function of $\mathcal{S}$, and also guarantees, together with the equation for ξ , the consistency conditions on the divergence of $E_{\bar{\psi}}$, which is to vanish as a consequence of the unconstrained gauge invariance of (6.16).

6.2 MIXED-SYMMETRY FERMIONS

The extension of these results to mixed-symmetry fermions proceeds along the same lines, and we can therefore confine ourselves to illustrating the main steps that lead to the final result. To begin with, in the gauge variation (2.6) of the field ψ ,

$$\delta \psi = \partial^i \epsilon_i, \tag{6.19}$$

one should isolate the contribution to the gauge parameters which is to vanish in the constrained theory. To this end, let us consider the decomposition

$$\epsilon_i = \epsilon^{(t)}_i + \gamma^k \epsilon^{(g)}_{i;k}, \tag{6.20}$$

where

$$\begin{aligned}
\gamma_{(i} \epsilon^{(t)}_{j)} &\equiv 0, \\
\gamma_{(i} \gamma^k \epsilon^{(g)}_{j);k} &\equiv \gamma_{(i} \epsilon_{j)}.
\end{aligned} \tag{6.21}$$

In this fashion, the parameters $\epsilon^{(g)}_{i;k}$ embody precisely the gauge freedom that is absent in the Labastida formulation, and that we would like to allow here without introducing new fields whose dimensions differ from that of ψ . In particular, the variation of the Labastida tensor (2.5) under the decomposition (6.20) reads

$$\delta \mathcal{S} = -\frac{i}{2} \partial^i \partial^j \left(D \delta^k_{(i} + 2 S^k_{(i} \right) \epsilon^{(g)}_{j);k} + \frac{i}{2} \gamma^k \partial^i \partial^j \gamma_{(i} \epsilon^{(g)}_{j);k}. \tag{6.22}$$

Guided by the solution of the problem for symmetric fermions presented in the previous section and from the general solution proposed for bosons in [2], we are thus led to introduce new compensator fields σ_k such that

$$\delta \sigma_k = \partial^l \epsilon^{(g)}_{l;k}. \tag{6.23}$$

This makes it possible to define the fully gauge invariant kinetic tensor

$$\mathcal{W}_\sigma = \mathcal{S} + i \left[(D - 2) \partial^j + 2 \partial^i S^j{}_i \right] \sigma_j + \gamma^k \mathcal{S}_k(\sigma), \quad (6.24)$$

where

$$\mathcal{S}_k(\sigma) = i (\not{\partial} \sigma_k - \partial^j \gamma_j \sigma_k) \quad (6.25)$$

is the Labastida tensor for σ_k , and where, let us reiterate, no higher derivatives appear, precisely because the compensators σ_k have the same physical dimensions as ψ .

The Bianchi identity for $\mathcal{W}_\sigma$,

$$\partial_i \mathcal{W}_\sigma - \frac{1}{2} \not{\partial} \gamma_i \mathcal{W}_\sigma - \frac{1}{2} \partial^j T_{ij} \mathcal{W}_\sigma - \frac{1}{6} \partial^j \gamma_{ij} \mathcal{W}_\sigma = \frac{i}{6} \partial^j \partial^k T_{(ij} \gamma_{k)} (\psi - \gamma^l \sigma_l), \quad (6.26)$$

takes a particularly nice form, so that the introduction of Lagrange multipliers is not expected to be problematic, and actually provides a clue to a more complete understanding of the structure of $\mathcal{W}_\sigma$. Indeed, the compensators σ_k allow the redefinition

$$\psi \rightarrow \psi - \gamma^k \sigma_k \quad (6.27)$$

of the field ψ , that has the virtue of identifying a combination of fields possessing the same gauge transformation as the *constrained* Labastida field under variations involving *unconstrained* gauge parameters. As we already observed, this is to be contrasted with the Stueckelberg shift (2.14). It is then possible to verify that, after the substitution (6.27), the Labastida tensor (2.5) takes the form of eq. (6.24).

One can thus follow a relatively straightforward route to a Lagrangian based on the tensor (6.24), starting from the Lagrangian defined by eq. (3.44), that we display again here for the sake of clarity,

$$\mathcal{L} = \frac{1}{2} \langle \bar{\psi}, \sum_{p,q=0}^N k_{p,q} \eta^p \gamma^q (\gamma^{[q]} \mathcal{S}^{[p]}) \rangle + \frac{i}{12} \langle \bar{\lambda}_{ijk}, T_{(ij} \gamma_{k)} \psi \rangle + \text{h.c.}, \quad (6.28)$$

with

$$k_{p,q} = \frac{(-1)^{p + \frac{q(q+1)}{2}}}{p! q! (p+q+1)!}, \quad (6.29)$$

where the constraints on the symmetrized triple γ -trace of ψ are enforced by Lagrange multipliers. Since (6.28) is gauge invariant under *constrained* gauge transformations, by construction it will become gauge invariant under unconstrained ones if one performs everywhere the substitution (6.27), that leads to

$$\begin{aligned} \mathcal{L} = & \frac{1}{2} \langle \bar{\psi} + \bar{\sigma}_k \gamma^k, \sum_{p,q=0}^N k_{p,q} \eta^p \gamma^q (\gamma^{[q]} \mathcal{W}_\sigma^{[p]}) \rangle \\ & + \frac{i}{12} \langle \bar{\lambda}_{ijk}, T_{(ij} \gamma_{k)} (\psi - \gamma^l \sigma_l) \rangle + \text{h.c.} \end{aligned} \quad (6.30)$$

Finally, in order to guarantee that the new compensators σ_k be effectively pure gauge, eq. (6.30) should be supplemented with further constraints, meant to enforce a linear relation between the

σ_k and the compensators $\xi_{ij}(\Psi)$ defined in (2.11), so that the complete Lagrangian takes the form

$$\begin{aligned} \mathcal{L} = & \frac{1}{2} \langle \bar{\psi} + \bar{\sigma}_k \gamma^k, \sum_{p,q=0}^N k_{p,q} \eta^p \gamma^q (\gamma^{[q]} \mathcal{W}_\sigma^{[p]}) \rangle \\ & + \frac{i}{12} \langle \bar{\lambda}_{ijk}, T_{(ij} \gamma_{k)} (\psi - \gamma^l \sigma_l) \rangle + \langle \bar{\chi}_k, \sigma_k - \partial^l \Pi^{ij}_{kl} \xi_{ij} \rangle + \text{h.c.}, \end{aligned} \quad (6.31)$$

where Π^{ij}_{kl} is the projector allowing to express the solution of eq. (6.20) as

$$\epsilon^{(g)}_{k;l} = \Pi^{ij}_{kl} \gamma_{(i} \epsilon_{j)}, \quad (6.32)$$

and where, summarizing, the transformations leaving eq. (6.31) invariant are

$$\begin{aligned} \delta \psi &= \partial^i \left(\epsilon^{(t)}_i + \gamma^k \epsilon^{(g)}_{i;k} \right), \\ \delta \sigma_k &= \partial^l \epsilon^{(g)}_{l;k}, \\ \delta \xi_{ij} &= \gamma_{(i} \epsilon_{j)}, \\ \delta \chi_k &= 0, \end{aligned} \quad (6.33)$$

with $\delta \lambda_{ijk}$ given in (3.41) after replacing ϵ with $\epsilon^{(t)}$. Our derivation of the Lagrangian (6.31) makes it also manifest that the equations of motion reduce generically to $\mathcal{S} = 0$, since the fields σ_k and ξ_{ij} can be gauged away simultaneously, while the Lagrange multipliers λ_{ijk} and χ_k can be expressed in terms of the physical field ψ by the equations for $\bar{\psi}$ and $\bar{\sigma}_k$, respectively.

We would like to conclude this section with some general comments on the unconstrained Lagrangians without higher-derivative terms that we proposed here and in [21, 2]. Since the compensators σ_k (as well as their bosonic counterparts θ_{ij} of [21, 2]) add *exactly* the amount of gauge freedom that was frozen in the Fronsdaal-Labastida formulation, we believe that the field content of these Lagrangians is the minimal possible one, if one requires that the theory be unconstrained and contain no higher-derivative operators. In addition, in the symmetric case there is still a one-to-one correspondence between the field contents of bosonic and fermionic theories, as was the case for the minimal Lagrangians with higher derivatives. Strictly speaking, this correspondence is not maintained with mixed symmetry, simply because some of the auxiliary fields bear different numbers of family indices, but this construction of single derivative Lagrangians for Fermi fields mirrors nonetheless the corresponding one for Bose fields of [2], which could prove useful when dealing with higher-spin supermultiplets. On the other hand, as for the bosonic fields discussed in [2], the gauge transformations for the σ_k compensators are typically rather involved, and we do not know them in closed form in general, beyond the symmetric case that was described in the previous section. This is clearly a disadvantage with respect to the minimal theory with higher derivatives, that in this sense appears technically simpler, not only in view of its reduced field content. Finally, we would like to stress again that, although they are somewhat unusual, higher derivatives accompany in our minimal Lagrangians fields that can be gauged away algebraically. In this sense, consistently with all the tests that one can perform at the present time [19], they are not expected to represent a real source of difficulties, even in future generalizations of our setting to the non-linear level.

7 CONCLUSIONS

In this paper we have constructed Lagrangian theories for free mixed-symmetry gauge fermions in a Minkowski background. These types of fields are inevitable in all dimensions $D > 5$, and indeed higher-spin fields of mixed symmetry are key ingredients of all massive string spectra. Whereas their dynamics is still poorly understood, it is difficult to escape the feeling that these excitations, together with their bosonic counterparts, should be held ultimately responsible for many of the most spectacular properties of String Theory. And, in a wider sense, that a closer look at their dynamics has a real potential to shed some light on the very meaning of String Theory and on its possible generalizations.

We based our construction on the important results obtained by Labastida in [30] (see also [9, 36, 37, 18, 24, 38, 39] for other relevant works on mixed-symmetry fields), where he did not arrive at a Lagrangian formulation, but identified nonetheless the constraints on gauge fields and gauge parameters that let the field equations propagate the correct degrees of freedom, as we show in Section 3.4.1. Since our main interest lies in unconstrained formulations, our basic kinetic spinor-tensors are gauge invariant completions of (2.5), that was introduced by Labastida in [30]. In their simplest form, they are obtained introducing compensator fields ξ_{ij} that transform proportionally to the constraints (2.9), but in analogy with the bosonic case of [2] the very fact that the constraints are not independent forces one to relate the ξ_{ij} to more fundamental objects, here denoted by Ψ_i , as in (2.11). We could thus generalize the basic dynamical setting of Labastida, defined by eqs. (2.5), (2.6) and (2.9),

$$\begin{aligned}\mathcal{S} &= i(\not{\partial}\psi - \partial^i\psi_i) = 0, \\ \delta\psi &= \partial^i\epsilon_i, \\ \gamma_{(i}\epsilon_{j)} &= 0,\end{aligned}\tag{7.1}$$

to its unconstrained counterpart, summarized by eqs. (2.13), (2.6), (2.11) and (2.12),

$$\begin{aligned}\mathcal{W} &= \mathcal{S} + i\partial^i\partial^j\xi_{ij} = 0, \\ \delta\psi &= \partial^i\epsilon_i, \\ \xi_{ij}(\Psi) &= \frac{1}{2}\gamma_{(i}\Psi_{j)}, \\ \delta\Psi_i &= \epsilon_i,\end{aligned}\tag{7.2}$$

that describes the same physical polarizations.

As in the bosonic construction of [2], our method to build the Lagrangians was largely driven by the Bianchi identities, but for Fermi fields we had to face a number of novel complications. These can be traced, one way or another, to the fact that fermionic constraints and compensator terms are *not* well distinct a priori, so that disentangling the truly independent contributions to the Lagrangians is not straightforward. This led, for two-family fields, to the *one-parameter* family of Lagrangians (2.57), equivalent to one another up to redefinitions of the Lagrange multipliers and whose derivation is spelled out in detail in Section 2.1. The general N -family Lagrangians illustrated in Sections 3.1 and 3.2 were also constructed exploiting the Bianchi identities (3.1), or their unconstrained counterparts (3.32). However, in this case we refrained from displaying the full one-parameter family, and confined our attention to a specific form of the Lagrangians that has the virtue of relative simplicity. Moreover, we also presented a derivation that stresses the self-adjoint nature of the resulting Rarita-Schwinger-like tensors.

Eqs. (3.7) and (3.28) are the complete Lagrangians in the constrained setting, and together with eq. (3.42), that provides the corresponding unconstrained extension, represent the main results of the present work.

We also discussed the reduction of the field equations to the Labastida form $\mathcal{S} = 0$, with particular attention to the sporadic low-dimensional cases where the naive procedure encounters some problems and, at the same time, new Weyl-like symmetries emerge. Just as we did for bosons in [2], for two-family Fermi fields we thus arrived at a classification of these special cases, while for general N -family fields we contented ourselves with a discussion of the general setup, together with the explicit analysis of some classes of examples.

We have analyzed the properties of spinor-tensors of mixed symmetry mostly with reference to a specific type of redundant presentation, via *multi-symmetric reducible* fields of the type $\psi_{\mu_1 \dots \mu_{s_1}, \nu_1 \dots \nu_{s_2}, \dots}$, that is capable of encompassing all possibilities. However, for the sake of completeness we have also described how the theory adapts itself to the case of *irreducible* spinor-tensors. In addition, in order to allow a better comparison with String Theory, we have explained how the formalism can be modified in order to deal with *multi-antisymmetric reducible* spinor-tensors.

While the presence of higher-derivative terms related to the compensators in the unconstrained theory is somehow the price to pay for the simplest possible setting, we also showed in Section 6 how to properly modify the form of $\mathcal{W}$ so as to recover a formulation that is a bit more complicated but has the virtue of involving only one-derivative terms. This can be attained with additional auxiliary fields, whose numbers depend only on the number of index families carried by the original gauge fields but not on their ranks, following [21, 2], albeit with some technical complications that emerge when one tries to define explicitly the corresponding gauge parameters.

The present work substantially completes the construction of “minimal” Lagrangian theories for massless, unconstrained higher-spin fields in a flat background. This program was developed in a series of previous papers exploiting the “metric-like” formalism, in an attempt to keep as close as possible to the conventional presentation for lower-spin theories, while also exhibiting the geometrical interpretation of the free dynamics, that is foreign to the original, constrained formulation of (Fang-)Fronsdal and Labastida. In particular, the works [15, 19, 21] were devoted to characterizing the role played by linearized higher-spin curvatures in the dynamics of free massless or massive higher-spin fields. On the other hand, in [17, 19, 21, 2], and finally in the present paper, we have presented a complete local Lagrangian description for symmetric higher-spin fields in flat and (A)dS backgrounds, and for fields of any symmetry type in flat backgrounds, making use of a minimal amount of additional auxiliary fields that make up for the direct use of higher-spin curvatures. This avoids all non-localities, but also bypasses, at the same time, the need for algebraic constraints of any sort on gauge fields and gauge parameters. In this local setting we mostly focused our attention on massless fields, but in a flat space time the treatment can be directly extended to the massive case, for instance via the harmonics of a Kaluza-Klein circle reduction. The description of (partially) massive higher-spin fields [41] on (A)dS backgrounds is less straightforward. It was discussed, in particular, in the second of [19] for symmetric tensors, while a number of features of the mixed-symmetry case were recently discussed in [39], extending the unfolded equations in Minkowski space time presented in [38]. These papers also shed more light on the flat limit of the (A)dS description, supporting the BMV conjecture of [37].

Other interesting aspects of the free theory can still benefit from further analysis. These

include the quantization of the Lagrangians we are proposing, the construction of supersymmetric higher-spin Lagrangians, possibly even resting on higher-spin analogues of the ordinary superalgebras, the analysis of (partially) massive, mixed symmetry representations on (A)dS backgrounds, and a closer look at the higher-spin geometry for the mixed symmetry, fermionic case. It is clear, however, that the real issue is now to try and reach a comparable understanding of the interactions, although progress in this respect is expected to be far more difficult and slower. Indeed, the radical difficulties that are met in any naive attempts to go beyond the linear regime for higher-spin fields stimulated along the years a number of different approaches to the problem (see [42, 43] for an inevitably incomplete list, including however some recent contributions). The approach initially proposed by Fradkin and Vasiliev and then extensively pursued by Vasiliev himself and by his collaborators [25, 26] (see also [44] for relevant contributions along the same lines) has proved, unquestionably, to be most successful to date, since it has led to non-linear equations for systems of infinitely many massless fully symmetric higher-spin fields. Still, a better understanding along these more conventional lines can definitely spur a better grasp of the systematics of these constructions. All this, in one way or another, supports the definite hope that the “metric form” of the theory of free higher-spin fields attained by now can provide some useful insights both into the nature of higher-spin interactions and, in a wider sense perhaps, into the very nature of String Theory.

Acknowledgments

We are very grateful to X. Bekaert and N. Boulanger for extended discussions at the beginning of this project. We are also grateful to the APC-Paris VII, to the École Polytechnique, to the Chalmers University of Technology, to the Scuola Normale Superiore di Pisa and to the “Galileo Galilei Institute” (GGI) of Florence for the kind hospitality extended to one or more of us at various stages while this work was in progress. Finally, we would like to thank the referee for urging us to extend Section 4 and to add Section 3.4.1. The present research was supported in part by APC-Paris VII, by Scuola Normale Superiore, by INFN, by the CNRS, in particular through the P2I program, by the MIUR-PRIN contract 2007-5ATT78, by the EU contracts MRTN-CT-2004-503369 and MRTN-CT-2004-512194, by the NATO grant PST.CLG.978785, and by the ERC Advanced Investigator Grant no. 226455 “Supersymmetry, Quantum Gravity and Gauge Fields” (SUPERFIELDS).

A NOTATION AND CONVENTIONS

In this paper we use the “mostly plus” convention for the space-time signature and resort to a compact notation, eliminating all space-time indices from tensor relations. The fields of interest are here multi-symmetric spinor-tensors $\psi_{\mu_1 \dots \mu_{s_1}, \nu_1 \dots \nu_{s_2}, \dots}$, whose sets of symmetric indices are referred to as “families”. In Section 5 we also describe how to adapt our results to the case of multi-form spinor-tensors, where the indices belonging to a given set are antisymmetrized, but in all the appendices, for definiteness, we refer explicitly to the case of multi-symmetric spinor-tensors, as in most of the present paper.

The fields $\psi_{\mu_1 \dots \mu_{s_1}, \nu_1 \dots \nu_{s_2}, \dots}$, in the following simply denoted by ψ , are thus fully symmetric under interchanges of pairs of indices belonging to the same set, but have no prescribed symmetry relating different sets, and are thus reducible $gl(D)$ spinor-tensors. As a result, they are perhaps less familiar than Young projected spinor-tensors, but are most convenient for the present discussion and play a natural role in String Theory. For instance, in the five superstring models these reducible spinor-tensors accompany products of bosonic oscillators of the type $\alpha_{-1}^{\mu_1} \dots \alpha_{-1}^{\mu_{s_1}} \alpha_{-2}^{\nu_1} \dots \alpha_{-2}^{\nu_{s_2}} \dots$, that are only symmetric under interchanges of pairs of identical oscillators, while the reducible multi-forms described in Section 5 accompany products of fermionic oscillators. In general, the massive superstring spectra contain spinor-tensors with some symmetric and some antisymmetric index sets, whose theory can be deduced from the results of this paper. Recovering conventional field theories from the present formulation requires suitable projections, and this procedure is discussed in Section 4.

Aside from these basic fields, Lagrangians and field equations involve γ -traces, traces, gradients and divergences of ψ , as well as Minkowski metric tensors related to one (or two) of the previous index sets and γ -matrices. As a result, “family indices” are often needed in order to specify the sets to which some tensor indices belong. As in [2], these family indices are here denoted by small-case Latin letters, and the Einstein convention for summing over pairs of them is used throughout. It actually proves helpful to be slightly more precise: *upper* family indices are thus reserved for operators, like a gradient, which *add* space-time indices, while *lower* family indices are used for operators, like a divergence, which *remove* them. For instance gradients, divergences and traces of a field ψ are denoted concisely by $\partial^i \psi$, $\partial_i \psi$ and $T_{ij} \psi$. This shorthand notation suffices to identify the detailed meaning of these symbols, so that

$$\begin{aligned} \partial^i \psi &\equiv \partial_{(\mu_1^i | \psi \dots, | \mu_2^i \dots \mu_{s_i+1}^i), \dots} , \\ \partial_i \psi &\equiv \partial^\lambda \psi \dots, \lambda \mu_1^i \dots \mu_{s_i-1}^i, \dots , \\ T_{ij} \psi &\equiv \psi \dots, \overset{\lambda}{\mu_1^i \dots \mu_{s_i-1}^i}, \dots, \lambda \mu_1^j \dots \mu_{s_j-1}^j, \dots , \end{aligned} \quad (\text{A.1})$$

where, as in [15, 16, 17, 19, 21, 2], we work with symmetrizations that are *not* of unit strength, but involve nonetheless the minimum possible number of terms, and we use round brackets to denote them. Thus, for instance, the product $T_{(ij} T_{kl)}$ here stands for $T_{ij} T_{kl} + T_{ik} T_{jl} + T_{il} T_{jk}$. Moreover, we use square brackets to denote antisymmetrizations.

In the mixed-symmetry case, one must also introduce mixed metric tensors

$$\eta^{ij} \psi \equiv \frac{1}{2} \sum_{n=1}^{s_i+1} \eta_{\mu_n^i (\mu_1^j | \psi \dots, \mu_1^i \dots \mu_{r \neq n}^i \dots \mu_{s_i+1}^i, \dots, | \mu_2^j \dots \mu_{s_j+1}^j), \dots} . \quad (\text{A.2})$$

Notice that the η^{ij} are rescaled for $i \neq j$, in order that the diagonal terms retain the conventional normalization. Aside from these basic operations, that were already introduced in [2], for the

Fermi fields treated in this paper one must also define some additional operations involving γ -traces,

$$\begin{aligned}\gamma^i \psi &\equiv \gamma(\mu_1^i | \psi \dots, | \mu_2^i \dots \mu_{s_i+1}^i), \dots, \\ \gamma_i \psi &\equiv \gamma^\lambda \psi \dots, \lambda \mu_1^i \dots \mu_{s_i-1}^i, \dots, \\ \emptyset \psi &\equiv \gamma_\lambda \partial^\lambda \psi_{\mu_1^1 \dots \mu_{s_1}^1}, \dots.\end{aligned}\tag{A.3}$$

The space-time γ -matrices introduced here satisfy the usual Clifford algebra

$$\{\gamma^\mu, \gamma^\nu\} = 2\eta^{\mu\nu},\tag{A.4}$$

and moreover

$$(\gamma_0)^\dagger = -\gamma_0, \quad (\gamma_{\mu \neq 0})^\dagger = \gamma_{\mu \neq 0}.\tag{A.5}$$

In the present paper we also use the antisymmetric basis for “family” γ -matrices, introducing

$$\gamma_{i_1 \dots i_n} \equiv \frac{1}{n!} \gamma[i_1 \gamma_{i_2} \dots \gamma_{i_n}],\tag{A.6}$$

and corresponding objects with raised indices. Furthermore, different groups of *symmetrized* indices borne by a given quantity are separated by colons, while a semicolon signals the beginning of a group of *antisymmetrized* indices.

In order to further simplify the combinatorics, as in [2] it proves convenient to introduce the scalar product

$$\langle \bar{\psi}, \chi \rangle \equiv \frac{1}{s_1! \dots s_n!} \bar{\psi}_{\mu_1^1 \dots \mu_{s_1}^1, \dots, \mu_1^n \dots \mu_{s_n}^n} \chi^{\mu_1^1 \dots \mu_{s_1}^1, \dots, \mu_1^n \dots \mu_{s_n}^n} \equiv \frac{1}{s_1! \dots s_n!} \bar{\psi} \chi.\tag{A.7}$$

Inside the brackets it is then possible to integrate by parts and to turn η 's into traces without introducing any s_i -dependent combinatoric factors, since

$$\langle \bar{\psi}, \partial^i \chi \rangle \equiv \frac{1}{s_1! \dots s_n!} \bar{\psi} \partial^i \chi = -\frac{s_i}{s_1! \dots s_n!} (\partial_i \bar{\psi}) \chi \equiv -\langle \partial_i \bar{\psi}, \chi \rangle,\tag{A.8}$$

and, for instance, if $i \neq j$,

$$\langle \bar{\psi}, \eta^{ij} \chi \rangle \equiv \frac{1}{s_1! \dots s_n!} \bar{\psi} \eta^{ij} \chi = \frac{1}{2} \frac{s_i s_j}{s_1! \dots s_n!} (T_{ij} \bar{\psi}) \chi \equiv \frac{1}{2} \langle T_{ij} \bar{\psi}, \chi \rangle,\tag{A.9}$$

where the reader should notice the somewhat unusual factor $\frac{1}{2}$, originating from our choice of normalization in eq. (A.2). In a similar fashion, for γ -matrices

$$\langle \bar{\psi}, \gamma^i \chi \rangle \equiv \frac{1}{s_1! \dots s_n!} \bar{\psi} \gamma^i \chi = \frac{s_i}{s_1! \dots s_n!} (\bar{\psi} \gamma_i) \chi \equiv \langle \bar{\psi} \gamma_i, \chi \rangle.\tag{A.10}$$

However, in the main body of this paper we are ignoring, for simplicity, the overall factors $\prod_{i=1}^N s_i!$, that should accompany the Lagrangians of multi-symmetric spinor-tensors to grant them the conventional normalization.

The scalar products prove quite convenient to derive Lagrangians and field equations for mixed-symmetry fields, but they would be as convenient for symmetric tensors. In this notation the Lagrangian for symmetric, or one-family, spinor-tensors, would simply read

$$\mathcal{L} = \frac{1}{2} \langle \bar{\psi}, \mathcal{W} - \frac{1}{2} \gamma \mathcal{W} - \frac{1}{2} \eta \mathcal{W}' \rangle + \frac{1}{4} \langle \bar{\xi}, \partial \cdot \mathcal{W} \rangle - \frac{1}{8} \langle \bar{\xi}, \partial \cdot \mathcal{W}' \rangle + \frac{1}{4} \langle \bar{\lambda}, \mathcal{Z} \rangle + \text{h.c.},\tag{A.11}$$

to be compared with the corresponding expression of [19],

$$\begin{aligned}\mathcal{L} = & \frac{1}{2} \bar{\psi} \left(\mathcal{W} - \frac{1}{2} \gamma \mathcal{W} - \frac{1}{2} \eta \mathcal{W}' \right) - \frac{3}{4} \binom{s}{3} \bar{\xi} \partial \cdot \mathcal{W}' \\ & + \frac{1}{2} \binom{s}{2} \bar{\xi} \partial \cdot \mathcal{W} + \frac{3}{2} \binom{s}{4} \bar{\lambda} \mathcal{Z} + \text{h.c.},\end{aligned}\tag{A.12}$$

that contains explicit combinatoric factors.

In order to take proper advantage of this compact notation, it is convenient to collect a number of identities that are used recurrently in this type of analysis. As was already pointed out in [2], these have a more complicated structure than their symmetric counterparts of [19], since they involve a genuinely new type of operation. This turns a tensor with s_i indices in the i -th group into others with $s_i + 1$ indices in the i -th group and $s_j - 1$ indices in the j -th group, according to

$$S^i_j \psi \equiv \psi_{\dots, (\mu_1^i \dots \mu_{s_i}^i |, \dots, | \mu_{s_i+1}^i) \mu_1^j \dots \mu_{s_j-1}^j, \dots}.\tag{A.13}$$

The new rules that are needed follow from the algebra of the various operators, and can be also derived from a realization in terms of bosonic oscillators, along the lines of bosonic String Field Theory and of [11, 12, 30]. The subalgebra that does not contain the operators built using γ -matrices reads

$$[\partial_i, \partial^j] = \square \delta_i^j,\tag{A.14}$$

$$[T_{ij}, \partial^k] = \partial_{(i} \delta_{j)}^k,\tag{A.15}$$

$$[\partial_k, \eta^{ij}] = \frac{1}{2} \delta_k^{(i} \partial^{j)},\tag{A.16}$$

$$[T_{ij}, \eta^{kl}] = \frac{D}{2} \delta_i^{(k} \delta_j^{l)} + \frac{1}{2} \left(\delta_i^{(k} S^l)_{j)} + \delta_j^{(k} S^l)_{i)} \right),\tag{A.17}$$

$$[S^i_j, \eta^{kl}] = \delta_j^{(k} \eta^{l)i},\tag{A.18}$$

$$[T_{ij}, S^k_l] = T_{l(i} \delta_{j)}^k,\tag{A.19}$$

$$[S^i_j, \partial^k] = \partial^i \delta_j^k,\tag{A.20}$$

$$[\partial_k, S^i_j] = \delta_k^i \partial_j,\tag{A.21}$$

$$[S^i_j, S^k_l] = \delta_j^k S^i_l - \delta_l^i S^k_j.\tag{A.22}$$

Notice that the commutators of the S^i_j operators, the key novelty of the mixed-symmetry case, build the $gl(N)$ Lie algebra if N index families are present.

The usual anti-commutation rules (A.4) of the Clifford algebra induce the following relations for the γ 's carrying family indices,

$$\{\gamma_i, \gamma_j\} = 2T_{ij},\tag{A.23}$$

$$\{\gamma^i, \gamma^j\} = 4\eta^{ij},\tag{A.24}$$

where the additional factor in the second of these relations originates from our choice of normalization for η^{ij} in eq. (A.2). The reader should notice that the structure of the Clifford algebra is only apparently transferred to the γ 's with family indices, that are actually more complicated because the off-diagonal T_{ij} and η^{ij} do not vanish in general. The difference is already manifest in the anticommutator

$$\{\gamma_i, \gamma^j\} = D \delta_i^j + 2S^j_i,\tag{A.25}$$

that also contains the S^i_j operators. The other relevant basic (anti)commutation relations are

$$[\gamma_i, \partial^j] = \not\partial \delta_i^j, \quad (\text{A.26})$$

$$[\partial_i, \gamma^j] = \not\partial \delta_i^j, \quad (\text{A.27})$$

$$[\gamma_i, \eta^{jk}] = \frac{1}{2} \delta_i^{(j} \gamma^{k)}, \quad (\text{A.28})$$

$$[T_{ij}, \gamma^k] = \gamma_{(i} \delta_{j)}^k, \quad (\text{A.29})$$

$$[S^i_j, \gamma^k] = \gamma^i \delta_j^k, \quad (\text{A.30})$$

$$[\gamma_k, S^i_j] = \delta_k^i \gamma_j, \quad (\text{A.31})$$

$$\{\gamma_i, \not\partial\} = 2\partial_i, \quad (\text{A.32})$$

$$\{\not\partial, \gamma^i\} = 2\partial^i, \quad (\text{A.33})$$

and the previous relations give rise to the basic rules collected in Appendix B.

Finally, in this paper we make extensive use of a number of standard tools related to the symmetric group. These include, in particular, the Young projectors Y , that are used repeatedly to separate irreducible components in family-index space. In the text the various components are often specified by ordered lists of the lengths of the rows for the associated diagrams, so that, for instance, the $\{3, 2\}$ component corresponds to

$$\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \\ \hline \end{array} \quad (\text{A.34})$$

In general the Young projectors Y can be built combining the contributions of different Young tableaux, that can be identified associating integer labels to the tensor indices to be projected and allowing within the given graph all their arrangements such that these integers grow from left to right and from top to bottom. In some cases, however, this simple procedure can fail to produce an orthogonal decomposition, which can still be attained by a further Graham-Schmidt orthogonalization. This difficulty is not present if, for any pair of tableaux, there is at least a couple of indices belonging to a row of the first that lie in the same column of the second, and vice versa. Let us stress that this problem is never to be faced in our constructions, as a result of the particular symmetry properties of our basic objects. Outside Section 5, in this paper we adopt the conventions of [2], and Young tableaux are then defined in the *symmetric* basis, so that the projector corresponding to a tableau containing n boxes takes the form

$$Y_\tau = \frac{\lambda(\tau)}{n!} S A, \quad (\text{A.35})$$

where S and A are the corresponding products of “row symmetrizers” and “column antisymmetrizers”. On the other hand, in Section 5 it is more convenient to work in the *antisymmetric* basis, where the roles of S and A are interchanged. Moreover, $\lambda(\tau)$ is the dimension of the associated representation of the symmetric group, that can be computed, for instance, counting the standard ways of filling the boxes of the corresponding diagram with the numbers $1, 2, \dots, n$ in increasing order from left to right and from top to bottom. We often denote the ratio that appears in eq. (A.35) as

$$\frac{\lambda(\tau)}{n!} = \frac{1}{h(\tau)}, \quad (\text{A.36})$$

where $h(\tau)$ is the “hook length” of the corresponding tableau. Similar techniques are used extensively in Section 4 to identify irreducible $gl(D)$ and Lorentz tensors.

The hook length h also enters the formulae for the dimensions of $gl(D)$ and $o(D)$ (spinorial) representations, that might prove useful to the reader. Thus, the number of components of an irreducible $gl(D)$ *spinor-tensor* with vector indices that are $\{s_1, \dots, s_N\}$ projected is

$$\dim_{gl(D)} [\{s_1, \dots, s_N\}] = \frac{2^{\lfloor \frac{D}{2} \rfloor}}{h} \prod_{i=1}^N \frac{(D + s_i - i)!}{(D - i)!}, \quad (\text{A.37})$$

while for the corresponding $o(D)$ *spinor-tensor* this number is

$$\dim_S [\{s_1, \dots, s_N\}] = \frac{2^{\lfloor \frac{D}{2} \rfloor}}{h} \prod_{i=1}^N \frac{(D + s_i - N - i)!}{(D - 2i)!} \prod_{j=i+1}^N (D + s_i + s_j - i - j + 1), \quad (\text{A.38})$$

to be compared with the number of components for the $o(D)$ *tensor* with the same Young projection in its vector indices:

$$\dim_T [\{s_1, \dots, s_N\}] = \frac{1}{h} \prod_{i=1}^N \frac{(D + s_i - N - i - 1)!}{(D - 2i)!} \prod_{j=i}^N (D + s_i + s_j - i - j). \quad (\text{A.39})$$

These last relations also appear in [12], but for a typo in the last products of eqs. (A.38) and (A.39).

We refrain from adding further details, since these and other related standard facts are discussed extensively in the literature, and in particular in [33].

B USEFUL IDENTITIES FOR FERMIONS

Using repeatedly the (anti)commutation rules of Appendix A yields the useful identities

$$[\partial_l, \eta^{i_1 j_1} \dots \eta^{i_p j_p}] = \frac{1}{2} \sum_{n=1}^p \prod_{r \neq n}^p \eta^{i_r j_r} \delta_l^{(i_n} \partial^{j_n)}, \quad (\text{B.1})$$

$$[T_{i_1 j_1} \dots T_{i_p j_p}, \partial^l] = \sum_{n=1}^p \prod_{r \neq n}^p T_{i_r j_r} \partial_{(i_n} \delta_{j_n)}^l, \quad (\text{B.2})$$

that were already presented in [2], and in a similar fashion

$$[\partial_l, \gamma^{k_1 \dots k_q}] = (-1)^{q+1} \delta_l^{[k_1} \gamma^{k_2 \dots k_q]} \not\partial + \delta_l^{[k_1} \partial^{k_2} \gamma^{k_3 \dots k_q]}, \quad (\text{B.3})$$

$$[\gamma_{k_1 \dots k_q}, \partial^l] = (-1)^{q+1} \not\partial \gamma_{[k_1 \dots k_{q-1}} \delta_{k_q]}^l + \gamma_{[k_1 \dots k_{q-2}} \partial_{k_{q-1}} \delta_{k_q]}^l, \quad (\text{B.4})$$

$$[\gamma_{k_1 \dots k_q}, \not\partial]_{(-1)^{q+1}} = 2 \gamma_{[k_1 \dots k_{q-1}} \partial_{k_q]}, \quad (\text{B.5})$$

$$[\not\partial, \gamma^{k_1 \dots k_q}]_{(-1)^{q+1}} = 2 \partial^{[k_1} \gamma^{k_2 \dots k_q]}, \quad (\text{B.6})$$

where the symbols $[\ , \]_{(-1)^{q+1}}$ can denote both commutators and anticommutators, depending on the sign of $(-1)^{q+1}$. They are widely used in Section 2 and in Section 3 in the construction of the unconstrained Lagrangians. For the reduction of the equation of motion to the Labastida form it is instead convenient to introduce the new operators

$$\gamma^{k_1 \dots k_q}_l = \gamma^{k_1 \dots k_q} \gamma_l - \gamma^{[k_1 \dots k_{q-1}} S^{k_q]}_l, \quad (\text{B.7})$$

that appear in the identities

$$[\gamma_l, \gamma^{k_1 \dots k_q}]_{(-1)^{q+1}} = (D - q + 1) \delta_l^{[k_1} \gamma^{k_2 \dots k_q]} + 2(-1)^{q+1} \gamma^{[k_1 \dots k_{q-1}} S^{k_q]}_l, \quad (\text{B.8})$$

$$[\gamma_{k_1 \dots k_q}, \gamma^l]_{(-1)^{q+1}} = (D - q + 1) \gamma_{[k_1 \dots k_{q-1}} \delta_{k_q]}^l + 2(-1)^{q+1} S^l_{[k_1} \gamma_{k_2 \dots k_q]}, \quad (\text{B.9})$$

$$[T_{lm}, \gamma^{k_1 \dots k_q}] = (-1)^{q+1} \delta_{(l}^{[k_1} \gamma^{k_2 \dots k_q]}_{|m)}, \quad (\text{B.10})$$

$$\begin{aligned} [\gamma_{lm}, \gamma^{k_1 \dots k_q}] &= -(D - q + 1)(D - q + 2) \delta_l^{[k_1} \delta_m^{k_2} \gamma^{k_3 \dots k_q]} \\ &\quad + (-1)^{q+1} (D - q + 1) \delta_{[l}^{[k_1} \gamma^{k_2 \dots k_{q-1}} S^{k_q]}_{|m]} \\ &\quad + (-1)^q (D - q + 2) \delta_{[l}^{[k_1} \gamma^{k_2 \dots k_q]}_{|m]} + 2 \gamma^{[k_1 \dots k_{q-1}} S^{k_q]}_{[l} S^{k_q]}_{|m]}, \end{aligned} \quad (\text{B.11})$$

$$[S^l_m, \gamma^{k_1 \dots k_q}] = (-1)^{q+1} \delta_m^{[k_1} \gamma^{k_2 \dots k_q]}_l, \quad (\text{B.12})$$

$$[\gamma_{k_1 \dots k_q}, S^l_m] = (-1)^{q+1} \gamma_{m[k_1 \dots k_{q-1}} \delta_{k_q]}^l. \quad (\text{B.13})$$

These are used together with the corresponding commutators involving products of metric tensors:

$$[\gamma_l, \eta^{i_1 j_1} \dots \eta^{i_p j_p}] = \frac{1}{2} \sum_{n=1}^p \prod_{r \neq n}^p \eta^{i_r j_r} \delta_l^{(i_n} \gamma^{j_n)}, \quad (\text{B.14})$$

$$\begin{aligned} [T_{lm}, \eta^{i_1 j_1} \dots \eta^{i_p j_p}] &= \frac{1}{2} \sum_{n=1}^p \left\{ D \prod_{r \neq n}^p \eta^{i_r j_r} \delta_l^{(i_n} \delta_m^{j_n)} \right. \\ &\quad \left. + \sum_{m < n} \prod_{r \neq m, n}^p \eta^{i_r j_r} \delta_{(l}^{(i_m} \eta^{j_m)} \delta_{|m)}^{(i_n} \delta^{j_n)}_{|m)} + \prod_{r \neq n}^p \eta^{i_r j_r} \delta_{(l}^{(i_n} S^{j_n)}_{|m)} \right\}, \end{aligned} \quad (\text{B.15})$$

$$\begin{aligned} [\gamma_{lm}, \eta^{i_1 j_1} \dots \eta^{i_p j_p}] &= \frac{1}{4} \sum_{n=1}^p \left\{ \sum_{m < n} \prod_{r \neq m, n}^p \eta^{i_r j_r} \delta_{[l}^{(i_m} \gamma^{j_m)} \delta_{|m]}^{(i_n} \delta^{j_n)}_{|m]} \right. \\ &\quad \left. + 2 \prod_{r \neq n}^p \eta^{i_r j_r} \delta_{[l}^{(i_n} \gamma^{j_n)}_{|m]} \right\}. \end{aligned} \quad (\text{B.16})$$

Finally, we are making a wide use of the following composition rules for family γ -matrices:

$$\gamma_{k_1 \dots k_q} \gamma_l = \gamma_{k_1 \dots k_q l} + \gamma_{[k_1 \dots k_{q-1}} T_{k_q] l}, \quad (\text{B.17})$$

$$\begin{aligned} \gamma_{k_1 \dots k_q} \gamma_{lm} &= \gamma_{k_1 \dots k_q lm} + (\gamma_{[k_1 \dots k_{q-1}} T_{|k_q] l} - \gamma_{[k_1 \dots k_{q-1}} T_{|k_q] m}) \\ &\quad + \gamma_{[k_1 \dots k_{q-2}} T_{k_{q-1} | m} T_{|k_q] l}. \end{aligned} \quad (\text{B.18})$$

Similar rules also apply when one raises indices, but the normalization of η^{ij} introduced in eq. (A.2) requires that this operation be supplemented by the substitution

$$T_{ij} \rightarrow 2\eta^{ij}. \quad (\text{B.19})$$

In order to derive the equations of motion, or in order to impose the condition (3.31) of self-adjointness of the constrained Lagrangians, it is also convenient to keep in mind the further, standard relation

$$\gamma_0 (\gamma_{k_1 \dots k_q})^\dagger \gamma_0 = (-1)^{\frac{(q-1)(q-2)}{2}} \gamma_{k_1 \dots k_q}. \quad (\text{B.20})$$

The previous identities also suffice to compute generic γ -traces of the kinetic tensors $\mathcal{S}$ and $\mathcal{W}$ and of the Bianchi identities. In particular, starting from the Fang-Fronsdal-Labastida tensor

$$\mathcal{S} = i \left(\not{\partial} \psi - \partial^i \psi_i \right), \quad (\text{B.21})$$

one can obtain

$$\begin{aligned} \prod_{r=1}^p T_{i_r j_r} \gamma_{k_1 \dots k_q} \mathcal{S} &= (-1)^q i (q+1) \not{\partial} \gamma_{k_1 \dots k_q} \prod_{r=1}^p T_{i_r j_r} \psi \\ &+ i (q+1) \gamma_{[k_1 \dots k_{q-1}} \partial_{k_q]} \prod_{r=1}^p T_{i_r j_r} \psi - i \sum_{n=1}^p \gamma_{[k_1 \dots k_{q-1}} T_{k_q] (i_n} \partial_{j_n)} \prod_{r \neq n}^p T_{i_r j_r} \psi \\ &- i \sum_{n=1}^p \gamma_{k_1 \dots k_q} (i_n \partial_{j_n}) \prod_{r \neq n}^p T_{i_r j_r} \psi - i \partial^l \gamma_{[k_1 \dots k_{q-1}} T_{k_q] l} \prod_{r=1}^p T_{i_r j_r} \psi \\ &- i \partial^l \gamma_{k_1 \dots k_q l} \prod_{r=1}^p T_{i_r j_r} \psi, \end{aligned} \quad (\text{B.22})$$

and this result is at the root of the derivation, in eq. (3.31), of the constrained Lagrangians via the identification of a constrained self-adjoint operator. It is also needed in a direct derivation of the equations of motion, and in this context the relevant γ -traces for a two-family gauge field read

$$\mathcal{S}_i = i \left(-2 \not{\partial} \psi_i + 2 \partial_i \psi - \partial^j T_{ij} \psi - \partial^j \gamma_{ij} \psi \right), \quad (\text{B.23})$$

$$T_{ij} \mathcal{S} = i \left(\not{\partial} T_{ij} \psi - \partial_{(i} \psi_{j)} - \partial^k T_{ij} \psi_k \right), \quad (\text{B.24})$$

$$\gamma_{ij} \mathcal{S} = i \left(3 \not{\partial} \gamma_{ij} \psi + 3 \gamma_{[i} \partial_{j]} \psi + \partial^k T_{k[i} \gamma_{j]} \psi - \partial^k \gamma_{ijk} \psi \right), \quad (\text{B.25})$$

$$\begin{aligned} T_{ij} \mathcal{S}_k &= i \left((2 T_{ij} \partial_k - T_{k(i} \partial_{j)}) \psi - \gamma_{k(i} \partial_{j)} \psi - 2 \not{\partial} T_{ij} \gamma_k \psi \right. \\ &\quad \left. - \partial^l T_{ij} T_{kl} \psi - \partial^l T_{ij} \gamma_{kl} \psi \right), \end{aligned} \quad (\text{B.26})$$

$$T_{ij} T_{kl} \mathcal{S} = i \left(\not{\partial} T_{ij} T_{kl} \psi - T_{ij} \partial_{(k} \psi_{l)} - T_{kl} \partial_{(i} \psi_{j)} - \partial^m T_{ij} T_{kl} \psi_m \right). \quad (\text{B.27})$$

Other identities that are useful for a direct construction of the field equations in the two-family case are:

$$\mathcal{W}_i = \mathcal{S}_i + 2i \partial^j \not{\partial} \xi_{ij} + i \partial^j \partial^k \gamma_i \xi_{jk}, \quad (\text{B.28})$$

$$T_{ij} \mathcal{W} = T_{ij} \mathcal{S} + 2i \square \xi_{ij} + 2i \partial^k \partial_{(i} \xi_{j)k} + i \partial^k \partial^l T_{ij} \xi_{kl}, \quad (\text{B.29})$$

$$\gamma_{ij} \mathcal{W} = \gamma_{ij} \mathcal{S} - 2i \partial^k \not{\partial} \gamma_{[i} \xi_{j]k} + 2i \partial^k \partial_{[i} \xi_{j]k} + i \partial^k \partial^l \gamma_{ij} \xi_{kl}, \quad (\text{B.30})$$

$$\begin{aligned} T_{ij} \gamma_k \mathcal{W} &= T_{ij} \gamma_k \mathcal{S} + 2i \square \gamma_k \xi_{ij} + 2i \not{\partial} \partial_{(i} \xi_{j)k} + 2i \partial^l \gamma_k \partial_{(i} \xi_{j)l} \\ &\quad + 2i \partial^l \not{\partial} T_{ij} \xi_{kl} + i \partial^l \partial^m T_{ij} \gamma_k \xi_{lm}, \end{aligned} \quad (\text{B.31})$$

$$\begin{aligned}
T_{ij} T_{kl} \mathcal{W} = & T_{ij} T_{kl} \mathcal{S} + 2i \square (T_{ij} \xi_{kl} + T_{kl} \xi_{ij}) + 2i (\partial_i \partial_{(k} \xi_{l)j} + \partial_j \partial_{(k} \xi_{l)i}) \\
& + 2i \partial^m (T_{ij} \partial_{(k} \xi_{l)m} + T_{kl} \partial_{(i} \xi_{j)m}) + i \partial^m \partial^n T_{ij} T_{kl} \xi_{mn}.
\end{aligned} \tag{B.32}$$

C PROOF OF SOME RESULTS USED IN SECTION 3

The construction of the Lagrangians for general Fermi fields of mixed symmetry presented in Sections 3.1 and 3.2 rests upon a number of statements that were referred to without presenting their proofs. In this Appendix we would like to provide the missing details:

- the proof that, for any given number of traces and antisymmetric γ -traces of the kinetic tensor $\mathcal{S}$, all Young projections in family indices with more than two columns vanish in the constrained theory;
- the proof of eq. (3.11), that links terms in the gauge variation of the Lagrangians involving projections with more than two columns to the Labastida constraints on the gauge parameters;
- the generic traces and γ -traces of the Bianchi identities and some of their projections that play an important role in our constructions;
- the actual link between eq. (3.38) and a combination of γ -traces of the Bianchi identities involving only two-column projected γ -traces of the unconstrained kinetic tensor $\mathcal{W}$.

These four points will be discussed in detail in the following four subsections.

YOUNG PROJECTIONS AND CONSTRAINTS

In the symmetric basis that we adopt in most of this paper for the Young tableaux, all projections with more than two columns of a generic combination of traces and antisymmetrized γ -traces can be expressed as a sum of terms where at least three family indices are symmetrized. We can now show that they are all directly related to the Labastida constraints (3.2) if the γ -traces are applied to ψ , or to the constraints (3.4) if the γ -traces are applied to $\mathcal{S}$.

To begin with, one can hide the traces that do not carry any of the symmetrized indices, because they do not play a role in the following analysis. Then, out of the possible placements of these indices the combination

$$T_{i(a} T_{bc)} \gamma_{k_1 \dots k_q} \tag{C.1}$$

is directly related to the constraints. This is so since the cycle can be directly extended to contain four indices, due to the presence of two identical tensors, while the traces commute with $\gamma_{k_1 \dots k_q}$. There is another possibility that involves only traces,

$$T_{i(a|} T_{j|b} T_{c)l} \gamma_{k_1 \dots k_q}. \tag{C.2}$$

It can be treated as in [2], rewriting it in the form

$$T_{i(a} T_{bc} T_{j)l} \gamma_{k_1 \dots k_q} - T_{ij} T_{(ab} T_{c)l} \gamma_{k_1 \dots k_q} - T_{i(a} T_{bc)} T_{j)l} \gamma_{k_1 \dots k_q}, \tag{C.3}$$

so that it vanishes manifestly in the constrained theory, since all these terms can be related to the constraints after extending the symmetrizations as pertains to products of identical T tensors. On the other hand, if one of the symmetrized indices is carried by a (multiple) γ -trace, one can resort to eq. (B.17), obtaining

$$\begin{aligned} T_{(ab\gamma_c)k_1\dots k_{q-1}} &= (-1)^{q+1} \gamma_{k_1\dots k_{q-1}} \gamma_{(a} T_{bc)} + (-1)^q \gamma_{[k_1\dots k_{q-2}} T_{k_{q-1}]} (a T_{bc)} , \\ T_{i(a|T_{j|b}\gamma_c)k_1\dots k_{q-1}} &= (-1)^{q+1} \gamma_{k_1\dots k_{q-1}} \gamma_{(a} T_{b|i} T_{|c)j} \\ &\quad + (-1)^q \gamma_{[k_1\dots k_{q-2}} T_{k_{q-1}]} (a T_{b|i} T_{|c)j} . \end{aligned} \quad (C.4)$$

All these terms therefore vanish either directly or after enlarging the cycle as above.

Alternatively, as in the bosonic case discussed in [2] one can also derive these results noticing that the expression

$$T_{i_1 j_1} \dots T_{i_{p-1} j_{p-1}} T_{(ab\gamma_c)k_1\dots k_{q-1}} , \quad (C.5)$$

admits all Young projections allowed by

$$T_{i_1 j_1} \dots T_{i_{p-1} j_{p-1}} T_{ab\gamma_c k_1\dots k_{q-1}} , \quad (C.6)$$

aside from the two-column one. Furthermore, acting on each irreducible component of (C.5) the permutation group can generate the entire corresponding irreducible subspaces, and in particular the irreducible components of (C.6) that it contains. This suffices to show that all Young projections of the generic expression (C.6) with more than two columns are actually related to the Labastida constraints. In fact, the previous observation shows that they can be expressed as linear combinations of (C.5) with similar quantities obtained permuting indices, and (C.5) is related to the Labastida constraints via the first of eqs. (C.4).

PROOF OF EQ. (3.11)

The presence of the scalar product makes it possible to extract the $\{3, 2^{p-1}\}$ component of

$$\langle T_{i_1 j_1} \dots T_{i_p j_p} \bar{\epsilon}_l \gamma_{k_1\dots k_q} , \partial_l (\gamma^{[q]} \mathcal{S}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1\dots k_q} \rangle \quad (C.7)$$

acting on the left entry with the projector associated to the *single* Young tableau

i_1	j_1	l
$\vdots$	$\vdots$	
i_p	j_p	
k_1		
$\vdots$		
k_q		

(C.8)

In fact, one can decompose $Y_{\{3, 2^{p-1}\}}$ into a sum of Young tableaux where only the projector associated to (C.8) is not annihilated when contracted against the right entry of the scalar

product, where $(\gamma^{[q]} \mathcal{S}^{[p]})$ carries the projection

$$\begin{array}{|c|c|} \hline i_1 & j_1 \\ \hline \vdots & \vdots \\ \hline i_p & j_p \\ \hline k_1 & \\ \hline \vdots & \\ \hline k_q & \\ \hline \end{array} \quad (C.9)$$

since all other tableaux would result in symmetrizations beyond a given a line.

Furthermore, one can recognize that the operations needed to build the projection associated to the tableau (C.8) differ from those implied by the tableau (C.9) only in the symmetrization of the three indices (i_1, j_1, l) . More precisely, denoting the tableau (C.8) by τ_1 and the tableau (C.9) by τ_2 and using the notation of eq. (A.35), when acting on the left entry of eq. (C.7)

$$Y_{\tau_2} = \frac{2}{h(\tau_2)} \tilde{S}_{\tau_2} A_{\tau_2}, \quad (C.10)$$

where the product $\tilde{S}_{\tau_2}$ of “row symmetrizers” does not include the operator $S_{(i_1, j_1)}$, because this symmetrization is already induced by the others. Acting with Y_{τ_1} gives instead

$$Y_{\tau_1} Y_{\tau_2} = \frac{1}{h(\tau_1)} S_{(i_1, j_1, l)} \tilde{S}_{\tau_2} A_{\tau_2} Y_{\tau_2} = \frac{h(\tau_2)}{2h(\tau_1)} S_{(i_1, j_1, l)} (Y_{\tau_2})^2, \quad (C.11)$$

so that, taking into account the ratio between the hook lengths, one ends up with

$$\begin{aligned} & \langle Y_{\{3, 2^{p-1}, 1^q\}} T_{i_1 j_1} \dots T_{i_p j_p} \bar{\epsilon}_l \gamma_{k_1 \dots k_q}, \partial_l (\gamma^{[q]} \mathcal{S}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q} \rangle \\ &= \frac{p(p+q+1)}{(p+1)(p+q+2)} \langle Y_{\tau_2} T_{i_1 j_1} \dots T_{i_p j_p} \bar{\epsilon}_l \gamma_{k_1 \dots k_q}, \partial_{(l|} (\gamma^{[q]} \mathcal{S}^{[p]})_{|i_1 j_1), \dots, i_p j_p; k_1 \dots k_q} \rangle. \end{aligned} \quad (C.12)$$

In a similar discussion presented in Appendix C of [2], these steps sufficed to recover the Labastida constraints on the gauge parameters, due to the presence of only identical T tensors. Here one needs to refine the arguments, because the family indices (i_1, j_1) that are symmetrized with l can be carried both by a trace T and by an antisymmetrized γ , as can be easily seen absorbing the symmetrizations induced by τ_2 in the right entry of the scalar product, so that

$$\begin{aligned} & \langle Y_{\{3, 2^{p-1}, 1^q\}} T_{i_1 j_1} \dots T_{i_p j_p} \bar{\epsilon}_l \gamma_{k_1 \dots k_q}, \partial_l (\gamma^{[q]} \mathcal{S}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q} \rangle \\ &= \langle \frac{2^p (q+1)!}{(p+q+1)!} T_{[i_1 | j_1} \dots T_{|i_p | j_p} \bar{\epsilon}_l \gamma_{|k_1 \dots k_q]}, \\ & \quad \frac{p(p+q+1)}{(p+1)(p+q+2)} \partial_{(l|} (\gamma^{[q]} \mathcal{S}^{[p]})_{|i_1 j_1), \dots, i_p j_p; k_1 \dots k_q} \rangle. \end{aligned} \quad (C.13)$$

In order to proceed, it is convenient to separate the terms in the antisymmetrization that differ in the position of (i_1, j_1) ,

$$\begin{aligned} & T_{[i_1 | j_1} \dots T_{|i_p | j_p} \bar{\epsilon}_l \gamma_{|k_1 \dots k_q]} = T_{i_1 j_1} T_{[i_2 | j_2} \dots T_{|i_p | j_p} \bar{\epsilon}_l \gamma_{|k_1 \dots k_q]} \\ & - \sum_{n=2}^p (-1)^{np} T_{i_1 j_n} T_{j_1 [i_2} T_{i_3 | j_{n+1}} \dots T_{|i_p | j_{n-1}} \bar{\epsilon}_l \gamma_{|k_1 \dots k_q]} \\ & - T_{[i_2 | j_2} \dots T_{|i_p | j_p} T_{|k_1 | j_1} \bar{\epsilon}_l \gamma_{|i_1 | k_1 \dots k_q]}. \end{aligned} \quad (C.14)$$

Forcing again the symmetrization in (i_1, j_1, l) , in the scalar product the second group of terms in eq. (C.14) becomes proportional to the first term, because

$$\begin{aligned}
& \sum_{n=2}^p (-1)^{np} T_{[i_2|j_n \dots T_{|i_{p-1}|j_{n-2} \dots T_{|i_p|(i_1 T_{j_1|j_{n-1}} \bar{\epsilon}|l) \gamma|k_1 \dots k_q]} \\
&= \sum_{n=2}^p (-1)^{np} T_{[i_2|j_n \dots T_{|i_{p-1}|j_{n-2} \dots T_{|i_p|(i_1 T_{j_1 l} \bar{\epsilon}_{j_{n-1}}) \gamma|k_1 \dots k_q]} \\
&- \sum_{n=2}^p (-1)^{np} T_{(i_1 j_1 T_l)[i_2 T_{i_3|j_{n+1} \dots T_{|i_p|j_{n-1}} \bar{\epsilon}_{j_n} \gamma|k_1 \dots k_q]} \\
&- (p-1) T_{[i_2|j_n \dots T_{|i_p|j_{n-1}} T_{(i_1 j_1 \bar{\epsilon} l) \gamma|k_1 \dots k_q]},
\end{aligned} \tag{C.15}$$

and all antisymmetrizations over four indices (including those induced on a couple of traces by a symmetrization over three indices) are annihilated by the right entry of the scalar product (C.13), so that only the last term survives.

In conclusion, the left entry of eq. (C.13) gives rise to two terms when one forces the symmetrization in (i_1, j_1, l) ,

$$\begin{aligned}
T_{[i_1|j_1 \dots T_{|i_p|j_p} \bar{\epsilon}_l \gamma|k_1 \dots k_q]} \longrightarrow \frac{1}{6} \Big\{ (p+1) T_{[i_2|j_n \dots T_{|i_p|j_{n-1}} T_{(i_1 j_1 \bar{\epsilon} l) \gamma|k_1 \dots k_q]} \\
- T_{[i_2|j_n \dots T_{|i_p|j_{n-1}} T_{|k_1|(i_1 \bar{\epsilon}_{j_1} \gamma l)|k_1 \dots k_q]} \Big\},
\end{aligned} \tag{C.16}$$

and imposing the remaining symmetrizations one can recover in each of them the projection associated to the Young tableau

i_2	j_2
$\vdots$	$\vdots$
i_p	j_p
k_1	
$\vdots$	
k_q	

(C.17)

However, this projection is automatically enforced by the right entry of the scalar product, so that

$$\begin{aligned}
& \langle Y_{\{3, 2^{p-1}, 1^q\}} T_{i_1 j_1} \dots T_{i_p j_p} \bar{\epsilon}_l \gamma_{k_1 \dots k_q}, \partial_l (\gamma^{[q]} \mathcal{S}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q} \rangle \\
&= \frac{p(p+q+1)}{(p+1)(p+q+2)} \langle \frac{1}{3(p+q+1)} Y_{\tau_3} \Big\{ (p+1) T_{[i_1|j_1 \dots T_{|i_p|j_p} \bar{\epsilon}_l \gamma|k_1 \dots k_q]} \\
&- T_{[i_2|j_n \dots T_{|i_p|j_{n-1}} T_{|k_1|(i_1 \bar{\epsilon}_{j_1} \gamma l)|k_1 \dots k_q]} \Big\}, \partial_{(l} (\gamma^{[q]} \mathcal{S}^{[p]})_{|i_1 j_1), \dots, i_p j_p; k_1 \dots k_q} \rangle \\
&= \langle (p+1) T_{i_2 j_2} \dots T_{i_p j_p} T_{(i_1 j_1 \bar{\epsilon} l) \gamma_{k_1 \dots k_q}} - T_{i_2 j_2} \dots T_{i_p j_p} T_{[k_1|(i_1 \bar{\epsilon}_{j_1} \gamma l)|k_2 \dots k_q]}, \\
&\frac{p}{3(p+1)(p+q+2)} \partial_{(l} (\gamma^{[q]} \mathcal{S}^{[p]})_{|i_1 j_1), \dots, i_p j_p; k_1 \dots k_q} \rangle,
\end{aligned} \tag{C.18}$$

where τ_3 denotes the tableau of eq. (C.17) and the last two lines lead to the result in eq. (3.11), using eq. (B.17).

BIANCHI IDENTITIES

The generic γ -trace of the constrained Bianchi identities

$$\mathcal{B}_l : \partial_l \mathcal{S} - \frac{1}{2} \not\partial \gamma_l \mathcal{S} - \frac{1}{2} \partial^m T_{lm} \mathcal{S} - \frac{1}{6} \partial^m \gamma_{lm} \mathcal{S} = 0 \quad (\text{C.19})$$

reads

$$\begin{aligned} (-1)^q \prod_{r=1}^p T_{i_r j_r} \gamma_{k_1 \dots k_q} \mathcal{B}_l : & \frac{1}{6} \sum_{n=1}^p \partial_{(i_n \gamma_{j_n) k_1 \dots k_q} l} \prod_{r \neq n}^p T_{i_r j_r} \mathcal{S} \\ & + \partial_{[k_1 \gamma_{k_2 \dots k_q} l]} \prod_{r=1}^p T_{i_r j_r} \mathcal{S} + \frac{q-1}{6} \partial_{[k_1 \gamma_{k_2 \dots k_q} l]} \prod_{r=1}^p T_{i_r j_r} \mathcal{S} \\ & - \frac{(-1)^q}{2} \sum_{n=1}^p \gamma_{k_1 \dots k_q} \partial_{(i_n T_{j_n) l}} \prod_{r \neq n}^p T_{i_r j_r} \mathcal{S} + \frac{1}{6} \sum_{n=1}^p \partial_{(i_n \gamma_{j_n) [k_1 \dots k_{q-1} T_{k_q} l]} \prod_{r \neq n}^p T_{i_r j_r} \mathcal{S} \\ & - \frac{1}{6} \sum_{n=1}^p \gamma_{l [k_1 \dots k_{q-1} T_{k_q} (i_n \partial_{j_n)} \prod_{r \neq n}^p T_{i_r j_r} \mathcal{S} + \frac{q+1}{6} \partial_{[k_1 \gamma_{k_2 \dots k_{q-1} T_{k_q} l]} \prod_{r=1}^p T_{i_r j_r} \mathcal{S} \\ & + \frac{(-1)^q}{6} \sum_{n=1}^p \gamma_{[k_1 \dots k_{q-2} T_{k_{q-1} | l} T_{| k_q} (i_n \partial_{j_n)} \prod_{r \neq n}^p T_{i_r j_r} \mathcal{S} \\ & - \not\partial \left\{ \frac{q+3}{6} \gamma_{k_1 \dots k_q} l + \frac{q-1}{6} \gamma_{[k_1 \dots k_{q-1} T_{k_q} l]} \right\} \prod_{r=1}^p T_{i_r j_r} \mathcal{S} - \frac{(-1)^q}{6} \partial^m \gamma_{k_1 \dots k_q} l m \prod_{r=1}^p T_{i_r j_r} \mathcal{S} \\ & - \frac{(-1)^q}{2} \partial^m \gamma_{k_1 \dots k_q} T_{lm} \prod_{r=1}^p T_{i_r j_r} \mathcal{S} - \frac{1}{6} \partial^m (\gamma_{l [k_1 \dots k_{q-1} T_{k_q} m} - \gamma_{m [k_1 \dots k_{q-1} T_{k_q} l]}) \prod_{r=1}^p T_{i_r j_r} \mathcal{S} \\ & + \frac{(-1)^q}{6} \partial^m \gamma_{[k_1 \dots k_{q-2} T_{k_{q-1} | l} T_{| k_q} m} \prod_{r=1}^p T_{i_r j_r} \mathcal{S} = 0. \end{aligned} \quad (\text{C.20})$$

In order to recover the result in eq. (3.26) for the $\{2^p, 1^{q+1}\}$ projection of this expression, we shall see shortly that the terms containing the same number of traces and an antisymmetrized γ -trace with the same number of indices become proportional when they are $\{2^p, 1^{q+1}\}$ -projected. Furthermore, some of the contributions in eq. (C.20) are annihilated by the Young projector, while the proportionality factors can be fixed resorting to the procedures explained in Appendix C of [2], that we shall also briefly recall in the present context.

First of all, it is convenient to recognize that the $\{2^p, 1^{q+1}\}$ component can be extracted acting with the projector associated to the *single* Young tableau

i_1	j_1
$\vdots$	$\vdots$
i_p	j_p
k_1	
$\vdots$	
k_q	
l	

(C.21)

on account of the symmetry properties of the combination $T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q} \mathcal{B}_l$. As a consequence, all symmetrizations in the generic set of three indices (i_m, j_m, k_n) vanish due to a standard property of Young tableaux. One can now use this fact to relate various terms, so that for instance

$$Y_{\{2^p, 1^{q+1}\}} \sum_{n=1}^p \partial_{(i_n T_{j_n})l} \prod_{r \neq n}^p T_{i_r j_r} \gamma_{k_1 \dots k_q} \mathcal{S} = -p Y_{\{2^p, 1^{q+1}\}} \partial_l T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q} \mathcal{S}. \quad (\text{C.22})$$

To conclude one must also consider that, in order to build the projection associated to (C.21), an antisymmetrization over l and all the k_n indices is required. The $\{2^p, 1^{q+1}\}$ projections of expressions that become proportional after this operation remain proportional with the same overall factor even after the full projection is enforced, and this finally leads to

$$\begin{aligned} & Y_{\{2^p, 1^{q+1}\}} T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q} \mathcal{B}_l : \\ & \left[(q+1) + \frac{q(q-1)}{6} + \frac{p}{2} + \frac{pq}{6} \right] Y_{\{2^p, 1^{q+1}\}} \partial_l T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q} \mathcal{S} \\ & + \frac{1}{6} Y_{\{2^p, 1^{q+1}\}} \sum_{n=1}^p \partial_{(i_n |} \prod_{r \neq n}^p T_{i_r j_r} \gamma_{| j_n) l k_1 \dots k_q} \mathcal{S} \\ & + (-1)^{q+1} \frac{q+3}{6} Y_{\{2^p, 1^{q+1}\}} \not\partial T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q} l \mathcal{S} \\ & - \frac{q+3}{6} Y_{\{2^p, 1^{q+1}\}} \partial^m T_{i_1 j_1} \dots T_{i_p j_p} T_{lm} \gamma_{k_1 \dots k_q} \mathcal{S} \\ & - \frac{1}{6} Y_{\{2^p, 1^{q+1}\}} \partial^m T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q} l m \mathcal{S} = 0. \end{aligned} \quad (\text{C.23})$$

Furthermore, in the two divergence terms only the two-column projected combination of the traces and the antisymmetrized γ -traces can contribute to the full two-column projection also involving the divergence. In a similar fashion, a two-column projection in all indices is enforced in the first gradient term. In fact, the manifest symmetries for its lower family indices are

$$\{2^p, 1^{q+1}\} \otimes \{1\} = \{2^p, 1^{q+2}\} \oplus \{2^{p+1}, 1^q\} \oplus \{3, 2^{p-1}, 1^{q+1}\}, \quad (\text{C.24})$$

but on the other hand this term results from the product of $(p+1)$ traces and a q -fold γ -trace. As a result, the irreducible components with less than four columns that it admits originate from the decomposition

$$\{2^{p+1}\} \otimes \{1^q\} = \{2^{p+1}, 1^q\} \oplus \{3, 2^p, 1^{q-1}\} \oplus \{3^2, 2^{p-1}, 1^{q-2}\} \oplus \dots, \quad (\text{C.25})$$

so that the corresponding term is actually $\{2^{p+1}, 1^q\}$ projected. This is no longer true for the last term of eq. (C.23), where the components displayed in eq. (C.24) and those carried by the product of p traces and a $(q+2)$ -fold γ ,

$$\{2^p\} \otimes \{1^{q+2}\} = \{2^p, 1^{q+2}\} \oplus \{3, 2^{p-1}, 1^{q+1}\} \oplus \{3^2, 2^{p-2}, 1^q\} \oplus \dots, \quad (\text{C.26})$$

share two admissible components. However, in the constrained setting one can elude this problem and simply select the $\{2^p, 1^{q+2}\}$ component, because it is the only non-vanishing one, so that eq. (C.23) directly leads to eq. (3.26), while this detailed analysis is the starting point for the next subsection.

ANALYSIS OF EQ. (3.38)

Most of the results of the previous subsection can be transferred to the unconstrained setting by simply replacing $\mathcal{S}$ with $\mathcal{W}$. Thus, the $\{2^p, 1^{q+1}\}$ projection of the generic γ -trace of the Bianchi identities (3.32),

$$Y_{\{2^p, 1^{q+1}\}} T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q} \mathcal{B}_l, \quad (\text{C.27})$$

reads

$$\begin{aligned} & \frac{(q+3)(p+q+2)}{6} Y_{\{2^p, 1^{q+1}\}} \partial_l (\gamma^{[q]} \mathcal{W}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q} \\ & + \frac{1}{6} Y_{\{2^p, 1^{q+1}\}} \sum_{n=1}^p \partial_{(i_n |} (\gamma^{[q+2]} \mathcal{W}^{[p-1]})_{\dots, i_r \neq n j_r \neq n, \dots; | j_n) l k_1 \dots k_q} \\ & + (-1)^{q+1} \frac{q+3}{6} \partial (\gamma^{[q+1]} \mathcal{W}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q l} \\ & - \frac{q+3}{6} \partial^m (\gamma^{[q]} \mathcal{W}^{[p+1]})_{i_1 j_1, \dots, i_p j_p, lm; k_1 \dots k_q} \\ & - \frac{1}{6} Y_{\{2^p, 1^{q+1}\}} \partial^m T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q lm} \mathcal{W} \\ & = \frac{1}{2} Y_{\{2^p, 1^{q+1}\}} T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q} \partial^m \partial^n \mathcal{Z}_{lmn}. \end{aligned} \quad (\text{C.28})$$

As we just saw, the peculiar feature of this expression is that the $\{2^p, 1^{q+1}\}$ projection in its free indices does not suffice to induce a two-column projection in all indices carried by the last $\mathcal{W}$ term. This is not a problem in the constrained setting, where the other projections vanish on account of the constraints (3.4), but here the last $\mathcal{W}$ term of eq. (C.28) can be only presented in this form, with the lower index m not taking part in the projection. However, an expression containing only two-column projected contributions obtains combining eq. (C.28) with a different product of traces and antisymmetric γ -traces of $\mathcal{B}_l$ carrying the same projection in the family indices, as we can now see. In particular, a $\{2^p, 1^{q+1}\}$ projection is also allowed for the product of $p-1$ traces and a $(q+2)$ -fold γ -trace acting on $\mathcal{B}_l$. In order to maintain the symmetry under interchanges of the (i_n, j_n) couples manifest, it is then natural to consider

$$Y_{\{2^p, 1^{q+1}\}} \sum_{n=1}^p \prod_{r \neq n}^p T_{i_r j_r} \gamma_{k_1 \dots k_q l (i_n \mathcal{B}_{j_n})}, \quad (\text{C.29})$$

that using the techniques of the previous subsection can be cast in the form

$$\begin{aligned} & p(q+1) \left\{ \frac{(q+3)(p+q+2)}{6} Y_{\{2^p, 1^{q+1}\}} \partial_l (\gamma^{[q]} \mathcal{W}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q} \right. \\ & + \frac{1}{6} Y_{\{2^p, 1^{q+1}\}} \sum_{n=1}^p \partial_{(i_n |} (\gamma^{[q+2]} \mathcal{W}^{[p-1]})_{\dots, i_r \neq n j_r \neq n, \dots; | j_n) l k_1 \dots k_q} \\ & + (-1)^{q+1} \frac{q+3}{6} \partial (\gamma^{[q+1]} \mathcal{W}^{[p]})_{i_1 j_1, \dots, i_p j_p; k_1 \dots k_q l} \\ & \left. - \frac{q+3}{6} \partial^m (\gamma^{[q]} \mathcal{W}^{[p+1]})_{i_1 j_1, \dots, i_p j_p, lm; k_1 \dots k_q} \right\} \end{aligned}$$

$$\begin{aligned}
& - \frac{p(q+3)}{6} Y_{\{2^p, 1^{q+1}\}} \partial^m T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q l m} \mathcal{W} \\
& + \frac{(-1)^q}{3} Y_{\{2^p, 1^{q+1}\}} \partial^m \sum_{n=1}^p \prod_{r \neq n}^p T_{i_r j_r} T_{m(i_n \gamma_{j_n}) k_1 \dots k_q l} \mathcal{W} \\
& = \frac{1}{2} Y_{\{2^p, 1^{q+1}\}} \sum_{n=1}^p \prod_{r \neq n}^p T_{i_r j_r} \gamma_{k_1 \dots k_q l (i_n |} \partial^m \partial^n \mathcal{Z}_{|j_n) m n} .
\end{aligned} \tag{C.30}$$

Hence, the combination

$$Y_{\{2^p, 1^{q+1}\}} \left\{ (p+2)(q+3) T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q} \mathcal{B}_l - \sum_{n=1}^p \prod_{r \neq n}^p T_{i_r j_r} \gamma_{k_1 \dots k_q l (i_n \mathcal{B}_{j_n})} \right\} \tag{C.31}$$

in eq. (3.38) contains the following terms with p traces and a $(q+2)$ -fold γ :

$$\begin{aligned}
& - \frac{1}{3} Y_{\{2^p, 1^{q+1}\}} \partial^m \left\{ (q+3) T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q l m} \mathcal{W} \right. \\
& \left. + (-1)^{q+1} \sum_{n=1}^p \prod_{r \neq n}^p T_{i_r j_r} T_{m(i_n \gamma_{j_n}) k_1 \dots k_q l} \mathcal{W} \right\} .
\end{aligned} \tag{C.32}$$

In order to verify that this expression is indeed $\{2^p, 1^{q+2}\}$ projected, one can first recognize that the $\{2^p, 1^{q+1}\}$ projection can be enforced by the projector associated to the single Young tableau displayed in eq. (C.21). Then, using the notation of eq. (A.35) and applying the column antisymmetrizer A associated to (C.21) to the expression within parentheses gives

$$\begin{aligned}
& A_{\{2^p, 1^{q+1}\}} \partial^m \left\{ (q+3) T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q l m} \mathcal{W} + (-1)^{q+1} \sum_{n=1}^p \prod_{r \neq n}^p T_{i_r j_r} T_{m(i_n \gamma_{j_n}) k_1 \dots k_q l} \mathcal{W} \right\} \\
& = p!(q+1)! \partial^m \left\{ (q+3) T_{[i_1 | j_1} \dots T_{|i_p | j_p} \gamma_{|k_1 \dots k_q l] m} \mathcal{W} \right. \\
& + (-1)^{q+1} (q+2) \sum_{n=1}^p T_{[i_1 | j_n} \dots T_{|i_{p-1} | j_{n-2}} T_{m j_{n-1}} \gamma_{|i_p k_1 \dots k_q l]} \mathcal{W} \\
& \left. + (-1)^{q+1} \sum_{n=1}^p T_{[i_1 | j_n} \dots T_{|i_{p-1} | j_{n-2}} T_{m | i_p |} \gamma_{j_{n-1} | k_1 \dots k_q l]} \mathcal{W} \right\} .
\end{aligned} \tag{C.33}$$

When acting with the row symmetrizer S , the last two terms become proportional, since they only differ because of an antisymmetrization over all the i_n indices and a single j_n index, that is annihilated by S . In conclusion, the right-hand side of eq. (C.33) is equivalent to

$$(q+3)p!(q+1)! \partial^m T_{[i_1 | j_1} \dots T_{|i_p | j_p} \gamma_{|k_1 \dots k_q l m]} \mathcal{W} , \tag{C.34}$$

and the remaining symmetrizations induced by S build an expression proportional to that associated to a Young tableau with the index m attached at the end of the longer column of (C.21). Due to the symmetry properties of the underlying combination, this suffices to build the Young projector $Y_{\{2^p, 1^{q+2}\}}$. In order to fix the proportionality factor and to reach the result displayed in eq. (3.38), one can then compare eq. (C.34) with an expression obtained acting with the column antisymmetrizer of the $\{2^p, 1^{q+2}\}$ projector,

$$A_{\{2^p, 1^{q+2}\}} T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q l m} \mathcal{W} = p!(q+2)! T_{[i_1 | j_1} \dots T_{|i_p | j_p} \gamma_{|k_1 \dots k_q l m]} \mathcal{W} . \tag{C.35}$$

Taking into account the hook lengths of the corresponding expressions,

$$h(\{2^p, 1^{q+1}\}) = \frac{p!(p+q+2)!}{q+2}, \quad h(\{2^p, 1^{q+2}\}) = \frac{p!(p+q+3)!}{q+3}, \quad (\text{C.36})$$

finally gives

$$\begin{aligned} & Y_{\{2^p, 1^{q+1}\}} \partial^m \left\{ (q+3) T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q l m} \mathcal{W} + (-1)^{q+1} \sum_{n=1}^p \prod_{r \neq n}^p T_{i_r j_r} T_{m(i_n \gamma_{j_n}) k_1 \dots k_q l} \mathcal{W} \right\} \\ &= (p+q+3) \partial^m Y_{\{2^p, 1^{q+2}\}} T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{k_1 \dots k_q l m} \mathcal{W}, \end{aligned} \quad (\text{C.37})$$

that when added to the other contributions contained in eqs. (C.28) and (C.30) proves the result in eq. (3.38).

D SOME PROPERTIES OF THE REDUCTION FOR N -FAMILY FIELDS

This last appendix collects some results on the reduction to $\mathcal{S} = 0$ of the field equations for N -family spinor-tensors that complement our discussion in Section 3.4.

Let us begin with some comments on the possibility that the tensor $\mathcal{S}$ itself vanish identically. To this end, let us resort again to an oscillator representation for the relevant operators, as in Section 5, that allows to rewrite $\mathcal{S}$ in the form

$$\mathcal{S} = i \partial^\mu \gamma^\nu \left(\eta_{\mu\nu} - u_\mu^i \frac{\partial}{\partial u_\nu^i} \right) \psi, \quad (\text{D.1})$$

where

$$\psi \equiv \frac{1}{s_1! \dots s_N!} u_{\mu_1}^1 \dots u_{\mu_{s_1}}^1 u_{\nu_1}^2 \dots u_{\nu_{s_2}}^2 \dots \psi^{\mu_1 \dots \mu_{s_1}, \nu_1 \dots \nu_{s_2}, \dots}. \quad (\text{D.2})$$

In general, the spinor-tensor $\mathcal{S}$ vanishes identically provided the tensor

$$\left(\eta_{\mu\nu} - u_\mu^i \frac{\partial}{\partial u_\nu^i} \right) \psi \quad (\text{D.3})$$

is γ -traceless. We can now see that the fully antisymmetric fields in $D = N$ mentioned in Section 3.4 are the unique class for which the combination (D.3) is identically zero. While it is relatively easy to verify that in these cases the only available component of $\mathcal{S}$ indeed vanishes, the absence of other solutions can be appreciated in two steps. To begin with, setting the trace of (D.3) to zero yields

$$\left(D - \sum_{i=1}^N s_i \right) \psi = 0, \quad (\text{D.4})$$

which in its turn implies the inequality $D \geq N$. On the other hand, for $\mu \neq \nu$ one ends up with

$$u_\mu^i \frac{\partial}{\partial u_\nu^i} = 0, \quad (\text{D.5})$$

that can be read as orthogonality conditions for the vectors u_μ^i and $\frac{\partial}{\partial u_\nu^i}$. More in detail, in order for (D.5) to hold at fixed μ , it would be necessary to have at least $(D-1)$ orthogonal

directions in the N -dimensional space to which those vectors belong, which can be possible if $D \leq N$. One can therefore conclude that $D = N$, so that all s_i are equal to 1.

Furthermore, in Section 3.4 we proposed the set of conditions (3.83), in order to select the shift transformations

$$\delta \mathcal{S} = \sum_{n=0}^{\lfloor \frac{m}{2} \rfloor} \eta^{i_1 j_1} \dots \eta^{i_n j_n} \gamma^{k_1 \dots k_{m-2n}} \Omega^{[n, m-2n]}_{i_1 j_1, \dots, i_n j_n; k_1 \dots k_{m-2n}} \quad (\text{D.6})$$

that leave eq. (3.77) invariant. Actually, we showed that eqs. (3.83) build these transformations in terms of Weyl-like ψ shifts with parameters Θ such that $\Omega \sim \mathcal{S}(\Theta)$. As a result, they are expected to preserve the Bianchi identities and the (on-shell) Labastida-like constraints induced on $\mathcal{S}$, provided the parameters also satisfy the triple γ -trace constraints (3.85). A missing step was an explicit proof that all transformations of this type, when combined with proper shifts of the $\mathcal{Y}_{ijk}$, are symmetries of the field equations for $\bar{\psi}$, a result that we verified in Section 3.4 only for the simplest shifts with a single γ^i .

While we did not arrive at a complete proof, we can nonetheless display the cancellation mechanism required to this effect for an interesting class of terms. In Sections 2.3 and 3.4 we repeatedly stressed that, when varying the $\mathcal{S}$ -dependent terms in eq. (3.77), all Young projections with more than two columns can be compensated by proper Ω -dependent shifts of the $\mathcal{Y}_{ijk}$ tensors. The rest is of the form

$$\begin{aligned} \delta E_{\bar{\psi}} &= \sum_{p, q=0}^N k_{p, q} \eta^{i_1 j_1} \dots \eta^{i_p j_p} \gamma^{k_1 \dots k_q} \times \\ &\times Y_{\{2^p, 1^q\}} \sum_{n=0}^{\lfloor \frac{m}{2} \rfloor} \sum_{r_1 < \dots < r_n} \prod_{s \neq r_l}^p T_{i_s j_s} \gamma_{[k_1 \dots k_{q-m+2n}]} \theta^{[n, m-2n]}(\Omega)_{i_{r_1} j_{r_1}, \dots, i_{r_n} j_{r_n}, | k_{q-m+2n+1} \dots k_q] }, \end{aligned} \quad (\text{D.7})$$

so that the symmetry is present provided the combinations $\theta^{[n, m-2n]}(\Omega)$ vanish whenever eqs. (3.83) hold. In the following, however, we shall only prove this result for the terms $\theta^{[0, m]}(\Omega)$, or $\theta^{[m]}$ for brevity, that are fully antisymmetric in their family indices and are thus relatively handy to control. In displaying their structure, it is convenient to follow an iterative approach, building $\theta^{[m+1]}$ from $\theta^{[m]}$ via proper redefinitions of the Ω parameters with m family indices that enter the latter quantity. In particular, for even values of m one can let

$$\Omega^{[n, m-2n]}_{i_1 j_1, \dots, i_n j_n; k_1 \dots k_{m-2n}} \rightarrow \gamma^l \Omega^{[n, m-2n+1]}_{i_1 j_1, \dots, i_n j_n; k_1 \dots k_{m-2n} l}, \quad 0 \leq n \leq \frac{m}{2}, \quad (\text{D.8})$$

while for odd values of m one can let

$$\begin{aligned} \Omega^{[n, m-2n]}_{i_1 j_1, \dots, i_n j_n; k_1 \dots k_{m-2n}} &\rightarrow \gamma^l \Omega^{[n, m-2n+1]}_{i_1 j_1, \dots, i_n j_n; k_1 \dots k_{m-2n} l}, \quad 0 \leq n \leq \frac{m-3}{2}, \\ \Omega^{[n, 1]}_{i_1 j_1, \dots, i_n j_n; k} &\rightarrow \gamma^l \left\{ \Omega^{[n, 2]}_{i_1 j_1, \dots, i_n j_n; kl} + \frac{1}{2} \Omega^{[n, 0]}_{i_1 j_1, \dots, i_n j_n, kl} \right\}, \quad n = \frac{m-1}{2}. \end{aligned} \quad (\text{D.9})$$

These redefinitions turn a shift of $\mathcal{S}$ with parameters carrying m family indices into the next member of the set (D.6), with $(m+1)$ family indices. One can therefore use them to determine how the field equations vary under shifts of $\mathcal{S}$ with $(m+1)$ family indices, starting from the redefinition

$$\theta^{[m]}_{k_1 \dots k_m} \rightarrow \gamma^l \tilde{\theta}^{[m]}_{l; k_1 \dots k_m} \quad (\text{D.10})$$

that the procedure induces on $\theta^{[m]}$ and substituting it in eq. (D.7) in order to determine the structure of $\theta^{[m+1]}$. Confining our attention to the $n = 0$ part in (D.7) gives

$$\begin{aligned}
\delta E_{\bar{\psi}} &= \sum_{p,q=0}^N \eta^{i_1 j_1} \dots \eta^{i_p j_p} \gamma^{k_1 \dots k_q} Y_{\{2^p, 1^q\}} \times \\
&\times \left\{ k_{p,q} \left[T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{[k_1 \dots k_{q-m}]} \gamma^l \right]_{(-1)^{q-m+1}} \tilde{\theta}^{[m]}_{l; |k_{q-m+1} \dots k_q]} \right. \\
&+ (-1)^{q+1} \frac{k_{p,q-1}}{q} T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{[k_1 \dots k_{q-m-1}} \tilde{\theta}^{[m]}_{k_{q-m}; k_{q-m+1} \dots k_q]} \\
&+ (-1)^q \frac{q+1}{p} k_{p-1,q+1} \sum_{n=1}^p \prod_{r \neq n}^p T_{i_r j_r} \gamma_{[k_1 \dots k_{q-m+1}} \tilde{\theta}^{[m]}_{(i_n; j_n) | k_{q-m+2} \dots k_q]} \\
&\left. + (-1)^{q+1} \frac{q+1}{p} k_{p-1,q+1} \sum_{n=1}^p \prod_{r \neq n}^p T_{i_r j_r} \gamma_{[k_1 \dots k_{q-m}} (i_n \tilde{\theta}^{[m]}_{j_n); |k_{q-m+1} \dots k_q]} \right\} + \dots,
\end{aligned} \tag{D.11}$$

where we are tracking only terms that can contribute to $\theta^{[m+1]}$. Taking into account the (anti)commutators (3.92) and the coefficients $k_{p,q}$ of eq. (3.28), eq. (D.11) finally becomes

$$\begin{aligned}
\delta E_{\bar{\psi}} &= \sum_{p,q=0}^N k_{p,q} \eta^{i_1 j_1} \dots \eta^{i_p j_p} \gamma^{k_1 \dots k_q} Y_{\{2^p, 1^q\}} \times \\
&\times \left\{ T_{i_1 j_1} \dots T_{i_p j_p} \gamma_{[k_1 \dots k_{q-m-1}} \mathcal{O}(D-m-2)^l_{|k_{q-m}} \tilde{\theta}^{[m]}_{l; |k_{q-m+1} \dots k_q]} \right. \\
&\left. + \sum_{n=1}^p \prod_{r \neq n}^p T_{i_r j_r} \gamma_{[k_1 \dots k_{q-m+1}} \tilde{\theta}^{[m]}_{(i_n; j_n) | k_{q-m+2} \dots k_q]} \right\} + \dots,
\end{aligned} \tag{D.12}$$

where we have introduced the operators

$$\mathcal{O}(\lambda)^l_m = \lambda \delta^l_m + 2 S^l_m. \tag{D.13}$$

Comparing this result with eq. (D.7), one can thus recognize that $\theta^{[m+1]}$ is of the form

$$\theta^{[m+1]}_{k_1 \dots k_{m+1}} \equiv \mathcal{O}(D-m-2)^l_{[k_1} \tilde{\theta}^{[m]}_{l; |k_2 \dots k_{m+1}]}, \tag{D.14}$$

while other contributions in eq. (D.12) are not relevant for the present discussion, but would enter the definition of other θ 's.

The next issue is to exhibit the Ω -dependence of the $\tilde{\theta}^{[m]}$ and $\theta^{[m]}$ spinor-tensors, and the $\mathcal{O}(\lambda)$ operators introduced in eq. (D.13) are particularly convenient to this end. In fact, let us begin by guessing the result,

$$\begin{aligned}
\theta^{[m]}_{k_1 \dots k_m} &= \sum_{n=0}^{\lfloor \frac{m}{2} \rfloor} (-1)^{\frac{(m-n)(m-n-1)}{2}} \prod_{r=1}^n S^{pr}_{|k_r|} \times \\
&\times \prod_{s=1}^{m-2n} \mathcal{O}(D-m-s)^{q_s}_{|k_{n+s}|} \Omega^{[n, m-2n]}_{p_1 | k_{m-n+1} |, \dots, p_n | k_m]; q_1 \dots q_{m-2n}},
\end{aligned} \tag{D.15}$$

where the indices within vertical bars are fully antisymmetrized and the products where the maximum value of the dummy index lies below the lower end are to be regarded as equal to unity.

Our task is now to prove that eq. (D.15) is indeed correct, performing the redefinitions (D.8) or (D.9) and commuting the additional γ -matrix past the other operators acting on the Ω 's. This identifies $\tilde{\theta}^{[m]}$, and multiplying it by a further $\mathcal{O}(D-m-2)$ as in eq. (D.14) should yield eq. (D.15) with $m \rightarrow m+1$. Moving the γ -matrix to the left of all the operators in eq. (D.15) transforms a generic term of the sum into

$$(-1)^{\frac{(m-n)(m-n-1)}{2}} \left(\gamma^l \prod_{r=1}^n S^{p_r}_{|k_r|} + \left[\prod_{r=1}^n S^{p_r}_{|k_r|}, \gamma^l \right] \right) \times \\ \times \prod_{s=1}^{m-2n} \mathcal{O}(D-m-s-2)^{q_s}_{|k_{n+s}|} \Omega^{[n, m-2n+1]}_{p_1 |k_{m-n+1}|, \dots, p_n |k_m|; q_1 \dots q_{m-2n} l}, \quad (\text{D.16})$$

while for odd values of m the last term of the sum becomes

$$\left(\gamma^l \prod_{r=1}^{\frac{m-1}{2}} S^{p_r}_{|k_r|} + \left[\prod_{r=1}^{\frac{m-1}{2}} S^{p_r}_{|k_r|}, \gamma^l \right] \right) \times \\ \times \left\{ \mathcal{O}(D-m-3)^q_{|k_{\frac{m+1}{2}}|} \Omega^{[\frac{m-1}{2}, 2]}_{p_1 |k_{\frac{m+3}{2}}|, \dots, p_{\frac{m-1}{2}} |k_m|; q l} \right. \\ \left. + \frac{1}{2} \mathcal{O}(D-m+1)^q_{|k_{\frac{m+1}{2}}|} \Omega^{[\frac{m+1}{2}, 0]}_{p_1 |k_{\frac{m+3}{2}}|, \dots, p_{\frac{m-1}{2}} |k_m|, q l} \right\}. \quad (\text{D.17})$$

The symmetries of the parameter $\Omega^{[n, m-2n+1]}$ then reduce the commutator in eq. (D.16) to

$$\left[\prod_{r=1}^n S^{p_r}_{|k_r|}, \gamma^l \right] \rightarrow \sum_{r=1}^n \gamma^{p_r} \delta_{|k_r|}^l \prod_{s \neq r}^n S^{p_s}_{|k_s|}, \quad (\text{D.18})$$

since they force $S^{p_s}_{|k_s|}$ and γ^{p_r} to commute.

One should now multiply $\tilde{\theta}^{[m]}$, that is obtained combining eqs. (D.16) and (D.17) and stripping off an overall γ -matrix, by the additional $\mathcal{O}(D-m-2)$ as in eq. (D.14). One can conveniently begin by considering the contributions in (D.16), whose first line thus turns into

$$(-1)^{\frac{(m-n)(m-n-1)}{2}} \left\{ (-1)^n \prod_{r=1}^n S^{p_r}_{|k_r|} \mathcal{O}(D-m-2)^l_{|k_{n+1}|} + 2 \left[S^l_{|k_1|}, \prod_{r=1}^n S^{p_r}_{|k_{r+1}|} \right] \right. \\ \left. + 2 \sum_{r=1}^n S^{p_r}_{|k_1|} \delta_{|k_{r+1}|}^l \prod_{s \neq r}^n S^{p_s}_{|k_s|} \right\}, \quad (\text{D.19})$$

after moving to the right the overall $\mathcal{O}(D-m-2)$, again on account of the symmetries of the Ω parameters. In a similar fashion, the commutator in eq. (D.19) reduces to

$$\left[S^l_{|k_1|}, \prod_{r=1}^n S^{p_r}_{|k_{r+1}|} \right] \rightarrow - \sum_{r=1}^n S^{p_r}_{|k_1|} \delta_{|k_{r+1}|}^l \prod_{s \neq r}^n S^{p_s}_{|k_s|}, \quad (\text{D.20})$$

so that it cancels against the last term in eq. (D.19), that originates from eq. (D.18). Keeping track of an additional factor $(-1)^{m-2n}$ from the reordering of the antisymmetric indices of $\Omega^{[n, m-2n+1]}$ finally leads to eq. (D.15) with $m \rightarrow m+1$, which vindicates our original guess, up to the terms in eq. (D.17), that however behave in the same fashion since

$$\mathcal{O}(\lambda)^l_{|i|} \mathcal{O}(\rho)^q_{|j|} \Omega^{[n, 0]}_{\dots, l q} = 2(\rho - \lambda - 2) S^l_{|i|} \Omega^{[n, 0]}_{\dots, |j| l}, \quad (\text{D.21})$$

as one can prove with arguments similar to those of eq. (2.108).

Having identified the structure of the $\theta^{[m]}$ spinor-tensors, we can finally show that they vanish when eqs. (3.83) hold. To this end, it is convenient to recast the sum (D.15) in the form

$$\begin{aligned} \theta^{[m]}_{k_1 \dots k_m} &= \sum_{n=0}^{\lfloor \frac{m}{2} \rfloor} (-1)^{\frac{(m+n-1)(m+n-2)}{2}} \left(\frac{m-2n}{m} \right) \prod_{r=1}^n S^{p_r}_{|k_r|} \prod_{s=1}^{m-2n-1} \mathcal{O}(D-m-s-1)^{q_s}_{|k_{n+s}|} \\ &\times \left\{ \mathcal{O}(D-m+n-1)^l_{|k_{m-n}|} \Omega^{[n, m-2n]}_{p_1 | k_{m-n+1} |, \dots, p_n | k_m |; l q_1 \dots q_{m-2n-1}} \right. \\ &\left. + \frac{(-1)^n (n+1)}{(m-2n)(m-2n-1)} \Omega^{[n+1, m-2(n+1)]}_{p_1 | k_{m-n+1} |, \dots, p_n | k_{m-1}, k_m |; q_1; q_2 \dots q_{m-2n-1}} \right\}. \end{aligned} \quad (\text{D.22})$$

Before proving that this is indeed possible, let us verify that eq. (D.22) suffices to reach our conclusion. In fact, the Ω -parameters can only contribute to eqs. (D.15) and (D.22) via their unique two-column Young projections in the family indices, since expanding them as in eq. (D.13) and making use of eqs. (3.83) one can effectively replace all the $S^{q_s}_{|k_{n+s}|}$ with various δ 's. As a result, all projections with more than two columns of the resulting terms vanish, simply because it is not possible to antisymmetrize them over $m-n$ indices. On the other hand, the terms within braces in (D.22) vanish if eqs. (3.83) hold, since they are just the antisymmetrization in its (j_r, k) indices of

$$\begin{aligned} &\mathcal{O}(D-m+n-1)^l_k \Omega^{[n, m-2n]}_{i_1 j_1, \dots, i_n j_n; l q_1 \dots q_{m-2n-1}} \\ &+ \frac{(n+1)}{(m-2n)(m-2n-1)} \Omega^{[n+1, m-2(n+1)]}_{i_1 j_1, \dots, i_n j_n, k; q_1; q_2 \dots q_{m-2n-1}} = 0, \end{aligned} \quad (\text{D.23})$$

that collects the contributions to eqs. (3.83) from two-column projected parameters. Let us stress that the $\mathcal{O}$ operators make it possible to deal with these expressions rather efficiently, so that we expect a similar behavior for the generic $\theta^{[n, m-2n]}$ appearing in eq. (D.7)²⁶.

We can now close the present appendix, and hence the whole paper, proving the identities (D.22). First of all, the symmetries of the Ω parameters make it possible to replace any of the $S^{p_r}_{|k_r|}$ operators in eq. (D.15) with $\frac{1}{2} \mathcal{O}(\lambda)^{p_r}_{|k_r|}$, with an arbitrary argument λ . One can thus recast the individual contributions labeled by n in eq. (D.15) in the form

$$\begin{aligned} &(-1)^{\frac{(m-n)(m-n-1)}{2}} \frac{n}{m(m-2n+1)} \prod_{r=1}^{n-1} S^{p_r}_{|k_r|} \prod_{s=1}^{m-2n+1} \mathcal{O}(D-m-s-1)^{q_s}_{|k_{n+s-1}|} \times \\ &\times \Omega^{[n, m-2n]}_{p_1 | k_{m-n+1} |, \dots, p_{n-1} | k_{m-1}, k_m |; q_1; q_2 \dots q_{m-2n+1}} \\ &- \frac{n(-1)^{\frac{(m-n)(m-n-1)}{2}}}{m(m-2n+1)} \prod_{r=1}^{n-1} S^{p_r}_{|k_r|} \left\{ 2 \sum_{s=1}^{m-2n+1} (-1)^{s+1} \prod_{a=1}^{s-1} \mathcal{O}(D-m-a-1)^{q_a}_{|k_{n+a-1}|} \times \right. \\ &\times S^{p_n}_{|k_{n+a}|} \prod_{b=s+1}^{m-2n+1} \mathcal{O}(D-m-b-1)^{q_{b-1}}_{|k_{n+b-1}|} - \frac{m(m-2n+1)}{n} S^{p_n}_{|k_n|} \times \\ &\left. \times \prod_{s=1}^{m-2n} \mathcal{O}(D-m-s)^{q_s}_{|k_{n+s}|} \right\} \Omega^{[n, m-2n]}_{p_1 | k_{m-n+1} |, \dots, p_n | k_m |; q_1 \dots q_{m-2n}}, \end{aligned} \quad (\text{D.24})$$

²⁶The proof that the bosonic field equations are invariant under the shift symmetries of the Bianchi identities, not presented explicitly in [2], also rests on the emergence of products of $\mathcal{O}$ operators, that are actually commuting when the conditions imposed by the Bianchi identities hold.

adding and subtracting the first term, that actually coincides with the second contribution within braces in (D.22) after replacing n with $n - 1$. Moving all “naked” S operators to the left of the $\mathcal{O}$ operators in this expression using

$$\mathcal{O}(\lambda)^{q_s}_{[i]} S^{p_n}_{[j]} \rightarrow -S^{p_n}_{[i]} \mathcal{O}(\lambda + 2)^{q_s}_{[j]}, \quad (\text{D.25})$$

the sum within braces can now be turned into

$$\begin{aligned} S^{p_n}_{[k_n]} & \left\{ 2 \sum_{s=1}^{m-2n+1} \prod_{a=1}^{s-1} \mathcal{O}(D - m - a + 1)^{q_a}_{[k_{n+a}]} \prod_{b=s}^{m-2n} \mathcal{O}(D - m - b - 2)^{q_b}_{[k_{n+b}]} \right. \\ & \left. - \frac{m(m-2n+1)}{n} \prod_{s=1}^{m-2n} \mathcal{O}(D - m - s)^{q_s}_{[k_{n+s}]} \right\} \Omega^{[n, m-2n]}_{p_1 | k_{m-n+1} |, \dots, p_n | k_m |; q_1 \dots q_{m-2n}}. \end{aligned} \quad (\text{D.26})$$

One can finally observe that the $\mathcal{O}$ operators in (D.15) commute,

$$\begin{aligned} & [\mathcal{O}(\lambda)^p_{[i]}, \mathcal{O}(\rho)^q_{[j]}] \Omega^{[n, m-2n]}_{\dots; \dots p \dots q \dots} \\ & = 4 [\mathcal{S}^p_{[i]}, \mathcal{S}^q_{[j]}] \Omega^{[n, m-2n]}_{\dots; \dots p \dots q \dots} = 0, \end{aligned} \quad (\text{D.27})$$

since the antisymmetrization in (i, j) tries to make the commutator symmetric in (p, q) . Furthermore, expanding the $\mathcal{O}$ operators, the resulting terms where only products of δ 's appear suggest that eq. (D.26) can be recast in the form

$$\begin{aligned} & (-1)^{\frac{(m+n-1)(m+n-2)}{2}} \left(\frac{m-2n}{m} \right) \prod_{r=1}^n S^{p_r}_{[k_r]} \prod_{s=1}^{m-2n-1} \mathcal{O}(D - m - s - 1)^{q_s}_{[k_{n+s}]} \\ & \times \mathcal{O}(D - m + n - 1)^l_{[k_{m-n}]} \Omega^{[n, m-2n]}_{p_1 | k_{m-n+1} |, \dots, p_n | k_m |; l q_1 \dots q_{m-2n-1}}, \end{aligned} \quad (\text{D.28})$$

which completes the proof. The factors accompanying these terms are indeed

$$\begin{aligned} & 2 \sum_{r=1}^{m-2n+1} \prod_{a=1}^{r-1} (D - m - a + 1) \prod_{b=r}^{m-2n} (D - m - b - 2) - \frac{m(m-2n+1)}{n} \prod_{a=1}^{m-2n} (D - m - a) \\ & = - \frac{(m-2n+1)(m-2n)}{n} \prod_{a=1}^{m-2n} (D - m - a - 1) (D - m + n - 1), \end{aligned} \quad (\text{D.29})$$

and agree with the structure in eq. (D.28) taking into account that in the resulting product one is to move the l index to the first position, which gives rise to an additional factor $(-1)^{m+1}$. While we have not fully proved eq. (D.28), we have verified that terms of higher order in S work out in a number of cases.

References

- [1] “Higher-Spin Gauge Theories”, Proceedings of the First Solvay Workshop, held in Brussels on May 12-14, 2004, eds. R. Argurio, G. Barnich, G. Bonelli and M. Grigoriev (Int. Solvay Institutes, 2006). This collection contains some contributions closely related, in spirit, to the present work, including:
M. Bianchi and V. Didenko, arXiv:hep-th/0502220; D. Francia and C. M. Hull, arXiv:hep-th/0501236; N. Bouatta, G. Compere and A. Sagnotti, arXiv:hep-th/0409068; X. Bekaert, S. Cnockaert, C. Iazeolla and M. A. Vasiliev, arXiv:hep-th/0503128; A. Sagnotti, E. Sezgin and P. Sundell, arXiv:hep-th/0501156. Other recent reviews include:
D. Sorokin, AIP Conf. Proc. **767** (2005) 172 [arXiv:hep-th/0405069]; A. Fotopoulos and M. Tsulaia, arXiv:0805.1346 [hep-th].
- [2] A. Campoleoni, D. Francia, J. Mourad and A. Sagnotti, Nucl. Phys. B **815** (2009) 289 [arXiv:0810.4350 [hep-th]].
- [3] J. Fang and C. Fronsdal, Phys. Rev. D **18** (1978) 3630.
- [4] L. P. S. Singh and C. R. Hagen Phys. Rev. D **9** (1974) 910.
- [5] T. Curtright, Phys. Lett. B **85** (1979) 219.
- [6] B. de Wit and D. Z. Freedman, Phys. Rev. D **21** (1980) 358; T. Damour and S. Deser, Annales Poincare Phys. Theor. **47** (1987) 277.
- [7] W. Siegel, Nucl. Phys. B **263** (1986) 93; W. Siegel and B. Zwiebach, Nucl. Phys. B **263** (1986) 105; T. Banks and M. E. Peskin, Nucl. Phys. B **264** (1986) 513; M. Kato and K. Ogawa, Nucl. Phys. B **212** (1983) 443; N. Ohta, Phys. Rev. D **33** (1986) 1681, Phys. Lett. B **179** (1986) 347, Phys. Rev. Lett. **56** (1986) 440 [Erratum-ibid. **56** (1986) 1316]; A. Neveu, H. Nicolai and P. C. West, Nucl. Phys. B **264** (1986) 573; A. Neveu and P. C. West, Nucl. Phys. B **268** (1986) 125; E. Witten, Nucl. Phys. B **268** (1986) 253.
- [8] C. S. Aulakh, I. G. Koh and S. Ouvry, Phys. Lett. B **173** (1986) 284; S. Ouvry and J. Stern, Phys. Lett. B **177** (1986) 335; W. Siegel and B. Zwiebach, Nucl. Phys. B **282** (1987) 125.
- [9] W. Siegel, Nucl. Phys. B **284** (1987) 632.
- [10] T. Curtright, Phys. Lett. B **165** (1985) 304.
- [11] J. M. F. Labastida and T. R. Morris, Phys. Lett. B **180** (1986) 101; J. M. F. Labastida, Phys. Rev. Lett. **58** (1987) 531.
- [12] J. M. F. Labastida, Nucl. Phys. B **322** (1989) 185.
- [13] C. Fronsdal, Phys. Rev. D **18** (1978) 3624.
- [14] A. Pashnev and M. Tsulaia, Mod. Phys. Lett. A **12** (1997) 861 [arXiv:hep-th/9703010], Mod. Phys. Lett. A **13** (1998) 1853 [arXiv:hep-th/9803207]; C. Burdik, A. Pashnev and M. Tsulaia, Nucl. Phys. Proc. Suppl. **102** (2001) 285 [arXiv:hep-th/0103143]; I. L. Buchbinder, A. Pashnev and M. Tsulaia, Phys. Lett. B **523** (2001) 338 [arXiv:hep-th/0109067], arXiv:hep-th/0206026; I. L. Buchbinder, V. A. Krykhtin and A. Pashnev, Nucl. Phys. B **711** (2005) 367 [arXiv:hep-th/0410215].

- [15] D. Francia and A. Sagnotti, Phys. Lett. B **543** (2002) 303 [arXiv:hep-th/0207002].
- [16] D. Francia and A. Sagnotti, Class. Quant. Grav. **20** (2003) S473 [arXiv:hep-th/0212185]; A. Sagnotti and M. Tsulaia, Nucl. Phys. B **682** (2004) 83 [arXiv:hep-th/0311257].
- [17] D. Francia and A. Sagnotti, Phys. Lett. B **624** (2005) 93 [arXiv:hep-th/0507144]. For a review see: D. Francia and A. Sagnotti, J. Phys. Conf. Ser. **33** (2006) 57 [arXiv:hep-th/0601199].
- [18] X. Bekaert and N. Boulanger, Commun. Math. Phys. **245** (2004) 27 [arXiv:hep-th/0208058], Phys. Lett. B **561** (2003) 183 [arXiv:hep-th/0301243], Commun. Math. Phys. **271** (2007) 723 [arXiv:hep-th/0606198]; P. de Medeiros and C. Hull, JHEP **0305** (2003) 019 [arXiv:hep-th/0303036].
- [19] D. Francia, J. Mourad and A. Sagnotti, Nucl. Phys. B **773** (2007) 203 [arXiv:hep-th/0701163], Nucl. Phys. B **804** (2008) 383 [arXiv:0803.3832 [hep-th]].
- [20] I. L. Buchbinder, A. V. Galajinsky and V. A. Krykhtin, Nucl. Phys. B **779** (2007) 155 [arXiv:hep-th/0702161].
- [21] D. Francia, Nucl. Phys. B **796** (2008) 77 [arXiv:0710.5378 [hep-th]], Fortsch. Phys. **56** (2008) 800 [arXiv:0804.2857 [hep-th]].
- [22] A. K. H. Bengtsson, Phys. Lett. B **182** (1986) 321; M. Henneaux and C. Teitelboim, in “Quantum Mechanics of Fundamental Systems, 2”, eds. C. Teitelboim and J. Zanelli (Plenum Press, New York, 1988), p. 113.
- [23] M. A. Vasiliev, Yad. Fiz. **32** (1980) 855 [Sov. J. Nucl. Phys. **32** (1980) 439]; V. E. Lopatin and M. A. Vasiliev, Mod. Phys. Lett. A **3** (1988) 257.
- [24] K. B. Alkalaev, O. V. Shaynkman and M. A. Vasiliev, Nucl. Phys. B **692** (2004) 363 [arXiv:hep-th/0311164], JHEP **0508** (2005) 069 [arXiv:hep-th/0501108], arXiv:hep-th/0601225.
- [25] E. S. Fradkin and M. A. Vasiliev, Phys. Lett. B **189** (1987) 89, Nucl. Phys. B **291** (1987) 141; M. A. Vasiliev, Phys. Lett. B **243** (1990) 378, Class. Quant. Grav. **8** (1991) 1387, Phys. Lett. B **285** (1992) 225. For a review, see: M. A. Vasiliev, Int. J. Mod. Phys. D **5** (1996) 763 [arXiv:hep-th/9611024].
- [26] M. A. Vasiliev, Phys. Lett. B **567** (2003) 139 [arXiv:hep-th/0304049].
- [27] C. Becchi, A. Rouet and R. Stora, Commun. Math. Phys. **42** (1975) 127, Annals Phys. **98** (1976) 287; I.V. Tyutin, preprint FIAN n. 39 (1975); E. S. Fradkin and G. A. Vilkovisky, Phys. Lett. B **55** (1975) 224; I. A. Batalin and G. A. Vilkovisky, Phys. Lett. B **69** (1977) 309; M. Henneaux, Phys. Rept. **126** (1985) 1.
- [28] J. Schwinger, “Particles, sources, and fields” (Addison-Wesley, Reading, MA, USA, 1970).
- [29] D. P. Sorokin and M. A. Vasiliev, Nucl. Phys. B **809** (2009) 110 [arXiv:0807.0206 [hep-th]].
- [30] J. M. F. Labastida, Phys. Lett. B **186** (1987) 365.

- [31] L. J. Dixon and J. A. Harvey, Nucl. Phys. B **274** (1986) 93; N. Seiberg and E. Witten, Nucl. Phys. B **276** (1986) 272; M. Bianchi and A. Sagnotti, Phys. Lett. B **247** (1990) 517; A. Sagnotti, arXiv:hep-th/9509080, Nucl. Phys. Proc. Suppl. **56B** (1997) 332 [arXiv:hep-th/9702093]; C. Angelantonj, Phys. Lett. B **444** (1998) 309 [arXiv:hep-th/9810214]; R. Blumenhagen, A. Font and D. Lust, Nucl. Phys. B **558** (1999) 159 [arXiv:hep-th/9904069]. For a review see: C. Angelantonj and A. Sagnotti, Phys. Rept. **371** (2002) 1 [Erratum-ibid. **376** (2003) 339] [arXiv:hep-th/0204089].
- [32] See, for instance: E. Dudas, J. Mourad and A. Sagnotti, Nucl. Phys. B **620**, 109 (2002) [arXiv:hep-th/0107081].
- [33] M. Hamermesh, “Group theory and its applications to physical problems” (Dover Publications, New York, NY, USA, 1969); I. V. Schensted, “A course on the application of group theory to Quantum Mechanics” (NEO Press, Peaks Island, ME, USA, 1976); B.E. Sagan. “The symmetric group”, second edition (Springer-Verlag, New York, NY, USA, 2001). See also: X. Bekaert and N. Boulanger, arXiv:hep-th/0611263.
- [34] Yu. M. Zinoviev, arXiv:0903.0262 [hep-th], arXiv:0904.0549 [hep-th].
- [35] W. Siegel, Phys. Lett. B **93** (1980) 170.
- [36] R. R. Metsaev, Phys. Lett. B **309** (1993) 39, Phys. Lett. B **354** (1995) 78; Class. Quant. Grav. **22** (2005) 2777 [arXiv:hep-th/0412311]; Yu. M. Zinoviev, arXiv:hep-th/0211233, arXiv:hep-th/0304067, arXiv:hep-th/0306292; I. L. Buchbinder, V. A. Krykhtin and H. Takata, Phys. Lett. B **641** (2006) 386 [arXiv:hep-th/0603212], Phys. Lett. B **656** (2007) 253 [arXiv:0707.2181 [hep-th]]; P. Y. Moshin and A. A. Reshetnyak, JHEP **0710** (2007) 040 [arXiv:0707.0386 [hep-th]]; I. L. Buchbinder, V. A. Krykhtin and L. L. Ryskina, arXiv:0902.1471 [hep-th].
- [37] L. Brink, R. R. Metsaev and M. A. Vasiliev, Nucl. Phys. B **586** (2000) 183 [arXiv:hep-th/0005136].
- [38] E. D. Skvortsov, JHEP **0807** (2008) 004 [arXiv:0801.2268 [hep-th]], Nucl. Phys. B **808** (2009) 569 [arXiv:0807.0903 [hep-th]].
- [39] N. Boulanger, C. Iazeolla and P. Sundell, arXiv:0812.3615 [hep-th], arXiv:0812.4438 [hep-th]; E. D. Skvortsov, arXiv:0904.2919 [hep-th].
- [40] S. Weinberg, Phys. Rev. **135** (1964) B1049; S. R. Coleman and J. Mandula, Phys. Rev. **159** (1967) 1251; R. Haag, J. T. Lopuszanski and M. Sohnius, Nucl. Phys. B **88** (1975) 257; C. Aragone and S. Deser, Phys. Lett. B **86** (1979) 161, Nuovo Cim. B **57** (1980) 33; F. A. Berends, J. W. van Holten, B. de Wit and P. van Nieuwenhuizen, J. Phys. A **13** (1980) 1643; P. Benincasa and F. Cachazo, arXiv:0705.4305 [hep-th]; M. Porrati, Phys. Rev. D **78** (2008) 065016 [arXiv:0804.4672 [hep-th]].
- [41] S. Deser and R. I. Nepomechie, Annals Phys. **154** (1984) 396; S. Deser and A. Waldron, Phys. Rev. Lett. **87** (2001) 031601 [arXiv:hep-th/0102166], Nucl. Phys. B **607** (2001) 577 [arXiv:hep-th/0103198], Phys. Lett. B **501**, 134 (2001) [arXiv:hep-th/0012014]; E. D. Skvortsov and M. A. Vasiliev, Nucl. Phys. B **756** (2006) 117 [arXiv:hep-th/0601095].
- [42] S. Weinberg, Phys. Rev. **133** (1964) B1318, Phys. Rev. **134** (1964) B882, Phys. Rev. **181** (1969) 1893; A. K. H. Bengtsson, I. Bengtsson and L. Brink, Nucl. Phys. B **227** (1983) 31,

- Nucl. Phys. B **227** (1983) 41; F. A. Berends, G. J. H. Burgers and H. Van Dam, Z. Phys. C **24** (1984) 247, Nucl. Phys. B **260** (1985) 295; A. K. H. Bengtsson, I. Bengtsson and N. Linden, Class. Quant. Grav. **4** (1987) 1333; S. Deser and Z. Yang, Class. Quant. Grav. **7** (1990) 1491.
- [43] X. Bekaert, N. Boulanger and S. Cnockaert, J. Math. Phys. **46** (2005) 012303 [arXiv:hep-th/0407102], JHEP **0601** (2006) 052 [arXiv:hep-th/0508048]; N. Boulanger, S. Leclercq and S. Cnockaert, Phys. Rev. D **73** (2006) 065019 [arXiv:hep-th/0509118]; X. Bekaert, N. Boulanger, S. Cnockaert and S. Leclercq, Fortsch. Phys. **54** (2006) 282 [arXiv:hep-th/0602092]; N. Boulanger and S. Leclercq, JHEP **0611** (2006) 034 [arXiv:hep-th/0609221]; A. Fotopoulos and M. Tsulaia, Phys. Rev. D **76** (2007) 025014 [arXiv:0705.2939 [hep-th]]; R. R. Metsaev, arXiv:0712.3526 [hep-th]; Yu. M. Zinoviev, Class. Quant. Grav. **26** (2009) 035022 [arXiv:0805.2226 [hep-th]]; N. Boulanger, S. Leclercq and P. Sundell, JHEP **0808** (2008) 056 [arXiv:0805.2764 [hep-th]]; X. Bekaert, E. Joung and J. Mourad, arXiv:0903.3338 [hep-th].
- [44] E. Sezgin and P. Sundell, JHEP **0109** (2001) 036 [arXiv:hep-th/0105001], Nucl. Phys. B **644** (2002) 303 [Erratum-ibid. B **660** (2003) 403] [arXiv:hep-th/0205131], Nucl. Phys. B **762** (2007) 1 [arXiv:hep-th/0508158]; J. Engquist, E. Sezgin and P. Sundell, Class. Quant. Grav. **19** (2002) 6175 [arXiv:hep-th/0207101].